เอกสารอ้างอิง

กนกภรณ์ อ่วมพราหมณ์. (2547). *ความคิดเห็นของมารดาที่มีต่อการพยาบาลในระยะคลอดที่ได้รับจากนักศึกษาพยาบาลศาสตร์บัณฑิต ชั้นปีที่ 3 วิทยาลัยพยาบาลพระจอมเกล้า จังหวัดเพชรบุรี*. (หน้า 6-9), สถาบันพระบรมราชชนก สำนักงานปลัดกระทรวงสาธารณสุข กระทรวงสาธารณสุข.

กมลชนก มากกา. (2552). *ความเจ็บปวดจากการดูดเสมหะของทารกเกิดก่อนกำหนดระหว่างขณะได้รับการห่อตัวกับการจัดท่าในหอผู้ป่วยหนักทารกแรกเกิดโรงพยาบาลพุทธชินราชพิษณุโลก.* พุทธชินราชเวชสาร ฉบับเพิ่มเติม 1, 406.

ขวัญใจ ปิยะวัฒนสกุล. (2546). *ผลของการห่อตัวต่อการตอบสนองต่อความเจ็บปวดจากการดูดเสมหะในท่อหลอดลมคอในทารกคลอดก่อนกำหนด.*วิทยานิพนธ์พยาบาลศาสตรมหาบัณฑิต สาขาวิชาการพยาบาลเด็ก, บัณฑิตวิทยาลัย มหาวิทยาลัยมหิดล.

จรัสศรี หินศิลป์. (2549). *การมีส่วนร่วมของมารดาในการดูแลทารกแรกเกิดที่มีภาวะเสี่ยงสูง*. พยาบาลศาสตรมหาบัณฑิต สาขาวิชาการพยาบาลเด็ก บัณฑิตวิทยาลัย มหาวิทยาลัยเชียงใหม่.

ฐิติพร อุดมกิตติ. (2540). *ผลของการเบี่ยงเบนความสนใจต่อความเจ็บปวดชนิดเฉียบพลันในเด็กวัยทารก.*วิทยานิพนธ์ พยาบาลศาสตรมหาบัณฑิต สาขาวิชาการพยาบาลเด็ก บัณฑิตวิทยาลัย มหาวิทยาลัยมหิดล.

ณัฐธยาน์ อังคประเสริฐกุล.(2551). *ผลของการให้น้ำนมมารดาต่อการตอบสนองความเจ็บปวดจากการเจาะเลือดส้นเท้าในทารกแรกเกิดคลอดครบกำหนด.*วิทยานิพนธ์ปริญญาพยาบาลศาสตรมหาบัณฑิต สาขาวิชาพยาบาลเด็ก บัณฑิตวิทยาลัย มหาวิทยาลัยมหิดล.

นิตยา สินปรุ. (2550). *ผลของการห่อตัวตามแนวปฏิบัติการพยาบาลทางคลินิกเพื่อลดความเจ็บปวดจากการเจาะเลือดบริเวณส้นเท้าในทารกแรกเกิด.* วิทยานิพนธ์พยาบาลศาสตรมหาบัณฑิต สาขาวิชาพยาบาลเด็ก บัณฑิตวิทยาลัย มหาวิทยาลัยมหิดล.

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved

เนตรทอง ทะยา, วิลาวัณย์ พิเชียรเสถียร, และ รัตนชฎาวรรณ อยู่นาค.(2545). *ความต้องการและการได้รับการตอบสนองของมารดาที่เข้ารับการรักษาในหออภิบาลทารกแรกเกิด.*รายงานการวิจัย.เชียงใหม่:คณะพยาบาลศาสตร์ มหาวิทยาลัยเชียงใหม่.

บุบผา แก้ววิเชียร.(2542). *คู่มือการพยาบาลการส่งเสริมการดูแลตนเองในมารดาหลังคลอด งานการพยาบาลผู้ป่วยสูติศาสตร์และนารีเวชวิทยา.* สังกัดฝ่ายการพยาบาลโรงพยาบาลมหาราชนครเชียงใหม่ คณะแพทย์ศาสตร์ มหาวิทยาลัยเชียงใหม่.

ประภาเพ็ญ สุวรรณ.(2526). *ทัศนคติ : การวัดการเปลี่ยนแปลง และพฤติกรรมอนามัย.* พิมพ์ครั้งที่ 2 กรุงเทพมหานคร : โดเดียนสโตร์.

ปัทมา กาคำ. (2540). *ผลการห่อตัวต่อการตอบสนองต่อความเจ็บปวดจากการเจาะส้นเท้าในทารกคลอดครบกำหนด.* วิทยานิพนธ์พยาบาลศาสตรมหาบัณฑิต สาขาวิชาการพยาบาลแม่และเด็ก, บัณฑิตวิทยาลัย มหาวิทยาลัยเชียงใหม่.

พรรณี คำอู. (2547). *ผลของการจัดท่าต่อการตอบสนองความเจ็บปวดจากการเจาะเลือดที่ส้นเท้าในทารกคลอดก่อนกำหนด.*วารสารสภาการพยาบาล,19(1), 70 – 82.

พีรภาพ คำแพง. (2539). *การศึกษาการรับรู้ของพยาบาลเกี่ยวกับความเจ็บปวด ความรู้และกิจกรรมการพยาบาลเพื่อบรรเทาความเจ็บปวดในทารกแรกเกิดที่ได้รับหัตถการ.* วิทยานิพนธ์พยาบาล ศาสตรมหาบัณฑิต สาขาวิชาการพยาบาลแม่และเด็ก, บัณฑิตวิทยาลัย มหาวิทยาลัยมหิดล.

พรพรรณ คีลาวงค์.(2554). *โปรแกรมการเลี้ยงดูทารกสำหรับมารดาหลังคลอดก่อนจำหน่ายจากโรงพยาบาลมหาราชนครเชียงใหม่.* วิทยานิพนธ์ศึกษาศาสตรมหาบัณฑิต สาขาวิชาการส่งเสริมสุขภาพ บัณฑิตวิทยาลัย มหาวิทยาลัยเชียงใหม่.

มนต์ตรา พันธุฟัก. (2552). *ผลของโปรแกรมการส่งเสริมบทบาทการเป็นมารดา ต่อความความรักใคร่ผูกพันระหว่างมารดาและทารก ความพึงพอใจในบทบาทของมารดา ความสามารถในการเรียนรู้พฤติกรรมทารก และการเจริญเติบโตของทารก.*รามาธิบดีพยาบาลสาร ;15(2): 149 – 161.

รัตนาวดี ชอนตะวัน. (2542). *ความรู้และการปฏิบัติกิจกรรมพยาบาลเพื่อลดความเจ็บปวดจากหัตการในทารก.* คณะพยาบาลศาสตร์ มหาวิทยาลัยเชียงใหม่.

ลิวรรณ อุนนาภิรักษ์ และคณะ. (2550). *พยาธิสรีรวิทยาทางการพยาบาล ภาควิชาการพยาบาลรากฐาน.*(หน้า110) คณะพยาบาลศาสตร์ มหาวิทยาลัยมหิดล.

สุชา จันทร์เอม. (2524). *จิตวิทยาทั่วไป.* บริษัทโรงพิมพ์ไทยวัฒนาพานิช จำกัด, กรุงเทพฯ.

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved

สุดาภรณ์ พยัคฆเรือง. (2545).*ผลของการจัดท่านอน ต่อความอิ่มตัวของออกซิเจนในเลือดแดง และความเจ็บปวดในทารก หลังผ่าตัดช่องท้อง*. วิทยานิพนธ์ พยาบาลศาสตรมหาบัณฑิต สาขาวิชาการพยาบาลเด็ก บัณฑิตวิทยาลัย มหาวิทยาลัยมหิดล.

สำนักนโยบายและยุทธศาสตร์. 2551. *"จำนวนและอัตราเกิดมีชีพ ต่อประชาการ 1,000 คนจำแนกรายภาคและเพศ พ.ศ. 2547 – 2551."*[ระบบออนไลน์]. แหล่งที่มา http://bps.ops.moph.go.th (25 มกราคม 2554).

เอื้องดอย ตันฑพงศ์ และคณะ.(2543).*ผลของการห่อตัวต่อการตอบสนองต่อความเจ็บปวดจากการเจาะเลือดบริเวณส้นเท้าในทารกคลอดก่อนกำหนด* วิทยานิพนธ์ปริญญาพยาบาลศาสตร์มหาบัณฑิต,บัณฑิตวิทยาลัย มหาวิทยาลัยมหิดล.

Akman I, Ozek E, Bilgen H, Ozdogan T, Cebeci D.(2002). *Sweet solutions and pacifiers for pain relief in newborn infants.*J Pain. 2002 Jun;3(3),199-202.

Ambuel, B.,Hamlett, K. W., Marx, C. M., & Blumer, J. L. (1992). *Assessing distress in pediatric intensive care environments: The COMFORT scale.* Journal of pediatric Psychology, 17(1),95 – 109.

Anand, K. J. S., & Hicky,P.R.(1987). *Pain and its effects in the human neonate and fetus.* The New England Journal of Medicine, 317(21),1321 – 1329.

Beacham, P.S. (2004). *Behavioral and physiological indicators of procedural and postoperative pain in high – risk infants.* JOGN, 33(2), 246 – 255.

Black, J. M.,& Mattassarin-Jacobs, E. (1993). *Pain assessment and intervention. Luckmann and Sorensen's Medical-surgical nursing*: A psychophysiologic approach (4 ed., pp. 311-358).Philadelphia,PA:W.B.Saunders.

Blass, E. M. (1997). *Milk-induced hypoalagesia in human newborns.* Pediatrics, 99, 825 – 829.

Blass, E. M., & Shan, A. (1995) *Pain – reducing properties of sucrose in human newborns*. Chem Sense, 310, 29 -35.

Campos, R. G. (1989). *Soothing pain-elicited distress in infants with swaddling and pacifiers.* Child Development, 60,781-792.

Carbajal, R., Chauvet, S., Couderc, S., & Olivier-Martin, M. (1999). *Randomized trial of analgesic effects of sucrose, glucose, and pacifiers in term neonates.* British Medical Journal, 38(319), 1393 – 1397.

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved

Carbajal, R., Veerapen, S., & Couderc, S.(2003). *Analgesic effects of breastfeeding in term neonates : randomized trial.* BMJ, 326, 13.

Cleve, L. V.,& Andrews, S. (1995). *Pain response of hospitalized neonates to veinpuncture activities.* American Journal maternal Child Nursing. 20 (May /June),148 -155.

Corff, K. E., Seideman, R., Venkataraman, P. S., Lutes, L., & Yates, B. (1995). *Facilitated tucking: A nonphamacologic comfortvmeasure for pain in preterm neonates.* JOGNN,24(2),143 – 147.

Emine Efe,&Sevim savaser. (2007).*The effect of two different methods used during peripheral venous blood collection on pain reduction in neonates.* Agri, 19:2, 49-50.

Fitzgerald, M.,& Anand, K.J. S. (1993). *Developmental neuroanatomy and neurophysiology of pain.* In N. L. Schechter et al.,(Eds), Pain in infant, children, and adolescent (pp. 11 - 31). Maryland: Williams & Wilkins.

Fitzgerald, M., & Howard, R. F. (2003). *The Neurobiological Basis of Pediatric Pain*.In Schechter, N., Berde, C. B., & Yaster, M. (Eds.), *Pain in infants, children,and adolescents.* (2nd ed.). (pp. 19 - 42). Philadelphia: Lippincott Williams & Wilkins.

Franck, L.S. and Gregory, G.A. *"Clinical Evaluation and Treatment of Infant Pain in the Neonatal Intensive Care Unit." In Pain in Infant, Children, and Adolescents*.(pp. 519 – 535). Edited by N.L. Schechter et. al. Maryland: Williams & Wilkins, 1993.

Gradin M, Eriksson M, Holmqvist G, Hastein A, Schollin J.(2002). *Pain reduction at vainipuncture in newborns: oral glucose compared with local anesthetic cream. Pediatrics.*110, 1053 -1057.

Grunau, R. V. E.,& Craig, K. D. (1987). *Pain expression in neonates: Facial action and cry.* Pain, 28,395 – 410.

International Association for the Study of Pain. (1994). *"Classification of chronic pain: Pain definitions*."[ระบบออนไลน์]. แหล่งที่มา http://www.iasp-pain .org. (20 มิถุนายน 2554).

Johnson, C. C., Stremler, R., Horton, L., & Friedman, A. (1997). *Single versus triple dose of sucrose for pain in preterm neonates. Pain Research and Management.* Abstracts, 2, 41.

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved

Kolasa, Blair J. (1969). *Introduction to Behavioral Science for Business*. New York: John Wiley and Sons, Inc.

Krechel, S. W., & Bildner, J. (1995). *CRIES: A new neonatal postoperative pain management score inittesting of validity and reliability.* Pediatric Anesthesia, 5,53 – 61.

Laiw, J. J. (2000). *Tactile stimulation and preterm infants.* Journal of perinatal and Neonatal Nursing, 14(1),84 – 103.

Larry Gray. MD, Lisa W. Miller, BA Barbara L. Phillipp, MD and Elliott M. Blass, PhD**. (**2002**).** *Breastfeeding Is Analgesic in Healthy Newborns.Padiatrics.*109, 590.

Lawrence, J., Aleock, D., McGrath, P.J., Kay, J., MacMurray, S. B., & Dulberg, C. (1993). *The development of a tool to assess neonatal pain.* Neonatal network, 12,59 – 66.

Logan P. (1999).*Vainipuncture versus heel prick for the collection of the newborn screening test.* J Adv Nurs 17,30 -36.

Macmilan, Shaver Kelly G. (1997). *Principles of Social Psychology.* Cambridge,Massachusetts:Winthrop Publeshers, Inc.

Mary Renfrew.(1989)**.** *Breast Feeding - Positioning Breast Holding and Latch-On.* Journal of Human Lactation**.**

Melzack, R.,& Wall, P.D.(1965). *Pain mechanism*: a new theory.Science,150,971-979.

Melzack, R.,& Wall, P.D.(1985).*Gate control theory*.In R. H. wilkins& S. S.

Mercer,R.T. (1985). *The process of maternal role attainment over the first year*. Nursing Research, 34, 198 – 204.

Mhipple, B. & Glynn, N.J.(1992). *Quantification of the effects of Listening to music as a noninvasive method of pain control.* Scholarly Inquiry for Nursing Practice: An International Journal, 6, 243 – 258.

Nancy Mohrbacher, IBCLC, Julie .(2003).*The Breastfeeding Answer Book*. (Third Revised Edition), by Stock, MA, IBCLC. La Leche League International, Schaumburg, Illinois.

Newton, M.S. (2000). *Family-Centered Care: Current realities in parent participation*. Pediatric Nursing, 26, 164 – 168.

Poets, C. F., & Southall, D. P. (1994), *Noninvasive monitoring of oxygenation in infants and children: Practical considerations and areas of concern. Pediatrics 93*(5),737 – 746.

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved

Ricardo Carbajal, Soocramanien Veerapen,Sophie Couderc, Myriam Jugie,Yves Ville.(2003). *Analgesic effect of breast feeding in term neonates: randomized controlled trial.* BMJ ,326 ,1

Savadra, M., Eland, J.M., & Tesler, M. (1990). *Pain Managemant In M.J. Craft, & J.A. Dehehy (Eds.), Nursing Intervention for infants and Children.* (pp. 304 - 318). Philadalphia: W.B. Saunders Co.

Schepp,K.(1995). *Psychometric assessment of the preferred participation scale for parent of hospitalized children.* Unpublished manuscript, University of Washington, School of Nursing, Seattle, WA.

Simon, S.H.P., & Tibboel,D. (2006). *Pain perception development and maturation.* Seminars in Fetal & Neonatal Medicine, 11,227 – 231.

Stevens, B. (1999). *Pain in infants.* In MaCaffery, M., & Pasero, C. (Eds.), Pain clinical manual. (2nd ed.). (pp. 626 -773). St. Louis:Mosby.

Steven, B. (2001). *Acute pain management in infants in the neonatal intensive care unit.* In finley, G.A., & McGrat, P.J. (Eds.), Acute and procedure pain in infants and children, progress in research and management Vol. 20. (pp. 101- 128). IASP Press, Seattle.

Steven, B. J., & Johnson, C.C.(1994). *Physiological responses of premature infants to a painful stimulus.* Nursing Research, 43(4),226 – 231.

Steven, B. J., Johnston, C. C., Petryshen, P., & Taddio, A (1996). *The premature infant pain profile: Development and initial validation.* Clinical Journal of pain, 12(1),13 -22.

Tremper, K. V. & Barker, S.J. (1989).*Pulse oximetry.* Anesthesiology, 70(1), 98-108.

Wong D, Eaton MH, Winkestein ML, Wilson D, Ahman E, Thomas PAD.(1999).*Nursing care of infants and children*: St. Louis: Mosby Company.

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved