TABLE OF CONTENTS

	, age
ACKNOWLEDGEMENT	i
ABSTRACT	ii
TABLE OF CONTENTS	vii
LIST OF TABLES	viii
LIST OF FIGURES	, x
INTRODUCTION	1
Organic matter and soil productivity Crop residues in rice-soybean cropping system Green manure as a source of organic matter and nitrogen	3 3 5 7
MATERIALS AND METHODS Informal and formal surveys Field experiment	10 10 11
RESULTS Informal and formal surveys Sources and management of organic matter Potential constraints in increasing soil	15 15 15
productivity through organic replenishment Field experiment Some characteristics of Sesbania rostrata Shoot dry matter of rice N-uptake in straw and grain of rice Rice yield and yield components Economic returns Changes in soil properties	26 29 29 30 31 34 38 43
DISCUSSION Burning rice straw Effects of organic matter on rice crop Effect of organic matter on soil Economic consideration	45 45 45 48 48
CONCLUSION	50
REFERENCES I B h t S I e S e I	52
APPENDIX	56
OUDDION UN VITAT	60

LIST OF TABLES

Table		Page
1.	Treatment combinations	11
2.	Soybean residue management in the farmers' fields	17
3.	The amount and nutrient content of soybean residues in the farmers' fields	18
4.	Rice straw management in farmers' fields	19
5.	The amount and nutrient content of rice straw in the farmers' fields	20
6.	The amount and management of soybean residues in the farmers' fields	21
7.	The amount and management of rice straw in the farmers'fields	21
8.	Animal manure and chemical fertilizer used in the rice fields and the soybean fields	23
9.	C/N ratio and the nutrient content of animal manure in the farmers' fields	24
10.	Farmers' activities during April-July	25
11.	Plant density, biomass, dry matter, nutrient accumulation and carbon content of <i>Sesbania rostrata</i> at 50 days old	29
12.	The shoot dry matter (t/ha) of rice as influenced by different N sources averaged over varieties at panicle initiation stage	y e 30
13.	Average N-uptake (kgN/ha) in straw of RD7 and KDML109 at different stages	5 3 2
14.	N-uptake (kgN/ha)in rice grain of various	s (e. 33
15.	The N concentration (%) in rice grain of variou treatments	s 34
16.	Grain yield (kg/ha) of rice in differen treatments	t 36
17.	The number of panicles per square-meter in differen	t 37

18.	The 1000-grain weight (g) of rice in different treatments	37
19.	The percentage of filled grain of rice in different treatments	38
20.	Total variable costs and return to individual treatments (B/ha)	40
21.	Marginal rate of return in treatments with different N sources comparing to no N	41
22.	Marginal rates of return of individual treatments comparing to no N at different prices of urea	42
23.	Maximum marginal rates of return treatments at different costs of land preparation and urea	42
24.	Soil analyses results before and after harvest	44
Appen	dix Table	
1	ANOVA of shoot dry matter (kg/ha) of rice at different stages	59
2	ANOVA of N-uptake (kg/ha) in rice straw at different	
	stages	59
3	ANOVA of N-uptake (kg/ha) in grain of two rice varieties in various N sources	59 60
3	ANOVA of N-uptake (kg/ha) in grain of two rice vari-	
	ANOVA of N-uptake (kg/ha) in grain of two rice varieties in various N sources ANOVA of grain yield (kg/ha) of two rice varieties in	60

Copyright[©] by Chiang Mai University All rights reserved

LIST OF FIGURES

Figure	
. The flow of organic matter in rice-soybean croppi system	ng 16
Green manure (GM) and rice-soybean cropping system Chiang Mai Valley in relation to rainfall and irrigat water distribution	
Shoot dry matter of rice as influenced by different sources averaged over varieties at panicle initiati stage(PI), flowering stage(F) and harvesting stage (on
. N-uptake in rice straw at panicle initiation stage(PI flowering stage (F) and harvesting stage(H)	32
ppendix Figure	
The layout of field experiment (Factorial in Randomiz Complete Block Design)	ed 57
Decision tree: minimum necessary criteria for gremanure production (Garrity and Flinn, 1988)	en 58

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่ Copyright[©] by Chiang Mai University All rights reserved