

## Table of Contents

	Page
Acknowledgements	I
Abstract	III
Thai Abstract	VI
Table of Contents	VIII
List of Tables	XV
List of Figures	XVII
List of Plates	XVIII
<b>Chapter I: General Introduction</b>	<b>1</b>
<b>1.1: Background</b>	<b>1</b>
<b>1.2: Rationale</b>	<b>4</b>
<b>1.3: Objectives</b>	<b>6</b>
<b>Chapter II: Literature Review</b>	<b>7</b>
<b>2.1: Fungi associated with mungbean and blackgram seeds</b>	<b>7</b>
<b>2.2: Pathogenicity of <i>Macrophomina phaseolina</i></b>	<b>8</b>
<b>2.3: Location of <i>Macrophomina phaseolina</i> in mungbean and blackgram seeds</b>	<b>10</b>
<b>2.4: Seed transmission of <i>Macrophomina phaseolina</i></b>	<b>11</b>
<b>2.5: Effect of fungi on seed quality</b>	<b>14</b>
<b>2.6: Effect of fungi on carbohydrate and protein content of seeds</b>	<b>15</b>
<b>2.7: Control of seed-borne diseases by seed treatment</b>	<b>18</b>
2.7.1: Seed treatment by hot water	19
2.7.2: Chemical seed treatment	20
2.7.3: Biological control	21

<b>Chapter III: Fungi Associated with Mungbean and Blackgram</b>	
<b>Seeds</b>	<b>27</b>
<b>3.1: An Overview</b>	<b>27</b>
<b>3.2: Materials and Methods</b>	<b>28</b>
3.2.1: Source of seeds	28
3.2.2: Incubation and detection technique	28
<b>3.3: Results</b>	<b>29</b>
3.3.1: Detection of <i>Macrophomina phaseolina</i> and other fungi associated with naturally infected seeds of mungbean.	29
3.3.2: Detection of <i>Macrophomina phaseolina</i> and other fungi associated with naturally infected seeds of blackgram.	39
<b>3.4: Discussion</b>	<b>45</b>
<b>Chapter IV: Pathogenicity of <i>Macrophomina phaseolina</i> as a Seed-borne Pathogen of Mungbean and Blackgram</b>	<b>48</b>
<b>4.1: An Overview</b>	<b>48</b>
<b>4.2: Materials and Methods</b>	<b>49</b>
4.2.1: Experimental materials	49
4.2.2: Preparation of pure culture of <i>M. phaseolina</i>	49
4.2.3: Preparation of suspension of <i>M. phaseolina</i> pure culture	50
4.2.4: Determination of pathogenicity of <i>M. phaseolina</i>	50
4.2.4.1: Seed inoculation method	52
4.2.4.2: Sprout dipping method	52
4.2.4.3: Soil inoculation method	53
4.2.4.4: Damping-off symptom test method	53
4.2.4.5: Test tube seedling symptom test	54
4.2.4.6: Toothpic inoculation method	54
<b>4.3: Results</b>	<b>55</b>
4.3.1: Seed inoculation method	55

4.3.2: Sprout dipping method	57
4.3.3: Soil inoculation method	60
4.3.4: Damping-off method	60
4.3.5: Test tube seedling symptom test	65
4.3.6: Toothpic inoculation method	65
<b>4.4: Discussion</b>	<b>72</b>
<b>Chapter V: Location of <i>Macrophomina phaseolina</i> in Mungbean and Blackgram Seed</b>	<b>76</b>
<b>5.1: An Overview</b>	<b>76</b>
<b>5.2: Materials and Methods</b>	<b>77</b>
5.2.1: Source of seed	77
5.2.2: Detection the location of <i>M. phaseoliana</i>	77
<b>5.3: Results</b>	<b>78</b>
<b>5.4: Discussion</b>	<b>78</b>
<b>Chapter VI: Seed Transmission of <i>Macrophomina phaseolina</i> in Mungbean and Blackgram Seeds</b>	<b>83</b>
<b>6.1: An Overview</b>	<b>83</b>
<b>6.2: Materials and Methods</b>	<b>84</b>
6.2.1: Seed source	84
6.2.2: Methods of studying seed to seedling transmission	85
<b>6.3: Results</b>	<b>86</b>
6.3.1: Seed transmission of <i>M. phaseolina</i> in mungbean	86
6.3.2: Seed transmission of <i>M. phaseolina</i> in blackgram	97
<b>6.4: Discussion</b>	<b>98</b>
<b>Chapter VII: Effect of <i>Macrophomina phaseolina</i> on Viability, Vigor and Storability of Mungbean and Blackgram Seeds</b>	<b>101</b>
<b>7.1: An Overview</b>	<b>101</b>
<b>7.2: Materials and Methods</b>	<b>102</b>
7.2.1: Source of seed	102

7.2.2: Isolation and preparation of pure culture and inoculum of <i>Macrophomina phaseolina</i>	102
7.2.3: Inoculation of <i>Macrophomina phaseolina</i> inoculum	103
7.2.4: Viability test	103
7.2.5: Vigor test	104
7.2.6: Storability test	104
<b>7.3: Results</b>	<b>105</b>
7.3.1: Viability test	105
7.3.1.1: Viability of mungbean	105
7.3.1.2: Viability of blackgram	105
7.3.2: Vigor test	107
7.3.2.1: Vigor test of mungbean	107
7.3.2.2: Vigor test of blackgram	107
7.3.3: Storability test	109
7.3.3.1: Storability of mungbean	109
7.3.3.2: Storability of blackgram	109
<b>7.4: Discussion</b>	<b>111</b>
<b>Chapter VIII: Effect of <i>Macrophomina phaseolina</i> on Carbohydrate and Protein Content in Mungbean and Blackgram Seeds</b>	<b>113</b>
<b>8.1: An Overview</b>	<b>113</b>
<b>8.2: Materials and Methods</b>	<b>114</b>
8.2.1: Source of sample	114
8.2.2: Infected and healthy seed separation and sample preparation	114
8.2.3: Carbohydrate analysis by 'Anthrone' method	114
8.2.3.1: Reagent preparation	114
8.2.3.2: Procedure	115
8.2.4: Protein analysis by Micro-Kjeldahl method	116
8.2.4.1: Reagent preparation	116
8.2.4.2: Procedure	117

8.2.4.3: Calculation	118
<b>8.3: Result</b>	<b>119</b>
8.3.1: Carbohydrate content	119
8.3.2: Protein content	119
<b>8.4: Discussion</b>	<b>119</b>
<b>Chapter IX: Control of <i>Macrophomina phaseolina</i> in Mungbean and Blackgram Seeds Through Seed Treatment</b>	<b>123</b>
<b>9.1: Physical Seed Treatment by Hot Water</b>	<b>123</b>
<b>9.1.1: An Overview</b>	<b>124</b>
<b>9.1.2: Materials and Methods</b>	<b>126</b>
9.1.2.1: Seed sample	126
9.1.2.2: Seed treatment	126
9.1.2.3: Evaluation of hot water treated seeds by blotter method	126
9.1.2.4: Plantation of hot water treated seeds in <i>in-vivo</i> condition	127
<b>9.1.3: Results</b>	<b>128</b>
9.1.3.1: Evaluation of hot water treated mungbean seeds by blotter method	128
9.1.3.2: Evaluation of hot water treated blackgram seeds by blotter method	130
9.1.3.3: Plantation of hot water treated seed in pot	130
<b>9.1.4: Discussion</b>	<b>134</b>
<b>9.2: Chemical Seed Treatment by Seed-dressing Fungicides</b>	<b>135</b>
<b>9.2.1: An Overview</b>	<b>135</b>
<b>9.2.2: Materials and Methods</b>	<b>136</b>
9.2.2.1: <i>In-vitro</i> trial	136
9.2.2.2: Seed treatment with effective fungicides	137
9.2.2.3: Incubation of treated seeds by blotter method	137
9.2.2.4: Evaluation of treated seeds <i>in-vivo</i> (pot experiment)	138
<b>9.2.3: Results</b>	<b>138</b>

9.2.3.1: <i>In-vitro</i> trial	138
9.2.3.2: Incubation of seeds with screened and effective fungicides by blotter method	143
9.2.3.3: Evaluation of treated seeds <i>in-vivo</i>	147
<b>9.2.4: Discussion</b>	<b>152</b>
<b>9.3: Biological Seed Treatments by Biological Antagonists</b>	<b>152</b>
<b>9.3.1: An Overview</b>	<b>152</b>
<b>9.3.2: Materials and Methods</b>	<b>155</b>
9.3.2.1: Screening of effective antagonists <i>in-vitro</i> against <i>M. phaseolina</i>	155
9.3.2.1.1: Isolation of <i>M. phaseolina</i>	155
9.3.2.1.2: Collection and preparation of antagonistic agents	156
9.3.2.1.3: Dual culture	156
9.3.2.1.4: Regression	158
9.3.2.1.5: Seed treatment with screened antagonists	158
9.3.2.1.5.1: Seed sample	158
9.3.2.1.5.2: Seed treatment	158
9.3.2.1.5.3: Evaluation of treated seed by blotter method	159
9.3.2.1.5.4: Evaluation of treated seed by <i>in-vivo</i> (pot culture)	159
<b>9.3.3: Results</b>	<b>159</b>
9.3.3.1: Screening of effective antagonists <i>in-vitro</i> against <i>M. phaseolina</i>	159
9.3.3.2: Regression	161
9.3.3.3: Seed treatment	169
9.3.3.3.1: Evaluation of treated seeds by blotter method	169
9.3.3.3.2: Evaluation of treated seeds <i>in-vivo</i> (pot culture)	170

<b>9.3.4: Discussion</b>	<b>175</b>
<b>Chapter X: Conclusions and Recommendations</b>	<b>179</b>
<b>References</b>	<b>182</b>
<b>Appendices</b>	<b>198</b>
<b>Curriculum vitae</b>	<b>226</b>

มหาวิทยาลัยเชียงใหม่  
Chiang Mai University

## LIST OF TABLES

Table	Description of Tables	Pages
3.01	Percentage of fungi associated with the seeds of different mungbean varieties and elite lines (average result of four replications, each replication contained 100 seeds).	30
3.02	Percentage of fungi associated with the seeds of blackgram varieties (average of four replications. Each replication contained 100 seeds).	40
4.01	Pathogenicity determination in mungbean and blackgram seeds by seed inoculation method	55
4.02	Pathogenicity determination in mungbean and blackgram sprouts by sprout dipping method	57
4.03	Pathogenicity determination in mungbean and blackgram seeds by soil inoculation method	61
4.04	Pathogenicity determination in mungbean and blackgram by damping-off method	64
4.05	Pathogenicity determination in mungbean and blackgram seeds by test tube seedling symptom test	66
4.06	Pathogenicity determination in mungbean and blackgram plants by toothpic inoculation method (mean results of ten plants)	68
5.01	Percentage of <i>M. phaseolina</i> incidence in different seed Components of mungbean and blackgram.	80
6.01	Types of infection resulting from <i>Macrophomina phaseolina</i> infected mungbean and blackgram seeds in blotter method (Based on 400 seeds)	87
6.02	Types of infection resulting from <i>Macrophomina phaseolina</i> infected mungbean and blackgram seeds in Test tube agar method (Based on 100 seeds)	90
6.03	Percentage of seed to seedling transmission of <i>Macrophomina phaseolina</i> (sand and soil method, based on 100 seeds)	93
7.01	Effect of <i>Macrophomina phaseolina</i> on different variables of seed viability in mungbean seeds (mean of four replications)	106



7.02	Effect of <i>Macrophomina phaseolina</i> on different variables of seed viability in blackgram seeds (mean of four replications)	106
7.03	Effect of <i>Macrophomina phaseolina</i> on different variables of vigor estimation in mungbean seeds (mean of four replications)	108
7.04	Effect of <i>Macrophomina phaseolina</i> on different variables of vigor estimation in blackgram seeds (mean of four replications)	108
7.05	Effect of <i>Macrophomina phaseolina</i> on different variables of storability estimation in mungbean seeds (mean of four replications)	110
7.06	Effect of <i>Macrophomina phaseolina</i> on different variables of storability in blackgram seeds (mean of four replications)	110
8.01	Percentage of carbohydrate content in healthy and diseased seeds of mungbean and blackgram	120
8.02	Percentage of protein content in healthy and inoculated seeds of mungbean and blackgram	120
9.01	Effect of hot water treatment on <i>Macrophomina phaseolina</i> infection and on germination when the seeds were planted in plastic pot with sterilized soil (mean of four replications)	132
9.02	Effect of fungicidal seed treatment on <i>Macrophomina phaseolina</i> infection and germination according to blotter method with promising fungicides (results show the average of four replications).	147
9.03	Fungicidal seed treatment against <i>Macrophomina phaseolina</i> infection and germination planting the seed in plastic pot with sterilized soil (mean of four replications)	148
9.04	Seed coating with promising antagonists for controlling <i>Macrophomina phaseolina</i> in mungbean and blackgram (results show the average of four replications)	170
9.05	Biological seed treatment with <i>Trichoderma</i> sp. Against <i>Macrophomina phaseolina</i> infection and germination when planting the seed in plastic pot with sterilized soil (mean of four replications).	171

## LIST OF FIGURES

Figure	Description of figures	Pages
3.01	Percent infection of <i>Macrophomina phaseolina</i> and germination of seeds in different mungbean and blackgram varieties and elite lines	31
9.01	Effect of hot water treatment in different temperatures and durations for controlling seed borne <i>Macrophomina phaseolina</i> in mungbean seed	129
9.02	Effect of hot water treatment in different temperature and durations for controlling seed-borne <i>Macrophomina phaseolina</i> in blackgram seeds	131
9.03	Percentage of decrease of radial growth of <i>Macrophomina phaseolina</i> over control in different fungicides	139
9.04	Radius of <i>Macrophomina phaseolina</i> as control	157
9.05	Radius of <i>Macrophomina phaseolina</i> after antagonism	157
9.06	Inhibition percentage of radial growth of <i>Macrophomina phaseolina</i> by different antagonists	160
9.07	The trend of inhibition of <i>Trichoderma hamatum</i> against <i>Macrophomina phaseolina</i>	166
9.08	The trend of inhibition of <i>Trichoderma harzianum</i> against <i>Macrophomina phaseolina</i>	166
9.09	The trend of inhibition of <i>Trichoderma viride</i> against <i>Macrophomina phaseolina</i>	167
9.10	The trend of inhibition of <i>Bacillus cereus</i> against <i>Macrophomina phaseolina</i>	167
9.11	The trend of inhibition of <i>Pseudomonas aeruginosa</i> against <i>Macrophomina phaseolina</i>	168
9.12	The trend of inhibition of <i>Pseudomonas putida</i> against <i>Macrophomina phaseolina</i>	168

## LIST OF PLATES

Plates	Description of Plates	Plates
3.01	1. Mungbean seed incubated by blotter method. 2. <i>Macrophomina phaseolina</i> infection appeared after incubation by blotter blotter method.	32
3.02	Microsclerotia (M) and pycnidia (P) on the <i>Macrophomina phaseolina</i> infected radicle of mungbean under stereomicroscope.	33
3.03	A. <i>Macrophomina phaseolina</i> infected mungbean seed Under stereomicroscope with numerous pycnidia. B. Pycnidia (P) of <i>Macrophomina phaseolina</i> on Mungbean seed under stereomicroscope.	34
3.04	Pycnidia (P) of <i>Macrophomina phaseolina</i> from mungbean seed under compound microscope.	35
3.05	Liberating of conidia (C) from ruptured pycnidium (P) of <i>Macrophomina phaseolina</i> of mungbean seed observed under compound microscope.	35
3.06	Conidia (C) from pycnidium of <i>Macrophomina phaseolina</i> from mungbean seed observed under compound microscope.	36
3.07	<i>Macrophomina phaseolina</i> infected mungbean seed spreading infection (I) to healthy seedlings by touching.	38
3.08	1. Blackgram seed incubated by blotter method. 2. <i>Macrophomina phaseolina</i> infection appeared after incubation by blotter method.	41
3.09	A. <i>Macrophomina phaseolina</i> infected blackgram seed Under stereomicroscope with dead radicle and without plumule B. Numerous pycnidia of <i>Macrophomina phaseolina</i> on Blackgram seed.	42
3.10	Microsclerotia (M) and pycnidia (P) on the <i>Macrophomina phaseolina</i> infected radicle of blackgram under stereo microscope.	43
3.11	<i>Macrophomina phaseolina</i> infected blackgram seed spreading infection (I) to healthy seedlings by touching.	43

3.12	Pycnidia (P) of <i>Macrophomina phaseolina</i> from blackgram seed under compound microscope.	44
3.13	Liberating of conidia (C) from ruptured pycnidium of <i>Macrophomina phaseolina</i> of blackgram seed observed under compound microscope.	46
3.14	Conidia (C) from pycnidium of <i>Macrophomina phaseolina</i> from blackgram seed observed under compound microscope.	46
4.01	Pure culture of <i>Macrophomina phaseolina</i> on artificial media (PDA) growing in different days.	51
4.02	Mungbean seedling from <i>Macrophomina phaseolina</i> inoculated (left) and noninoculated (right) seed.	56
4.03	Blackgram seedling from <i>Macrophomina phaseolina</i> inoculated (left) and non-inoculated (right) seed.	56
4.04	Mungbean seedlings after dipping the sprout in <i>Macrophomina phaseolina</i> suspension (left) and in sterilized water (right).	58
4.05	Blackgram seedlings after dipping the sprout in <i>Macrophomina phaseolina</i> suspension (left) and in sterilized water (right).	59
4.06	Seed rot from <i>Macrophomina phaseolina</i> inoculated soil.	62
4.07	<i>Macrophomina phaseolina</i> infected cotyledonary leaves.	62
4.08	Infection of <i>Macrophomina phaseolina</i> is extending from infected cotyledonary leaves.	62
4.09	Lesion after infection of <i>Macrophomina phaseolina</i> on the stem.	63
4.10	Lesion on the leaf after <i>Macrophomina phaseolina</i> infection.	63
4.11	Seedling died just after emergence due to <i>Macrophomina phaseolina</i> infection.	63
4.12	Emerging seedling of mungbean from <i>Macrophomina phaseolina</i> inoculated plate (left) and from noninoculated plate (right).	64

4.13	Emerging seedling of blackgram from <i>Macrophomina phaseolina</i> inoculated plate (left) and from non-inoculated plate (right).	64
4.14	Mungbean seedling in <i>Macrophomina phaseolina</i> inoculated agar medium (left and middle), mungbean seedling in non-inoculated medium (right).	67
4.15	Blackgram seedling in <i>Macrophomina phaseolina</i> inoculated medium (left and middle), blackgram seedling in non-inoculated medium (right).	68
4.16	Mungbean stem inoculation with <i>Macrophomina phaseolina</i> . Control treatment (left) and inoculated stem (right).	69
4.17	Blackgram stem inoculation with <i>Macrophomina phaseolina</i> . Control treatment (left) and inoculated stem (right).	69
4.18	Spreading the lesion causing break down of stem followed by producing numerous pycnidia and microsclerotia of <i>Macrophomina phaseolina</i> in mungbean.	70
4.19	Spreading the lesion causing break down of stem followed by producing numerous pycnidia and microsclerotia of <i>Macrophomina phaseolina</i> in blackgram.	70
4.20	Mungbean internode inoculated with <i>Macrophomina phaseolina</i> . Control treatment (left) and inoculated internode resulted charcoal like necrosis (right).	71
4.21	Blackgram internode inoculated with <i>Macrophomina phaseolina</i> . Control treatment (left) and inoculated internode resulted charcoal like necrosis (right).	71
4.22	Spreading of lesion due to <i>Macrophomina phaseolina</i> infection to upward and downward directions in mungbean causing death of whole plant.	73
4.23	Spreading of lesion due to <i>Macrophomina phaseolina</i> infection to upward and downward directions in blackgram causing death of whole plant.	73
4.24	Control (left) and <i>Macrophomina phaseolina</i> inoculated plant (right) of mungbean after 21 days of inoculation.	74

4.25	Control (left) and <i>Macrophomina phaseolina</i> inoculated plant (right) of blackgram after 21 days of inoculation.	74
5.01	Detection of site of <i>M. phaseolina</i> in different parts of mungbean seed.	79
5.02	Detection of site of <i>M. phaseolina</i> in different parts of blackgram seed by blotter method.	79
5.03	Pycnidia and sclerotia of <i>M. phaseolina</i> were observed in the seed coat of mungbean under stereo-binocular microscope.	81
5.04	Pycnidia and sclerotia of <i>M. phaseolina</i> were observed in the seed coat of blackgram under stereo-binocular microscope.	81
6.01	<i>Macrophomina phaseolina</i> transmitted mungbean seed.	87
6.02	Pycnidia of <i>Macrophomina phaseolina</i> formation on the seed coat of mungbean.	89
6.03	Pycnidia and microsclerotia of <i>Macrophomina phaseolina</i> on the un-germinated seed.	89
6.04	<i>Macrophomina phaseolina</i> transmitted mungbean seed in test tube agar method.	90
6.05	Un-germinated seed with heavy mycelia of <i>Macrophomina phaseolina</i> observed in test-tube agar method	91
6.06	<i>Macrophomina phaseolina</i> transmitted seedling (left) and without transmitted healthy seedling (right) observed in test-tube agar method	91
6.07	Transmission of <i>Macrophomina phaseolina</i> to the cotyledonary leaves of mungbean planted in sand at 4 days.	94
6.08	Transmission of <i>Macrophomina phaseolina</i> to the cotyledonary leaves of mungbean planted in soil at 4 days.	94

6.09	Spreading of infection due to <i>Macrophomina phaseolina</i> from the cotyledonary leaves to true leaves (left) and a healthy seedling without infection (right) at the age of 3 days.	95
6.10	Eventual death of seedling due to <i>Macrophomina phaseolina</i> infection (left) and a healthy seedling (right) at the age of 5 days.	95
6.11	In sand, <i>Macrophomina phaseolina</i> transmitted (left) and healthy seedlings (right) of mungbean at the age of 6 days.	96
6.12	In soil, <i>Macrophomina phaseolina</i> transmitted (left) and healthy seedlings (right) of mungbean at the age of 6 days.	96
9.01	Control treatment (without treated) and hot water treated (at most effective time and duration) seedlings of mungbean at the age of 8 days after plantation.	133
9.02	Control treatment (without treated) and hot water treated (at most effective time and duration) seedlings of blackgram at the age of 8 days after plantation	133
9.03	Radial growth of <i>Macrophomina phaseolina</i> on Benlate mixed PDA after 3 days of plating.	140
9.04	Radial growth of <i>Macrophomina phaseolina</i> on Benlate mixed PDA after 5 days of plating.	140
9.05	Radial growth of <i>Macrophomina phaseolina</i> on Dithane M-45 mixed PDA after 3 days of plating.	141
9.06	Radial growth of <i>Macrophomina phaseolina</i> on Dithane M-45 mixed PDA after 5 days of plating.	141
9.07	Radial growth of <i>Macrophomina phaseolina</i> on Thiram mixed PDA after 3 days of plating.	142
9.08	Radial growth of <i>Macrophomina phaseolina</i> on Thiram mixed PDA after 5 days of plating.	142
9.09	Radial growth of <i>Macrophomina phaseolina</i> on Vitavax mixed PDA after 3 days of plating.	144

9.10	Radial growth of <i>Macrophomina phaseolina</i> on Vitavax mixed PDA after 5 days of plating.	144
9.11	Radial growth of <i>Macrophomina phaseolina</i> on Captan mixed PDA after 3 days of plating.	145
9.12	Radial growth of <i>Macrophomina phaseolina</i> on Captan mixed PDA after 5 days of plating.	145
9.13	Radial growth of <i>Macrophomina phaseolina</i> on Metalexyl mixed PDA after 3 days of plating.	146
9.14	Radial growth of <i>Macrophomina phaseolina</i> on Metalexyl mixed PDA after 5 days of plating.	146
9.15	Comparison of control (untreated) and Benlate treated seedlings of mungbean at the age of 8 days after plantation	149
9.16	Comparison of control (untreated) and Benlate treated seedlings of blackgram at the age of 8 days after plantation	149
9.17	Comparison of control (untreated) and Dithane M-45 treated seedlings of mungbean at the age of 8 days after plantation	150
9.18	Comparison of control (untreated) and Dithane M-45 treated seedlings of blackgram at the age of 8 days after plantation.	150
9.19	Comparison of control (untreated) and Thiram treated seedlings of mungbean at the age of 8 days after plantation.	151
9.20	Comparison of control (untreated) and Benlate treated seedlings of blackgram at the age of 8 days after plantation	151
9.21	Control and dual culture of different <i>Trichoderma</i> spp. against <i>Macrophomina phaseolina</i> at 3rd day.	162
9.22	Control and dual culture of different <i>Trichoderma</i> spp. against <i>Macrophomina phaseolina</i> at 5th day.	163



9.23	Control and dual culture of different bacterial antagonists against <i>Macrophomina phaseolina</i> at 3rd day.	164
9.24	Control and dual culture of different bacterial antagonists against <i>Macrophomina phaseolina</i> at 5th day.	165
9.25	Comparison of control (untreated) and <i>Trichoderma harzianum</i> treated seedlings of mungbean at the age of 8 days after plantation	172
9.26	Comparison of control (untreated) and <i>Trichoderma harzianum</i> treated seedlings of blackgram at the age of 8 days after plantation	172
9.27	Comparison of control (untreated) and <i>Trichoderma hamatum</i> treated seedlings of mungbean at the age of 8 day after plantation	173
9.28	Comparison of control (untreated) and <i>Trichoderma hamatum</i> treated seedlings of blackgram at the age of 8 days after plantation	173
9.29	Comparison of control (untreated) and <i>Trichoderma viride</i> treated seedlings of mungbean at the age of 8 days after plantation	174
9.30	Comparison of control (untreated) and <i>Trichoderma viride</i> treated seedlings of blackgram at the age of 8 days after plantation	174