

TABLE OF CONTENTS

	page
Acknowledgement	iv
Abstract in Thai	vii
Abstract in English	xi
List of tables	xv
List of figures	1
Chapter 1 Introduction	3
Chapter 2 Literature review	13
Chapter 3 Material and Methods	15
<i>Part I Studies on factors influencing growth and in vitro tuberization.</i>	16
Experiment 1 Effect of sucrose concentration	18
Experiment 2 Effect of coconut water concentration	22
Experiment 3 Effects of coconut water and BAP concentrations	26
Experiment 4 Effects of IAA, BAP and coconut water concentrations in the dark condition	36
Experiment 5 Effects of illumination and sucrose concentration	44
<i>Part II Histological study on tuberization.</i>	54
1. The first tuberization of a seedling	55
2. The tuberization of a mature plant	60
<i>Part III Studies on the changing of internal macro elements, free amino acids and some growth regulators in mature plants during tuberization.</i>	68
1. Analysis of macro elements	68
2. Analysis of free amino acid contents	68
3. Analysis of free indoleacetic acid (IAA) and abscisic acid (ABA) contents	68
Chapter 4 Results	68
<i>Part I Studies on factors influencing growth and in vitro tuberization.</i>	68
Experiment 1 Effect of sucrose concentration	68
Experiment 2 Effect of coconut water concentration	72
Experiment 3 Effects of coconut water and BAP concentrations	76
Experiment 4 Effects of IAA, BAP and coconut water concentrations in the dark condition	86
Experiment 5 Effects of illumination and sucrose concentration	94
Conclusion results in Part I	100
<i>Part II Histological study on tuberization.</i>	105
1. The first tuberization of a seedling	105
2. The tuberization of a mature plant	110
<i>Part III Studies on the changing of internal macro elements, free amino acids and some growth regulators in mature plants during tuberization.</i>	118
1. Fresh weight, dry weight and water contents	118

2. Analysis of macro elements	68
3. Analysis of free amino acid contents	69
4. Analysis of free indoleacetic acid (IAA) and abscisic acid (ABA) contents	72
Chapter 5 Discussions	
Part I Studies on factors influencing growth and in vitro tuberization.	74
Part II Histological study on tuberization.	84
Part III Studies on the changing of internal macro elements, free amino acids and some growth regulators in mature plants during tuberization.	86
Chapter 6 Conclusions	90
References	92
Appendices	
Appendix A Compositions of CMU ₁ medium	97
Appendix B Histological preparation	98
Appendix C Chemical analysis	100
Curriculum vitae	112

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
 Copyright© by Chiang Mai University
 All rights reserved

LIST OF TABLES

Table		page
1	Combinations of coconut water and BAP concentrations.	14
2	Combinations of growth regulators and coconut water in each treatment.	14
3	Combinations of illumination and sucrose concentrations.	15
4	Effect of sucrose concentration on leaf-emerging percentage of <i>Pecteilis sagarikii</i> seedlings during culturing for 10 weeks.	18
5	Effect of sucrose concentration on shoot height, number of leaf and leaf size of <i>Pecteilis sagarikii</i> seedlings after culturing for 10 weeks.	19
6	Effect of sucrose concentration on tuberization percentage of <i>Pecteilis sagarikii</i> seedlings during culturing for 10 weeks.	19
7	Effect of sucrose concentration on number of tuber, tuber width and tuber length, and oval-shape tuber percentage of <i>Pecteilis sagarikii</i> seedlings after culturing for 10 weeks.	21
8	Effect of sucrose concentration on rooting percentage of <i>Pecteilis sagarikii</i> seedlings during culturing for 10 weeks.	22
9	Effect of sucrose concentration on number of root, root width and root length of <i>Pecteilis sagarikii</i> seedlings after culturing for 10 weeks.	22
10	Effect of coconut water concentration on leaf-emerging percentage of <i>Pecteilis sagarikii</i> seedlings during culturing for 10 weeks.	23
11	Effect of coconut water concentration on shoot height, number of leaf and leaf size of <i>Pecteilis sagarikii</i> seedlings after culturing for 10 weeks.	23
12	Effect of coconut water concentration on tuberization percentage of seedlings during culturing for 10 weeks.	24
13	Effect of coconut water concentration on number of tuber, tuber size and oval-shape tuber percentage of <i>Pecteilis sagarikii</i> seedlings after culturing for 10 weeks.	24
14	Effect of coconut water concentration on rooting percentage of seedlings during culturing for 10 weeks.	25
15	Effect of coconut water concentration on number of root, root width and root length of <i>Pecteilis sagarikii</i> seedlings after culturing for 10 weeks.	25
16	Effects of coconut water and BAP concentrations on leaf-emerging percentages of the protocorms during culturing for 10 weeks.	26
17	Main effect of coconut water concentration on shoot height, number of leaf and leaf size of <i>Pecteilis sagarikii</i> seedlings after culturing for 10 weeks.	26

Table	page
18	Main effect of BAP concentration on shoot height, number of leaf and leaf size of <i>Pecteilis sagarikii</i> seedlings after culturing for 10 weeks. 27
19	Effects of coconut water and BAP concentrations on shoot height, number of leaf and leaf size of <i>Pecteilis sagarikii</i> seedlings after culturing for 10 weeks. 28
20	Effects of coconut water and BAP concentrations on tuberization percentages of <i>Pecteilis sagarikii</i> seedlings during culturing for 10 weeks. 28
21	Main effect of coconut water concentration on number of tuber and tuber size of <i>Pecteilis sagarikii</i> seedlings after culturing for 10 weeks. 29
22	Main effect of BAP concentration on number of tuber and tuber size of <i>Pecteilis sagarikii</i> seedlings after culturing for 10 weeks. 29
23	Effects of coconut water and BAP concentrations on number of tuber, tuber size and oval-shape tuber percentage of <i>Pecteilis sagarikii</i> seedlings after culturing for 10 weeks. 30
24	Effects of coconut water and BAP concentrations on rooting percentages of <i>Pecteilis sagarikii</i> seedlings during culturing for 10 weeks. 31
25	Main effect of coconut water concentration on number of root, root width and root length of <i>Pecteilis sagarikii</i> seedlings after culturing for 10 weeks. 32
26	Main effect of BAP concentration on number of root, root width and root length of <i>Pecteilis sagarikii</i> seedlings after culturing for 10 weeks. 32
27	Effects of coconut water and BAP concentrations on number of root, root width and root length of <i>Pecteilis sagarikii</i> seedlings after culturing for 10 weeks. 33
28	Effects of coconut water and BAP concentrations on protocorm-like bodies forming percentages of <i>Pecteilis sagarikii</i> seedlings during culturing for 10 weeks. 34
29	Main effect of coconut water concentration on number of protocorm-like bodies of <i>Pecteilis sagarikii</i> seedlings after culturing for 10 weeks. 34
30	Main effect of BAP concentration on number of protocorm-like bodies of <i>Pecteilis sagarikii</i> seedlings after culturing for 10 weeks. 35
31	Effects of coconut water and BAP concentrations on number of protocorm-like bodies of <i>Pecteilis sagarikii</i> seedlings after culturing for 10 weeks. 35
32	Effects of IAA, BAP and coconut water concentrations on leaf-emerging percentages of protocorms during culturing for 10 weeks. 36
33	Effects of IAA, BAP and coconut water concentrations on shoot height, number of leaf and leaf size of <i>Pecteilis sagarikii</i> seedlings after culturing for 10 weeks. 37

Table		Page
34	Effects of IAA, BAP and coconut water concentrations on tuberization percentages of seedlings during culturing for 10 weeks.	37
35	Effects of IAA, BAP and coconut water concentrations on number of tuber, tuber size and oval-shape tuber percentage of <i>Pecteilis sagarikii</i> seedlings after culturing for 10 weeks.	38
36	Effects of IAA, BAP and coconut water concentrations on rooting percentages of seedlings during culturing for 10 weeks.	40
37	Effects of IAA, BAP and coconut water concentrations on number of root, root width and root length of <i>Pecteilis sagarikii</i> seedlings after culturing for 10 weeks.	40
38	Effects of IAA, BAP and coconut water concentrations on total soluble sugar in various tissues of <i>Pecteilis sagarikii</i> seedlings after culturing for 10 weeks.	41
39	Effects of IAA, BAP and coconut water concentrations on starch contents in various tissues of <i>Pecteilis sagarikii</i> seedlings after culturing for 10 weeks.	42
40	Effects of illumination and sucrose concentration on leaf-emerging percentages of <i>Pecteilis sagarikii</i> seedlings during culturing for 12 weeks.	44
41	Main effect of illumination on shoot height, number of leaf and leaf size of <i>Pecteilis sagarikii</i> seedlings after culturing for 12 weeks.	45
42	Main effect of sucrose concentration on shoot height, number of leaf and leaf size of <i>Pecteilis sagarikii</i> seedlings after culturing for 12 weeks.	45
43	Effects of illumination and sucrose concentration on shoot height, number of leaf and leaf size of <i>Pecteilis sagarikii</i> seedlings after culturing for 12 weeks.	46
44	Effects of illumination and sucrose concentrations on tuberization percentages of <i>Pecteilis sagarikii</i> seedlings during culturing for 12 weeks.	47
45	Main effect of illumination on number of tuber and tuber size of <i>Pecteilis sagarikii</i> seedlings after culturing for 12 weeks.	47
46	Main effect of sucrose concentration on number of tuber and tuber size of <i>Pecteilis sagarikii</i> seedlings after culturing for 12 weeks.	48
47	Effects of illumination and sucrose concentration on number of tuber and tuber size of <i>Pecteilis sagarikii</i> seedlings after culturing for 12 weeks.	48
48	Effects of illumination and sucrose concentrations on rooting percentages of <i>Pecteilis sagarikii</i> seedlings during culturing for 12 weeks.	49
49	Main effect of illumination on number of root, root width and root length of <i>Pecteilis sagarikii</i> seedlings after culturing for 12 weeks.	49

Table	page
50	Main effect of sucrose concentrations on number of root, root width and root length of <i>Pecteilis sagarikii</i> seedlings after culturing for 12 weeks. 50
51	Effects of illumination and sucrose concentrations on number of root, root width and root length of <i>Pecteilis sagarikii</i> seedlings after culturing for 12 weeks. 50
52	Effects of illumination and sucrose concentrations on total soluble sugar in various tissues of <i>Pecteilis sagarikii</i> seedlings after culturing for 12 weeks. 52
53	Effects of illumination and sucrose concentrations on starch contents in various tissues of <i>Pecteilis sagarikii</i> seedlings after culturing for 12 weeks. 52
54	Changing of fresh weight (FW), dry weight (DW) and water content percentage (WC) in tuber, shoot and root in different growing stages of <i>Pecteilis sagarikii</i> grown in a green house. 68
55	Concentrations of nitrogen, phosphorus, potassium, calcium and magnesium in a tuber and shoot in different growing stages of <i>Pecteilis sagarikii</i> grown in a green house. 69
56	Concentrations of free amino acids and distribution in <i>Pecteilis sagarikii</i> tuber in different growing stages. 70
57	Concentrations of free amino acids and distribution in <i>Pecteilis sagarikii</i> shoot in different growing stages. 71
58	Concentrations of free amino acids and distribution in <i>Pecteilis sagarikii</i> root in different growing stages. 72
59	Concentrations of free indole-3-acetic acid (IAA), abscisic acid (ABA), and their distributions in <i>Pecteilis sagarikii</i> tuber and shoot in different growing stages. 73

LIST OF FIGURES

Figure		page
1	<i>Pecteilis sagarikii</i> Seidenfaden inflorescence.	4
2	<i>Pecteilis sagarikii</i> seedlings showing two different tuber shapes and the new tuber primordia (arrow).	20
3	<i>Pecteilis sagarikii</i> seedlings cultured in CMU ₁ medium with different sucrose concentrations at 0 1 2 3 5 and 7 % (w/v) for 10 weeks.	21
4	<i>Pecteilis sagarikii</i> seedlings cultured in CMU ₁ medium with different coconut water concentrations after culturing for 10 weeks.	25
5	<i>Pecteilis sagarikii</i> seedlings cultured in various supplemented CMU ₁ medium after culturing for 10 weeks.	31
6	<i>Pecteilis sagarikii</i> seedling cultured on CMU ₁ medium supplemented with coconut water at 15 % and BAP at 2.0 mg/l produced protocorm-like bodies (arrow) at the joining position between shoot and its protocorm.	34
7	<i>Pecteilis sagarikii</i> seedlings cultured in various tested medium in continuous darkness after culturing for 10 weeks.	39
8	Longitudinal cut of <i>Pecteilis sagarikii</i> seedlings after staining with iodine solution, showing starch accumulation in different parts.	43
9	<i>Pecteilis sagarikii</i> seedlings cultured in two light conditions.	51
10	The longitudinal free-hand sections of <i>Pecteilis sagarikii</i> seedlings staining with iodine solution showing starch accumulation in their different plant parts.	53
11	Various stages of <i>Pecteilis sagarikii</i> development observed under a stereomicroscope, 20 weeks after seed sowing.	55
12	<i>Pecteilis sagarikii</i> showed various stages of seedlings in a culture vessel, 20 weeks after seed sowing.	55
13	A longitudinal section of <i>Pecteilis sagarikii</i> seedling showing a tuber primordium in the early stage of tuberization.	56
14	A longitudinal section of <i>Pecteilis sagarikii</i> seedling showing a tuber primordium began to grow downwards.	57
15	A longitudinal section of <i>Pecteilis sagarikii</i> seedling showing a tuber stalk extension and a shoot bud formation.	58
16	A longitudinal section of <i>Pecteilis sagarikii</i> seedling by using a freezing-microtome and staining with iodine solution showing a tuber expansion and starch accumulation in different parts.	59
17	The magnified tuber cells showing various sizes of starch granules.	59
18	A tuber of <i>Pecteilis sagarikii</i> in the early sprouting stage.	60
19	The vegetative stage of <i>Pecteilis sagarikii</i> .	61

Figure	page
20	The pre-flowering stage of <i>Pecteilis sagarikii</i> . 61
21	The reproductive stage of <i>Pecteilis sagarikii</i> . 62
22	A new tuber produced a tuber-shoot bud (arrow) and continued to increase in the tuber size. 62
23	A young floral spike and a new tuber continued to increase in sizes. 63
24	A whole plant showing a complete new-formed tuber (arrow). 63
25	A longitudinal section of a vegetative shoot showing a tuber primordial bud originated at its lowest part of the plant monopodial stem. 64
26	A longitudinal section of a tuber primordium showing the extension of a tuber primordial meristem and leaf primordia. 65
27	A longitudinal section of a new tuber showing the formation of a new tuber-shoot bud. 66
28	A longitudinal section of a new tuber using a freezing microtome. 67
29	A longitudinal section of a seedling showing the formation of a new tuber primordium. 80
30	A longitudinal section of protocorm showed the multiple formation of apical meristematic dots. 81