

## TABLE OF CONTENTS

	<b>Page</b>
<b>Acknowledgements</b>	iii
<b>Abstract in English</b>	v
<b>Abstract in Thai</b>	ix
<b>Table of contents</b>	xii
<b>List of tables</b>	xv
<b>List of figures</b>	xviii
<b>Chapter 1 Introduction</b>	1
<b>Chapter 2 Literature review</b>	4
2.1 General information of genus <i>Phalaenopsis</i>	4
2.2 Classification of genus <i>Phalaenopsis</i>	6
2.3 Crossability in genus <i>Phalaenopsis</i>	10
2.4 Molecular markers for genetic analysis	12
<b>Chapter 3 Materials and Methods</b>	24
3.1 Analysis of genetic relationship of genus <i>Phalaenopsis</i> and related genera, <i>Doritis</i> and <i>Kingidium</i> , by RAPD technique	24
3.2 Studies on crossability of genus <i>Phalaenopsis</i> and related genera, <i>Doritis</i> and <i>Kingidium</i>	36

	<b>Page</b>
3.3 Characterizations of F <sub>1</sub> progenies derived from intersectional and intergeneric hybrids of <i>Phalaenopsis</i> and related genera, <i>Doritis</i> and <i>Kingidium</i> , by RAPD technique	40
3.4 Specific marker for flower color pattern of <i>Phalaenopsis cornu-cervi</i> by AFLP technique	41
<b>Chapter 4 Results</b>	<b>48</b>
4.1 Analysis of genetic relationship of genus <i>Phalaenopsis</i> and related genera, <i>Doritis</i> and <i>Kingidium</i> , by RAPD technique	48
4.2 Studies on crossability of genus <i>Phalaenopsis</i> and related genera, <i>Doritis</i> and <i>Kingidium</i>	64
4.3 Characterizations of F <sub>1</sub> progenies derived from intersectional and intergeneric hybrids of <i>Phalaenopsis</i> and related genera, <i>Doritis</i> and <i>Kingidium</i> , by RAPD technique	70
4.4 Specific marker for flower color pattern of <i>Phalaenopsis cornu-cervi</i> by AFLP technique	95
<b>Chapter 5 Discussions</b>	<b>100</b>
5.1 Analysis of genetic relationship of genus <i>Phalaenopsis</i> and related genera, <i>Doritis</i> and <i>Kingidium</i> , by RAPD technique	100
5.2 Studies on crossability of genus <i>Phalaenopsis</i> and related genera, <i>Doritis</i> and <i>Kingidium</i>	105

	<b>Page</b>
5.3 Characterizations of F <sub>1</sub> progenies derived from intersectional and intergeneric hybrids of <i>Phalaenopsis</i> and related genera, <i>Doritis</i> and <i>Kingidium</i> , by RAPD technique	108
5.4 Specific marker for flower color pattern of <i>Phalaenopsis</i> <i>cornu-cervi</i> by AFLP technique	111
<b>Chapter 6 Conclusions</b>	116
<b>References</b>	121
<b>Appendices</b>	130
Appendix A Chemical reagents preparation	131
Appendix B Data matrices of present (1) and absent (0) DNA band	135
<b>Curriculum vitae</b>	153

## LIST OF TABLES

Table		Page
1	Geographical distributions of plant materials used in this study.	25
2	Flower descriptions of plant materials used in this study.	26
3	List of 20 decamer primers and their sequences used for RAPD technique.	35
4	Intersectional hybridization of <i>Phalaenopsis</i> species.	38
5	Intrasectional hybridization of <i>Phalaenopsis</i> species.	39
6	Intergeneric hybridization of genus <i>Phalaenopsis</i> and related genera, <i>Doritis</i> and <i>Kingidium</i> .	39
7	List of primers and their sequences used for AFLP technique.	46
8	Sixty-four primer combinations used for AFLP technique.	47
9	Number of DNA bands of 36 samples from 30 species of <i>Phalaenopsis</i> and 2 related genera, <i>Doritis</i> and <i>Kingidium</i> , generated by OPAK10, OPD03, OPF01, OPF02, OPF09 and OPF14 primers.	55
10	Gene diversity and number of polymorphic loci among 8 sections of <i>Phalaenopsis</i> and 2 related genera, <i>Doritis</i> and <i>Kingidium</i> .	56

<b>Table</b>		<b>Page</b>
11	Genetic distance of 36 samples from 30 species of <i>Phalaenopsis</i> (PH01 - PH29) and 2 related genera, <i>Doritis</i> (D01 - D03) and <i>Kingidium</i> (K01 - K04), based on 6 primer combinations, OPAK10, OPD03, OPF01, OPF02, OPF09 and OPF14.	60
12	Genetic distance of 8 sections of <i>Phalaenopsis</i> and 2 related genera, <i>Doritis</i> and <i>Kingidium</i> , based on 6 primer combinations, OPAK10, OPD03, OPF01, OPF02, OPF09 and OPF14.	63
13	Results of intersectional hybridization of <i>Phalaenopsis</i> species.	66
14	Results of intrasectional hybridization of <i>Phalaenopsis</i> species.	67
15	Results of intergeneric hybridization of genus <i>Phalaenopsis</i> and related genera, <i>Doritis</i> and <i>Kingidium</i> .	67
16	Number of pollinated flowers and number of fruit setting from intersectional, intrasectional and intergeneric hybridization of genus <i>Phalaenopsis</i> and related genera, <i>Doritis</i> and <i>Kingidium</i> .	68
17	Number of seedlings per fruit obtained <i>in vitro</i> from 24 crosses.	69
18	Number of transplanted seedlings and number of survival plantlets after 6 months transplanting of 7 crosses.	69
19	Leaf descriptions of <i>P. schilleriana</i> (FP), <i>P. cornu-cervi</i> (MP) and their ten progenies (H1 - H10).	71
20	Flower descriptions of <i>P. schilleriana</i> (FP), <i>P. cornu-cervi</i> (MP) and their ten progenies (H1 - H10).	71

<b>Table</b>		<b>Page</b>
21	Leaf descriptions of <i>D. pulcherrima</i> ‘dwarf’ (FP), <i>P. equestris</i> (MP) and their ten progenies (H1 - H10).	73
22	Flower descriptions of <i>D. pulcherrima</i> ‘dwarf’ (FP), <i>P. equestris</i> (MP) and their ten progenies (H1 - H10).	74
23	Leaf descriptions of <i>D. pulcherrima</i> ‘dwarf’ (FP), <i>K. minus</i> (MP) and their ten progenies (H1 - H10).	76
24	Flower descriptions of <i>D. pulcherrima</i> ‘dwarf’ (FP), <i>K. minus</i> (MP) and their ten progenies (H1 - H10).	77

## LIST OF FIGURES

Figure		Page
1	Flower of <i>Phalaenopsis</i> species in section <i>Phalaenopsis</i> .	27
2	Flower of <i>Phalaenopsis</i> species in section <i>Proboscidioides</i> .	27
3	Flower of <i>Phalaenopsis</i> species in section <i>Parishianae</i> .	28
4	Flower of <i>Phalaenopsis</i> species in section <i>Polychilos</i> .	28
5	Flower of <i>Phalaenopsis</i> species in section <i>Stauroglottis</i> .	28
6	Flower of <i>Phalaenopsis</i> species in section <i>Fuscatae</i> .	29
7	Flower of <i>Phalaenopsis</i> species in section <i>Amboinenses</i> .	29
8	Flower of <i>Phalaenopsis</i> species in section <i>Zebrinae</i> .	29
9	Flower of species from genus <i>Doritis</i> .	30
10	Flower of species from genus <i>Kingidium</i> .	30
11	Three groups of <i>Phalaenopsis cornu-cervi</i> used in the AFLP analysis, reddish brown flower (a - c), yellow flower with reddish brown bar and spot (d - g), and pure yellow flower (no spot or bar) (h - i).	44
12	RAPD profiles of PH01 - PH18 generated by OPAK10 primer.	49
13	RAPD profiles of PH19 - D03 generated by OPAK10 primer.	49
14	RAPD profiles of PH01 - PH18 generated by OPD03 primer.	50
15	RAPD profiles of PH19 - D03 generated by OPD03 primer.	50
16	RAPD profiles of PH01 - PH18 generated by OPF01 primer.	51

<b>Figure</b>		<b>Page</b>
17	RAPD profiles of PH19 - D03 generated by OPF01 primer.	51
18	RAPD profiles of PH01 - PH18 generated by OPF02 primer.	52
19	RAPD profiles of PH19 - D03 generated by OPF02 primer.	52
20	RAPD profiles of PH01 - PH18 generated by OPF09 primer.	53
21	RAPD profiles of PH19 - D03 generated by OPF09 primer.	53
22	RAPD profiles of PH01 - PH18 generated by OPF14 primer.	54
23	RAPD profiles of PH19 - D03 generated by OPF14 primer.	54
24	Dendrogram of 36 samples from 30 species of <i>Phalaenopsis</i> and 2 related genera, <i>Doritis</i> and <i>Kingidium</i> , based on Nei' s (1972) genetic distance clustering by the UPGMA method from 6 primer combinations.	62
25	Dendrogram of 8 sections of <i>Phalaenopsis</i> and 2 related genera, <i>Doritis</i> and <i>Kingidium</i> , based on Nei' s (1972) genetic distance clustering by the UPGMA method from 6 primer combinations.	63
26	Flowers of <i>P. schilleriana</i> (FP), <i>P. cornu- cervi</i> (MP) and their ten progenies (H1 - H10).	72
27	Flowers of <i>D. pulcherrima</i> 'dwarf' (FP), <i>P. equestris</i> (MP) and their ten progenies (H1 - H10).	75
28	Flowers of <i>D. pulcherrima</i> 'dwarf' (FP), <i>K. minus</i> (MP) and their ten progenies (H1 - H10).	78



<b>Figure</b>		<b>Page</b>
29	RAPD profiles obtained from <i>P. schilleriana</i> (FP), <i>P. cornu-cervi</i> (MP) and their ten progenies (H1 - H10) after amplification with OPAK10 primer.	82
30	RAPD profiles obtained from <i>P. schilleriana</i> (FP), <i>P. cornu-cervi</i> (MP) and their ten progenies (H1 - H10) after amplification with OPD03 primer.	82
31	RAPD profiles obtained from <i>P. schilleriana</i> (FP), <i>P. cornu-cervi</i> (MP) and their ten progenies (H1 - H10) after amplification with OPF01 primer.	83
32	RAPD profiles obtained from <i>P. schilleriana</i> (FP), <i>P. cornu-cervi</i> (MP) and their ten progenies (H1 - H10) after amplification with OPF02 primer.	83
33	RAPD profiles obtained from <i>P. schilleriana</i> (FP), <i>P. cornu-cervi</i> (MP) and their ten progenies (H1 - H10) after amplification with OPF09 primer.	84
34	RAPD profiles obtained from <i>P. schilleriana</i> (FP), <i>P. cornu-cervi</i> (MP) and their ten progenies (H1 - H10) after amplification with OPF14 primer.	84
35	Principle component analysis (PCA) diagram illustrated the genetic relationship of <i>P. schilleriana</i> (FP), <i>P. cornu-cervi</i> (MP) and their ten progenies (H1 - H10) evaluated with 6 primer combinations, OPAK10, OPD03, OPF01, OPF02, OPF09 and OPF14.	85

<b>Figure</b>	<b>Page</b>
36	87
RAPD profiles obtained from <i>D. pulcherrima</i> ‘dwarf’ (FP), <i>P. equestris</i> (MP) and their ten progenies (H1 - H10) after amplification with OPAK10 primer.	
37	87
RAPD profiles obtained from <i>D. pulcherrima</i> ‘dwarf’ (FP), <i>P. equestris</i> (MP) and their ten progenies (H1 - H10) after amplification with OPF01 primer.	
38	88
RAPD profiles obtained from <i>D. pulcherrima</i> ‘dwarf’ (FP), <i>P. equestris</i> (MP) and their ten progenies (H1 - H10) after amplification with OPF02 primer.	
39	88
RAPD profiles obtained from <i>D. pulcherrima</i> ‘dwarf’ (FP), <i>P. equestris</i> (MP) and their ten progenies (H1 - H10) after amplification with OPF09 primer.	
40	89
Principle component analysis (PCA) diagram illustrated the genetic relationship of <i>D. pulcherrima</i> ‘dwarf’ (FP), <i>P. equestris</i> (MP) and their ten progenies (H1 - H10) evaluated with 4 primer combinations, OPAK10, OPF01, OPF02 and OPF09.	
41	91
RAPD profiles obtained from <i>D. pulcherrima</i> ‘dwarf’ (FP), <i>K. minus</i> (MP) and their ten progenies (H1 - H10) after amplification with OPAK10 primer.	

<b>Figure</b>		<b>Page</b>
42	RAPD profiles obtained from <i>D. pulcherrima</i> ‘dwarf’ (FP), <i>K. minus</i> (MP) and their ten progenies (H1 - H10) after amplification with OPD03 primer.	92
43	RAPD profiles obtained from <i>D. pulcherrima</i> ‘dwarf’ (FP), <i>K. minus</i> (MP) and their ten progenies (H1 - H10) after amplification with OPF02 primer.	92
44	RAPD profiles obtained from <i>D. pulcherrima</i> ‘dwarf’ (FP), <i>K. minus</i> (MP) and their ten progenies (H1 - H10) after amplification with OPF09 primer.	93
45	RAPD profiles obtained from <i>D. pulcherrima</i> ‘dwarf’ (FP), <i>K. minus</i> (MP) and their ten progenies (H1 - H10) after amplification with OPF14 primer.	93
46	Principle component analysis (PCA) diagram illustrated the genetic relationship of <i>D. pulcherrima</i> ‘dwarf’ (FP), <i>K. minus</i> (MP) and their ten progenies (H1 - H10) valuated with 5 primer combinations, OPAK10, OPD03, OPF02, OPF09 and OPF14.	94
47	AFLP profiles of <i>P. cornu-cervi</i> generated by 64 primer combinations.	97
48	AFLP profiles of 12 <i>P. cornu-cervi</i> generated by 4 <i>Eco</i> RI and <i>Mse</i> I primer combinations.	98
49	DNA sequences of two specific DNA bands of <i>P. cornu-cervi</i> .	99