

## TABLE OF CONTENTS

|   | <b>Page</b> |
|---|-------------|
| <b>Acknowledgements</b>   | iii         |
| <b>Abstract (English)</b>   | v           |
| <b>Abstract (Thai)</b>  | viii        |
| <b>Table of Contents</b>  | xi          |
| <b>List of Tables</b>   | xiii        |
| <b>List of Figures</b>  | xvi         |
| <b>List of Appendices</b>   | xxviii      |
| <b>List of Publications</b>   | xxx         |
| <br>  |             |
| <b>Chapter 1</b> Introduction   | 1           |
| <b>Chapter 2</b> Biodiversity and taxonomy  | 36          |
| <b>Chapter 3</b> The genus <i>Oxydothis</i> and its phylogenetic relationship<br>within the Xylariales                  | 150         |
| <b>Chapter 4</b> Morphology and phylogenetic study of leaf spotting<br>fungal species associated with terrestrial palms | 174         |
| <b>Chapter 5</b> Phylogenetic relationship of <i>Dictyochoeta wallichianensis</i><br>within Chaetosphaeriaceae          | 202         |
| <b>Chapter 6</b> Fungal community variation on decaying fronds of Thai Dwarf<br>Fishtail Palm                           | 218         |

**TABLE OF CONTENTS (CONTINUED)**

|   | <b>Page</b> |
|---|-------------|
| <b>Chapter 7</b> Analysis of vertical spatial distribution of fungal community<br>on Thai Dwarf Fishtail Palm | 256         |
| <b>Chapter 8</b> Endophytic and epiphytic fungi of Thai Dwarf Fishtail Palm                                   | 289         |
| <b>Chapter 9</b> Conclusions  | 314         |
| <b>References</b>   | 320         |
| <b>Appendices</b>   | 352         |
| <b>Curriculum Vitae</b>   | 403         |
| <b>Publications</b>   | 408         |

## LIST OF TABLES

| Table   | Page |
|---|------|
| 1.1 The most commonly Ascomycetes genera identified from palms in Brunei, Ecuador, and Malaysia during 1988-1994  | 27   |
| 1.2 List of the most common Ascomycetes genera encountered on palms   | 28   |
| 2.1 List of palmicolous fungi found on <i>Wallichia siamensis</i> and other palms during the present study        | 38   |
| 3.1 Comparison of <i>Oxydothis wallichianensis</i> with similar <i>Oxydothis</i> species                          | 163  |
| 3.2 Comparison of <i>Oxydothis inaequalis</i> with <i>O. asymmetrica</i>  | 164  |
| 3.3 Comparison of <i>Oxydothis</i> sp. FIH 019 with similar <i>Oxydothis</i> species                              | 164  |
| 3.4 Comparison of <i>Oxydothis cyrtostachicola</i> with J- <i>Oxydothis</i> Species                               | 165  |
| 3.5 Morphological comparison of <i>Oxydothis</i> with Amphisphaeriaceae and Hyponectriaceae                       | 172  |
| 4.1 Morphological comparison of <i>Cercospora arecacearum</i> with other <i>Cercospora</i> species from Arecaceae | 183  |

## LIST OF TABLES (CONTINUED)

| <b>Table</b>   | <b>Page</b> |
|--|-------------|
| 6.1 Two tail <i>t</i> -test tables showing comparison of mean data between fungal diversity indices on pinna and primary rachis of <i>Wallichia siamensis</i>  | 239         |
| 6.2 Two tail <i>t</i> -test tables showing comparison of mean data between fungal diversity indices on pinna and secondary rachis of <i>Wallichia siamensis</i>  | 240         |
| 6.3 Two tail <i>t</i> -test tables showing comparison of mean data between fungal diversity indices on primary and secondary rachis of <i>Wallichia siamensis</i>  | 241         |
| 6.4 Percentage indicator values (IndVal percentage) for all taxa with frequency of occurrence (FO) > 25%   | 244         |
| 6.5 A proportion of fungal community diversity on various species of palms   | 250         |
| 7.1 Two tail <i>t</i> -test tables showing comparison of mean data between Shanon-Weiner ( $H'$ ) fungal diversity indices on hanging-fallen fronds and primary-secondary rachis of <i>Wallichia siamensis</i> | 262         |
| 8.1 Isolation rate of fungal community on <i>Wallichia siamensis</i> at five different site based on tissue types and combination of tissue types-method (experimental design)                                 | 296         |

**LIST OF TABLES (CONTINUED)**

| <b>Table</b>  | <b>Page</b> |
|---|-------------|
| 8.2 Table showing Sørensen index of similarity among different type of microhabitat   | 305         |
| 8.3 Descriptive, test of homogeneity of variances, and ANOVA tables generated from One-Way ANOVA analysis showing the statistical test of isolation rate of surface disinfected and non-surface disinfected samples | 311         |

## LIST OF FIGURES

| Figure   | Page |
|--|------|
| 1.1 Morphology structure of Thai Dwarf Fishtail Palm<br>( <i>Wallichia siamensis</i> ) | 3    |
| 1.2 Structure of several economically important palm seeds                             | 7    |
| 1.3 General morphology structure of palms  | 10   |
| 1.4 Structures of palm leaves  | 11   |
| 1.5 Palm growth habits   | 13   |
| 1.6 A climbing palm, <i>Calamus</i> sp.  | 13   |
| 1.7 Illustration map of early Cretaceous or late Jurassic period                       | 16   |
| 1.8 Illustration map of late Cretaceous period   | 18   |
| 1.9 Illustration of Wallace's line in Malay Archipelago                                | 19   |
| 2.1 Micrographs of <i>Anthostomella limitata</i> on <i>Wallichia siamensis</i>         | 60   |
| 2.2 Micrographs of <i>Anthostomella puiggarii</i> on <i>Wallichia siamensis</i>        | 63   |
| 2.3 Micrographs of <i>Apiospora siamicola</i> <b>sp. nov.</b> (from holotype)          | 65   |
| 2.4 Micrographs of <i>Arecomyces foliicola</i> <b>sp. nov.</b> (from holotype)         | 69   |
| 2.5 Micrographs of <i>Botryosphaeria obtusa</i> on <i>Wallichia siamensis</i>          | 73   |
| 2.6 Micrographs of <i>Chaetospermum chaetosporum</i><br>on <i>Wallichia siamensis</i>  | 75   |
| 2.7 Micrographs of <i>Chaetosphaeria fusiformis</i> on <i>Wallichia siamensis</i>      | 77   |

## LIST OF FIGURES (CONTINUED)

| Figure   | Page |
|--|------|
| 2.8 Micrographs of <i>Fasciatispora petrakii</i> on <i>Wallichia siamensis</i>             | 79   |
| 2.9 Micrographs of <i>Glomerella cingulata</i> on <i>Wallichia siamensis</i>               | 81   |
| 2.10 Micrographs of <i>Guignardia uniappendiculatum</i> <b>sp. nov.</b><br>(from holotype) | 83   |
| 2.11 Line drawings of <i>Mycosphaerella wallichiae</i> sp. nov.<br>(from holotype)         | 86   |
| 2.12 Micrographs of <i>Myelosperma tumidum</i> on <i>Wallichia siamensis</i>               | 88   |
| 2.13 Micrographs of <i>Neorehmia arecae</i> on <i>Wallichia siamensis</i>                  | 91   |
| 2.14 Micrographs of <i>Ophioceras tenuisporum</i> on <i>Wallichia siamensis</i>            | 93   |
| 2.15 Micrographs of <i>Oxydothis daemonoropsicola</i> on<br><i>Wallichia siamensis</i>     | 95   |
| 2.16 Micrographs of <i>Oxydothis inaequalis</i> (from holotype)                            | 97   |
| 2.17 Micrographs of <i>Oxydothis wallichianensis</i> (from holotype)                       | 99   |
| 2.18 Micrographs of <i>Pemphidium rattanicola</i> on <i>Wallichia siamensis</i>            | 101  |
| 2.19 Micrographs of <i>Pestalospaeria elaeidis</i> on <i>Wallichia siamensis</i>           | 103  |
| 2.20 Micrographs of <i>Pestalotiopsis guepinii</i> on <i>Wallichia siamensis</i>           | 105  |
| 2.21 Micrographs of <i>Protocreopsis pertusa</i> on <i>Wallichia siamensis</i>             | 107  |

## LIST OF FIGURES (CONTINUED)

| Figure   | Page |
|--|------|
| 2.22 Micrographs of <i>Pseudohalonectria palmicola</i> on<br><i>Wallichia siamensis</i>    | 109  |
| 2.23 Micrographs of <i>Rachidicola obclavatum</i> <b>sp. nov.</b> (from holotype)          | 111  |
| 2.24 Micrographs of <i>Rosellinia victoriae</i> on <i>Wallichia siamensis</i>              | 113  |
| 2.25 Micrographs of <i>Submersisphaeria suthepensis</i> <b>sp. nov.</b><br>(from holotype) | 115  |
| 2.26 Micrographs of <i>Terriera brevis</i> on <i>Wallichia siamensis</i>                   | 117  |
| 2.27 Micrographs of <i>Apiosporina rhapsicola</i> <b>sp. nov.</b> (from holotype)          | 119  |
| 2.28 Micrographs of <i>Astrosphaeriella caryotae</i> <b>sp. nov.</b> (from holotype)       | 121  |
| 2.29 Micrographs of <i>Astrosphaeriella palmicola</i> <b>sp. nov.</b> (from holotype)      | 124  |
| 2.30 Line drawings of <i>Fasciatispora ujungkulonensis</i> (from holotype)                 | 126  |
| 2.31 Line drawings of <i>Guignardia arengae</i> <b>sp. nov.</b> (from holotype)            | 130  |
| 2.32 Micrographs of <i>Linocarpon nonappendiculatum</i> <b>sp. nov.</b><br>(from holotype) | 132  |
| 2.33 Micrographs of <i>Lophiostoma livistonocola</i> <b>sp. nov.</b> (from holotype)       | 135  |
| 2.34 Line drawings of <i>Mycosphaerella arecacearum</i> <b>sp. nov.</b><br>(from holotype) | 137  |
| 2.35 Line drawings of <i>Mycosphaerella borassi</i> <b>sp. nov.</b> (from holotype)        | 139  |



## LIST OF FIGURES (CONTINUED)

| <b>Figure</b>   | <b>Page</b> |
|---|-------------|
| 2.36 Line drawings of <i>Oedocephalum formosus</i> <b>sp. nov.</b> (from holotype)  | 142         |
| 2.37 Micrographs of <i>Oxydothis cyrtostachicola</i> (from holotype)  | 144         |
| 2.38 Micrographs of <i>Terriera livistonae</i> <b>sp. nov.</b> (from holotype)  | 146         |
| 2.38 Micrographs of <i>Venturia frondicola</i> <b>sp. nov.</b> (from holotype)  | 148         |
| 3.1 Micrographs of <i>Oxydothis wallichianensis</i> (from the holotype)   | 156         |
| 3.2 Micrographs of <i>Oxydothis inaequalis</i> (from holotype)  | 158         |
| 3.3 Micrographs of <i>Oxydothis cyrtostachicola</i> (from holotype)   | 161         |
| 3.4 Phylogenetic tree based on partial 28S nrDNA sequence data<br>representing placement of members of the <i>Oxydothis</i><br>within class Sordariomycetes                       | 166         |
| 3.5 Phylogenetic tree based on ITS nrDNA sequence data representing<br>placement of members of the <i>Oxydothis</i> within the Xylariales   | 167         |
| 4.1 <i>Cercospora arecearum</i> (from holotype)   | 180         |
| 4.2 Single parsimonious tree based on ITS nrDNA sequence data<br>representing placement of <i>Cercospora arecearum</i> within<br>representatives of the family Mycosphaerellaceae | 186         |
| 4.3 Line drawings of <i>Fusarium sansainensis</i> <b>sp. nov.</b> (from holotype)   | 188         |

## LIST OF FIGURES (CONTINUED)

| <b>Figure</b>   | <b>Page</b> |
|---|-------------|
| 4.4 Micrographs of <i>Fusarium sansainensis</i> <b>sp. nov.</b> (from holotype)   | 189         |
| 4.5 The best parsimonious tree based on ITS nrDNA sequence data representing placement of <i>Fusarium sansainensis</i> within representatives of the family Nectriaceae               | 193         |
| 5.1 Micrographs of <i>Dictyochaeta wallichianensis</i> <b>sp. nov.</b> (from holotype)  | 208         |
| 5.2 Micrographs of <i>Dictyochaeta wallichianensis</i> <b>sp. nov.</b> on PDA (from holotype)   | 209         |
| 5.3 The best parsimonious tree based on ITS nrDNA sequence data representing placement of <i>Dictyochaeta wallichianensis</i> within representatives of the family Chaetosphaeriaceae | 211         |
| 6.1 Species area curve of fungal number identified against number of samples examined   | 227         |
| 6.2 Dominance diversity curve showing the proportion of fungal abundance on different frond types of <i>Wallichia siamensis</i>   | 228         |
| 6.3 Distribution of all taxa calculated from total specimens of <i>Wallichia siamensis</i>  | 229         |
| 6.4 Distribution of all taxa recorded from primary rachis of <i>Wallichia siamensis</i>   | 230         |

## LIST OF FIGURES (CONTINUED)

| <b>Figure</b>  | <b>Page</b> |
|--|-------------|
| 6.5 Distribution of all taxa recorded from secondary rachis of <i>Wallichia siamensis</i>  | 231         |
| 6.6 Distribution of all taxa recorded from pinna of <i>Wallichia siamensis</i>   | 232         |
| 6.7 Histogram of Shannon-Weiner diversity index (H') and evenness index (E') showing the diversity and evenness levels of fungal community on different tissue types of <i>Wallichia siamensis</i> | 233         |
| 6.8 Histogram of Shannon-Weiner diversity index (H') and evenness index (E') showing the diversity and evenness levels of fungal community on different sites                                      | 234         |
| 6.9 Box plot showing diversity level of fungal community on total samples collected of <i>Wallichia siamensis</i>  | 235         |
| 6.10 Box plots showing proportion of diversity level of fungal community on samples collected from different tissue types of <i>Wallichia siamensis</i>  | 235         |
| 6.11 Box plot showing abundance of fungal community on total samples collected of <i>Wallichia siamensis</i>   | 236         |
| 6.12 Box plots showing abundance of fungal community on samples collected from different tissue types of <i>Wallichia siamensis</i>  | 237         |

## LIST OF FIGURES (CONTINUED)

| <b>Figure</b>  | <b>Page</b> |
|--|-------------|
| 6.13 Histogram showing similarity of fungal community occurs on different tissue types (as microhabitat) of <i>Wallichia siamensis</i>   | 238         |
| 6.14 Dendrogram for hierarchical clustering generated using Ward method and rescaled distance cluster combine matrix of all taxa with $\Sigma$ records > 10 from total samples collected | 242         |
| 6.15 Dendrogram representing hierarchical clustering (using Centroid method) of the pooled data from all specimens collected at five sites   | 243         |
| 6.16 Box plots showing distribution of <i>Ophioceras tenuisporum</i> on <i>Wallichia siamensis</i> based on tissue types   | 245         |
| 6.17 Box plots showing distribution of <i>Lophiostoma macrostomum</i> on <i>Wallichia siamensis</i> based on tissue types  | 246         |
| 6.18 Box plots showing distribution of <i>Ophioceras tenuisporum</i> on <i>Wallichia siamensis</i> based on tissue types   | 247         |
| 6.19 Correspondence Analysis (CA) showing a relationship between taxa (FO > 10%) and tissue types (as microhabitat)  | 248         |
| 6.20 Proportion number of fungi overlapping among different tissue types of <i>Wallichia siamensis</i>   | 254         |

## LIST OF FIGURES (CONTINUED)

| <b>Figure</b>   | <b>Page</b> |
|---|-------------|
| 7.1 Species area curve of number of fungi identified against number of samples examined   | 262         |
| 7.2 Proportion all taxa on fronds of <i>Wallichia siamensis</i> grouped based on pseudo-systematic position of a taxon                    | 263         |
| 7.3 Distribution of all taxa on <i>Wallichia siamensis</i> based on frequency of occurrence data (FO > 10)                                | 265         |
| 7.4 Qualitative distribution of all taxa recorded on <i>Wallichia siamensis</i> based on experimental design                              | 265         |
| 7.5 Quantitative distribution of all taxa recorded on <i>Wallichia siamensis</i> based on experimental design                             | 266         |
| 7.6 Qualitative distribution of all taxa recorded on <i>Wallichia siamensis</i> based on tissue types                                     | 266         |
| 7.7 Quantitative distribution of all taxa recorded on <i>Wallichia siamensis</i> based on tissue types                                    | 267         |
| 7.8 Qualitative distribution of all taxa recorded on <i>Wallichia siamensis</i> based on combination of experimental design-tissue types  | 268         |
| 7.9 Quantitative distribution of all taxa recorded on <i>Wallichia siamensis</i> based on combination of experimental design-tissue types | 268         |

## LIST OF FIGURES (CONTINUED)

| <b>Figure</b>   | <b>Page</b> |
|---|-------------|
| 7.10 Average number of record per sample all taxa recorded on <i>Wallichia siamensis</i> based on experimental design   | 269         |
| 7.11 Average number of record per sample all taxa recorded on <i>Wallichia siamensis</i> based on tissue types  | 270         |
| 7.12 Average number of record per sample all taxa recorded on <i>Wallichia siamensis</i> based on combination of experimental design-tissue types                                     | 271         |
| 7.13 Dominance diversity curve showing the proportion of fungal abundance on <i>Wallichia siamensis</i> based on combination of experimental design-tissue types as microhabitat      | 272         |
| 7.14 Two dimensional of Multi Dimensional Scaling (MDS) ordination of all taxa with frequency of occurrence > 10% based on total abundances and Euclidean distance model              | 274         |
| 7.15 Dendrogram for hierarchical clustering (using Ward method) of all taxa with $\Sigma$ records > 10 from total samples collected based on rescaled distance cluster combine matrix | 275         |

**LIST OF FIGURES (CONTINUED)**

| <b>Figure</b>   | <b>Page</b> |
|---|-------------|
| 7.16 Correspondence Analysis (CA) of all taxa with frequency of occurrence > 10% and combination of experimental design-tissue types (as microhabitat)                    | 276         |
| 7.17 Contingency table of Correspondence Analysis (CA) showed relationship between taxa and microhabitats (represented by combination of experimental design-tissue type) | 277         |
| 7.18 Box plots showing distribution of dominant species on <i>Wallichia siamensis</i> after one year decomposition based on experimental design and tissue types          | 278         |
| 7.19 Box plots showing proportion of diversity level of fungal community on <i>Wallichia siamensis</i> grouped based on experimental design and tissue types              | 280         |
| 7.20 Box plots showing Margalef's index species richness of fungal community of <i>Wallichia siamensis</i> based on combination of experimental design and tissue types   | 280         |
| 7.21 Box plots showing abundances of fungal community of <i>Wallichia siamensis</i> based on experimental design and tissue types   | 281         |

## LIST OF FIGURES (CONTINUED)

| <b>Figure</b>  | <b>Page</b> |
|--|-------------|
| 7.22 Graphic showing distribution of fungal community on hanging fronds of <i>Wallichia siamensis</i> based on species richness dataset                          | 282         |
| 7.23 Graphic showing distribution of fungal community on fallen fronds of <i>Wallichia siamensis</i> based on species richness dataset                           | 283         |
| 7.24 Graphic showing similarities among selected microhabitats based on Sørensen index analysis  | 284         |
| 7.25 The major biogeographic zones for microbial colonization of plant surfaces as a function of exposure and nutrient availability                              | 286         |
| 8.1 Bar errors showing mean of total fungal individual records based on tissue types as microhabitat   | 297         |
| 8.2 Distribution of all taxa on <i>Wallichia siamensis</i> based on abundances data ( $\Sigma$ records > 5)  | 298         |
| 8.3 Box plots showing the distribution of the most dominant species on <i>Wallichia siamensis</i> based on tissue types as microhabitat                          | 299         |
| 8.4 Bar errors showing mean of total fungal individual records based on tissue types and experimental design (surface disinfection and non-surface disinfection) | 300         |
| 8.5 Bar errors showing mean of total fungal individual records on different tissue types and site  | 301         |



**LIST OF FIGURES (CONTINUED)**

| <b>Figure</b>  | <b>Page</b> |
|--|-------------|
| 8.6 Graphic showing distribution of fungal community on <i>Wallichia siamensis</i> generated from non-surface disinfection method based on species abundances dataset  | 302         |
| 8.7 Graphic showing fungal community overlap generated from surface- disinfection and non surface- disinfection methods  | 302         |
| 8.8 Graphic showing distribution of endophytic fungal community on <i>Wallichia siamensis</i> based on species richness dataset  | 303         |
| 8.9 Graphic showing distribution of endophytic fungi community on <i>Wallichia siamensis</i> based on abundances dataset   | 304         |
| 8.10 Dendrogram for hierarchical clustering (using Ward method) of all taxa with frequency of occurrence > 5% from total samples collected based on rescaled distance cluster combine matrix                 | 307         |
| 8.11 Two dimensional of symmetric plot generated from multiple correspondence analysis showing the relationship among seven dominant taxa (represented by genus name), tissue types, and experimental design | 308         |

## LIST OF APPENDICES

| <b>Appendix</b>  | <b>Page</b> |
|--|-------------|
| 1 Sequences used in phylogenetic analysis of <i>Oxydothis</i> within Xylariales  | 352         |
| 2 Sequences used in the phylogenetic analysis of <i>Cercospora arecacearum</i> and related species   | 355         |
| 3 Sequences used in the analysis of <i>Fusarium sansainensis</i> and related species   | 357         |
| 4 Sequences used in the phylogenetic analysis of <i>Dictyochaeta wallichianensis</i>   | 360         |
| 5 List of all taxa found on fronds of <i>Wallichia siamensis</i> are shown with total number recorded, % Occurrence and % Abundance                              | 363         |
| 6 List of taxa recorded on <i>Wallichia siamensis</i> with $\Sigma$ records, % Occurrence, % Recurrence and % Abundance at each microhabitat and site            | 366         |
| 7 List of all taxa found on fronds of <i>Wallichia siamensis</i> are shown with total number recorded and % Abundance  | 377         |
| 8 List of taxa recorded on <i>Wallichia siamensis</i> with $\Sigma$ records, % Occurrence, % Recurrence and % Abundance at each microhabitat, treatment and site | 380         |
| 9 Frequency occurrence of all taxa recorded is represented based on various grouping of microhabitat   | 386         |

## LIST OF APPENDICES (CONTINUED)

| <b>Appendix</b>  | <b>Page</b> |
|--|-------------|
| 10 List of all taxa found on fronds of <i>Wallichia siamensis</i> are shown with total number recorded, % Abundance and % Occurrence                   | 390         |
| 11 List of taxa recorded on <i>Wallichia siamensis</i> with $\Sigma$ records, % Occurrence, % Recurrence and % Abundance at each microhabitat and site | 393         |

## LIST OF PUBLICATIONS

1. **Hidayat, I.**, Jeewoon, R., To-anun, C., and Hyde, K. D. (2006) The genus *Oxydothis*: new palmicolous taxa and phylogenetic relationship within *Xylariales*. *Fungal Diversity* **23**: 159-179.
2. **Hidayat, I.**, Meeboon, J., and To-anun, C. (2007). *Anthostomella* and *Fasciatispora* species (Xylariaceae) from palms including *F. ujungkulonensis* sp. nov. *Mycotaxon* **102**: 347–354.
3. **Hidayat, I.**, Meeboon, J., K.D. Hyde, and C. To-anun. (2007). A study on microfungi associated with necrotic leaves of palms. In: *Proceedings of the International Conference on 'Integration of Science and Technology for Sustainable Development* (Eds. Soytong, K. and Hyde K. D.), KMITL, Ladkrabang, Bangkok, Thailand, 26-27 April 2007. p. 279-285
4. To-anun, C., Nguenhom, J., Meeboon, J., and **Hidayat, I.** (2009). Two fungi associated with necrotic leaflets of areca palms (*Areca catechu*). *Mycological Progress* **8**: 115-121.