

## TABLE OF CONTENTS

|   | <b>Page</b> |
|---|-------------|
| <b>Acknowledgements</b>                           | iii         |
| <b>Abstract(Thai)</b>                             | v           |
| <b>Abstract(English)</b>                          | vii         |
| <b>List of Tables</b>                             | xiii        |
| <b>List of Figures</b>                            | xv          |
| <b>Abbreviations and symbols</b>                  | xvii        |
| <b>Chapter 1</b>                                  | 1           |
| Introduction                                      | 1           |
| <b>Chapter 2</b>                                  | 4           |
| Literature review                                 | 4           |
| 2.1 Crop improvement                              | 4           |
| 2.2 Hybrid rice                                   | 5           |
| 2.3 Haploid and doubled haploid plants production | 7           |
| 2.4 <i>In vitro</i> techniques                    | 8           |
| 2.4.1 Anther and pollen culture                   | 10          |
| 2.4.1.1 The benefit of anther culture             | 11          |
| 2.4.1.2 Modes of androgenesis                     | 13          |
| 2.4.1.3 Factors affect on androgenesis            | 13          |
| 2.4.1.4 Stage of the microspores and stress       | 14          |
| Pretreatment                                      |             |

|   |    |
|---|----|
| 2.4.2 Callus cultures   | 16 |
| 2.4.3 Cell suspension cultures  | 19 |
| 2.4.4 Embryogenesis in suspension cultures  | 21 |
| 2.4.4.1 Subculturing  | 22 |
| 2.4.4.2 Subculturing hazards  | 23 |
| 2.5 The components of plant tissue culture media                                      | 24 |
| 2.5.1 Macro- and Micro-nutrients  | 24 |
| 2.5.2 Growth regulators   | 26 |
| 2.6 Genetic stability   | 27 |
| 2.7 Doubling chromosome number  | 27 |
| 2.8 Somaclonal and gametoclonal variations  | 28 |
| 2.8.1 Sources of somaclonal variation   | 29 |
| 2.8.2 Factors determining somaclonal variation  | 29 |
| 2.8.3 Applications somaclonal variation in plant<br>breeding                          | 30 |
| 2.9 Chromosomal analysis of <i>in vitro</i> cultured tissue and<br>regenerated plants | 31 |
| 2.10 Auxin effect in tissue culture   | 32 |
| 2.10.1 Induction of callus growth   | 32 |
| 2.10.2 Organ cultures   | 33 |
| 2.10.3 Embryogenesis  | 34 |
| 2.11 Growth regulator shock   | 35 |
| 2.12 Synthetic seeds (Artificial seeds)   | 36 |
| 2.12.1 Advantages of synthetic seeds  | 40 |

|                  |  |           |
|------------------|--|-----------|
|                  | 2.12.2 Limitations of synthetic seeds                                    | 41        |
| <b>Chapter 3</b> | <b>Material and methods</b>  | <b>45</b> |
|                  | 3.1 Plant materials  | 47        |
|                  | 3.2 Chemical reagents  | 47        |
|                  | 3.2.1 <i>In vitro</i> culture media                                      | 47        |
|                  | 3.2.2 Haploid diploidization (Chromosome doubling)                       | 49        |
|                  | 3.2.3 Chromosome analysis  | 49        |
|                  | 3.2.4 Synthetic seed production  | 49        |
|                  | 3.3 Instrument   | 49        |
|                  | 3.4 Equipments   | 50        |
|                  | 3.5 Methods  | 51        |
|                  | 3.5.1 Pollen culture   | 51        |
|                  | 3.5.2 Caulogenesis inducement  | 52        |
|                  | 3.5.3 Hormone Shock, Doubling chromosome<br>and Embryogenesis inducement | 53        |
|                  | 3.5.4 Chromosome analysis  | 53        |
|                  | 3.5.5 Synthetic seed production  | 54        |
|                  | 3.5.6 Statistical Analysis   | 55        |
| <b>Chapter 4</b> | <b>Results and Discussion</b>  | <b>56</b> |
|                  | 4.1 Pre-Treatment  | 56        |
|                  | 4.2 Caulogenesis inducement  | 61        |
|                  | 4.3 Embryogenesis inducement and doubling<br>chromosome                  | 65        |
|                  | 4.4 Synthetic seed production  | 75        |

|                         |   |            |
|-------------------------|---|------------|
|                         | 4.5 Germination and conversion of encapsulated<br>embryoids | 81         |
| <b>Chapter 5</b>        | <b>Conclusions</b>  | <b>83</b>  |
| <b>References</b>       |   | <b>85</b>  |
| <b>Appendices</b>       |   | <b>105</b> |
| <b>Appendix A</b>       |   | <b>106</b> |
| <b>Appendix B</b>       |   | <b>114</b> |
| <b>Appendix C</b>       |   | <b>115</b> |
| <b>Appendix D</b>       |   | <b>117</b> |
| <b>Appendix E</b>       |   | <b>123</b> |
| <b>Curriculum vitae</b> |   | <b>131</b> |

## LIST OF TABLES

| <b>Table</b> |   | <b>Page</b> |
|--------------|---|-------------|
| 4.1          | Effect of low temperature treatment on development stage of F1 pollen. <sup>a</sup>   | 59          |
| 4.2          | Effect of low temperature treatment on development stage of H1 pollen. <sup>a</sup>   | 60          |
| 4.3          | Effect of low temperature treatment on F1 and H1 androgenesis in LS media. <sup>a</sup>   | 61          |
| 4.4.         | Influence of the various LS media formulas on the anther culture response of F1 hybrid rice. <sup>a</sup>   | 70          |
| 4.5.         | Influence of the various LS media formulas on the anther culture response of H1 hybrid rice. <sup>a</sup>   | 71          |
| 4.6.         | Influence of the various LS media formulas on the caulogenesis of F1 and H1 anther hybrid Rice. <sup>a</sup>  | 72          |
| 4.7          | Influence of LS liquid media supplemented with different concentration of colchicine and 2,4-D on the embryogenesis of F1 hybrid rice. <sup>a</sup> | 73          |
| 4.8          | Effect of different colchicine and 2,4-D concentrations on chromosome doubling in F1 hybrid rice anther culture. <sup>a</sup>                       | 74          |

- 4.9 Result of storage time on inducing desiccation tolerance synthetic seed germination percentage and speed of germination after dehydration was persued until seeds lost 80% of their moisture contents 78
- 4.10 Effect of different benomyl concentration of encapsulating applied to embryoid of KDML 105 x SPR 1 (H1) 80

## LIST OF FIGURES


| <b>Figure</b> |   | <b>Page</b> |
|---------------|---|-------------|
| 4.1           | Stage of microspore development of hybrid rice (a) early uninucleate (b) mid- uninucleate (c) late- uninucleate (d) binucleate.   | 58          |
| 4.2           | The difference of callus induction from anther on LS media formula no.10 after 4 weeks of culture (a) F1 hybrid anther and (b) H1 anther. The difference of organogenesis formation in anther culture on LS media formula no.4 promoted organogenesis in F1 anther culture (c) and in H1 anther culture (d) after 4 weeks of culture. Embryogenic callus formation (e) and plant regeneration (f) from anther culture after 4 weeks of culture. | 69          |
| 4.3           | Differentiation of embryoids after culture calli in LS media supplemented 0.2 g /L colchicine and 100 $\mu$ M 2,4-D (a). compare with conventional anther culture method (b) after 8 weeks.   | 69          |
| 4.4           | Analyzed metaphase chromosome of ELSs after (a). treated with 0.2 g /L colchicine and 100 $\mu$ M 2,4-D and (b) treated with over 0.3 % colchicines (denature chromosome).  | 69          |
| 4.5           | Synthetic seed (a). during dehydration persued until the seeds lost 80% and (b) compare size of Synthetic seed before and after dehydration.  | 79          |

- 4.6 Synthetic seed (a). during dehydration persued until the seeds lost 80% and (b) compare size of Synthetic seed before and after dehydration. 79
- 4.7 F1 hybrid seed germination compare with H1 synthetic seed (a). 4 days, (b) 14 days and synthetic seeds after culturing on LS media 14 days (c). 82



### ABBREVIATIONS AND SYMBOLS

|       |   |                                |
|-------|---|--------------------------------|
| ABA   | = | Absciscic acid                 |
| BA    | = | 6-benzylaminopurine            |
| °C    | = | Degree of Celsius              |
| 2,4-D | = | 2,4-Dichlorophenoxyacetic acid |
| cm    | = | Centimeter                     |
| cv.   | = | Cultivar                       |
| DW    | = | Dry weight                     |
| EDTA  | = | Ethylenediaminetetraacetate    |
| e.g.  | = | Exempli gratia (for example)   |
| g     | = | Gram                           |
| GA    | = | Gibberellic acid               |
| hrs   | = | Hours                          |
| IAA   | = | Indole-3-acetic acid           |
| IBA   | = | Indole-3-butyric acid          |
| i.e.  | = | Id est (it is or that is)      |
| L     | = | Liter                          |
| LS    | = | Linsmaier and Skoog            |
| mg    | = | Milligram                      |
| mg/L  | = | Milligram per liter            |
| ml    | = | Milliliter                     |



|         |   |                                  |
|---------|---|----------------------------------|
| mm      | = | Millimeter                       |
| mM      | = | Millimole                        |
| MS      | = | Murashige and Skoog              |
| NAA     | = | $\alpha$ -Naphthaleneacetic acid |
| rpm     | = | Revolutions per minute           |
| UV      | = | Ultraviolet (light)              |
| v/v     | = | volume/volume (concentration)    |
| w/v     | = | weight/volume (concentration)    |
| %       | = | Percent                          |
| $\mu$ M | = | Micromole                        |

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่

Copyright© by Chiang Mai University  
All rights reserved