

TABLE OF CONTENTS

	PAGE
ACKNOWLEDGEMENT	iii
ABSTRACT	v
TABLE OF CONTENTS	ix
LIST OF TABLES	xi
LIST OF FIGURES	xii
ABBREVIATIONS	xiv
1. INTRODUCTION	1
2. OBJECTIVES	15
3. LITERATURE REVIEWS	16
4. MATERIALS AND METHODS	
4.1 Equipment	21
4.2 Sample used in the studies	21
4.3 Methods	22
4.3.1 Urinary microalbumin	22
4.3.2 Urinary NAG activity	23
4.3.3 HbA _{1C} determination	23
4.3.4 Fructosamine	24
4.3.5 Blood urea nitrogen (BUN)	24
4.3.6 Creatinine	25
4.3.7 Glucose	25
4.3.8 Urinalysis	25

4.4 Method of studies	26
4.4.1 Precision studies	26
4.4.2 Linearity studies of MA and NAG	26
4.4.3 Accuracy studies	26
4.4.4 Stability of BPB and CNP-NAG reagents	27
4.4.5 Clinical values of urinary MA and NAG determinations.	28
4.4.6 Normal reference values	28
4.4.7 Correlation of the tests	29
5. RESULTS	
5.1 Precision of methods .	30
5.2 Linearity studies of MA and NAG determination methods.	30
5.3 Accuracy studies	30
5.4 Stability of BPB and CNP-NAG reagents	31
5.5 Correlation of the tests	31
5.6 Cut-off predictive levels of MA and NAG	32
5.7 Clinical values of urinary MA and NAG determinationms.	32
5.8 Reference values	33
6. DISCUSSION	53
7. REFERENCES	60
8. APPENDIX	68
9. CURRICULUM VITAE	77

LIST OF TABLES

TABLE	PAGE
1. Precision of methods used for serum glucose, BUN, creatinine and fructosamine determinations. (n=10)	34
2. Precision of the NAG activity determination using CNP-NAG as substrate. (n=10)	34
3. Precision of the urinary microalbumin determination with bromphenol blue (BPB) method. (n=10)	35
4. Reference ranges for urinary NAG activity, MA and serum fructosamine in 100 healthy adults	35
5. Analytical recovery (%) of standard NAG (64.52 U/l) activity added into urine sample. (Average of 4 determinations)	36
6. Analytical recovery (%) of standard MA (100 mg/dl) added into urine sample. (Average of 4 determinations)	36
7. Sensitivity, specificity of urinary MA (cut-off 2.97 mg/gm creat.), and NAG activity (cut-off 32.4 U/gm creat.)	37
8. Positive and negative predictive values of urinary MA (cut-off 2.97 mg/gm creat.), NAG activity (cut-off 32.4 U/gm creat.) in diabetics at different prevalences .	37
9. Sensitivity, specificity of urinary MA (cut-off 3.48 mg/gm creat.), and NAG activity (cut-off 37.2 U/gm creat.) in diabetics (using ROC curve for cut-off level determination).	38
10. Positive and negative predictive value of urinary MA (cut-off 3.48 mg/gm creat.), NAG activity (cut-off 37.2 U/gm creat.) in diabetics at different prevalences (using ROC curve for cut-off level determination).	38

LIST OF FIGURES

FIGURE	PAGE
1. Distribution of glucose and fructosamine in within-run (A) and between-run (B) precision studies.	39
2. Distribution of BUN and creatinine in within-run (A) and between-run (B) precision studies.	40
3. Distribution of normal and high levels MA in within-run (A) and between-run (B) precision studies.	41
4. Distribution of normal and high levels of NAG activity in within-run (A) and between-run (B) precision studies.	42
5. Calibration curve of standard human albumin.	43
6. Calibration curve of standard NAG.	43
7. Stability of bromphenol blue reagent. (n=4)	44
8. Stability of CNP-NAG reagent. (n=4)	45
Correlation of MA and NAG activity in diabetics. (n=220)	45
9. Correlation of MA and NAG activity in patients with diabetic nephropathy. (n=14)	45
10. Correlation of MA an HbA _{1C} in diabetics. (n=220)	46
11. Correlation of NAG and HbA _{1C} in diabetics. (n=220)	46
12. Correlation of MA an HbA _{1C} in patients with diabetic nephropathy. (n=14)	47
13. Correlation of NAG activity an HbA _{1C} in patients with diabetic nephropathy. (n=14)	47
14. Correlation of MA and fructosamine in diabetics. (n=220)	48
15. Correlation of NAG activity and fructosamine in diabetics. (n=220)	48
16. Correlation of MA and fructosamine in patients with diabetic nephropathy. (n=14)	49
18. Correlation of NAG activity and fructosamine in patients with diabetic nephropathy. (n=14)	49
19. Correlation of fructosamine and HbA _{1C} in diabetics. (n=220)	50
20. Distribution of NAG activity in normal control group (n=100) and diabetics (n=220).	51

20. Distribution of MA in normal control group (n=100) and diabetics (n=220).	51
21. Receiver operating characteristics curve (ROC curve) of MA level in diabetics. (n=220)	52
23. Receiver operating characteristics curve (ROC curve) of MA level in diabetics. (n=220)	52

มหาวิทยาลัยเชียงใหม่
Chiang Mai University

ABBREVIATIONS

α -KG	Alpha ketoglutarate
°C	Degree Celcius
%CV	Percentage coefficient of variation
BPB	Bromphenol blue
CHD	Coronary heart disease
CNP-NAG	2-Chloro-4-nitrophenyl-N-acetyl- β -D- glucosaminide
DKA	Diabetic ketoacidosis
DM	Diabetes mellitus
DMF	1-Deoxy-1-morpholinofructose
DN	Diabetic nephropathy
ESRD	End stage renal disease
G-6-P	Glucose-6-phosphate
G-Hb	Glycosylated hemoglobin
GLDH	Glutamate dehydrogenase
gm.	Gram
gm creat.	Gram creatinine
Hb	Hemoglobin
HLA	Human Lenkocyte Antigens
IDDM	Insulin-dependent diabetes mellitus
MA	Microalbumin
NADH	Nicotinamide adenine dinucleotide
NAG	N-Acetyl- β -D-glucosaminidase
NBT	Nitroblue tetrazolium
NHHC	Non-ketotic hyperglycemic hyperosmotic coma

NIDDM	Non-insulin-dependent diabetes mellitus
r	Coefficient of correlation
ROC	Receiver operating characteristics
U/l	Unit per litre

มหาวิทยาลัยเชียงใหม่
Chiang Mai University