

REFERENCES

1. Weedt W., Harrison M. (1985). Measuring recovery of arm-hand function in stroke patient: A comparison of the Brunnstrom-Fugl-Meyer test and the Action Research Arm test. *Physiother Can*, 37, 65-70.
2. Duncan P., Goldstein L., Horner R., Landsma., Samsa G., Matchar D. (1994). Similar motor recovery of upper and lower extremities after stroke. *Stroke*, 25, 1181-8.
3. Olsen T. (1990). Arm and leg paresis as outcome predictors in stroke rehabilitation. *Stroke*, 21, 247-51.
4. Winstein C., Rose D., Chui H., (2001). Recovery and rehabilitation of arm use after stroke. *J Stroke Cerebrovasc Dis*, 10, 197.
5. Nakayama H., Jorgenson H., Raaschou H., Olson T. (1994). Recovery of upper extremity function in stroke patients: The Copenhagen stroke study. *Arch Phys Med Rehabil*, 75, 394-8.
6. Ernst E. (1990). A review of stroke rehabilitation and physiotherapy. *Stroke* , 21, 1081-5.
7. Coot S., Stokes E. (2001). Physiotherapy for upper extremity dysfunction following stroke. *Phys Ther Rev*, 75, 394-8.

8. Wade D., Hewer R., Wood V., Skilbeck C., Ismail H. (1983). The hemiplegic arm after stroke: measurement and recovery. *J Neurol Neurosurg Psychiatry*, 46, 521-4.
9. Kalra L., Dale P., Crome P. (1993). Improving stroke rehabilitation: A controlled study. *Stroke*, 24, 1462-7.
10. Blanton S., Wolf S. (1999). An application of upper extremity constraint-induced movement therapy in a patient with subacute stroke. *Phys Ther*, 79, 847- 53.
11. Taub E., Carlson J., Seifert A., Birbaumer N. (1994). Overcoming learned non-use: A new approach to treatment in physical medicine. *Clinical Applied Psychophysiology*. Plenum Press:185-220.
12. Carr J., Shepherd R. (1987). A motor relearning programme for stroke: 2nd edition. Oxford, UK: Heinemann Medical.
13. Consensus panel on the management of the hemiplegic arm and hand. (2002). Hamilton. Ontario, Canada: The Heart and Stroke Foundation of Ontario.
14. Butefisch C., Hummelsheim H., Denzler P., Mauritz KH. (1995). Repetitive training of isolated movements improves the outcome of motor rehabilitation of the centrally paretic hand. *J Neurol Sci*, 130, 59-68.
15. Feys H.M., De Weerd W.J., Selz B.E., Cox Steck G.A., Spichiger R., Vereeck L.E., et al. (1998). Effect of a therapeutic intervention for the hemiplegic upper limb in the acute phase after stroke: a single-blind, randomized, controlled multicenter trial. *Stroke*, 29, 785-92.

16. Whittall J., McCombe W.S., Silver K.H., Macko R.F. (2000). Repetitive bilateral arm training with rhythmic auditory cueing improves motor function in chronic hemiparetic stroke. *Stroke*, 31, 2390-5.
17. Mudie M.H., Matyas T.A. (1996). Upper extremity retraining following stroke: Effects of bilateral practice. *J Neurologic Rehabil*, 10, 167-84.
18. Van der Lee J., Wagenaar R., Lankhorst G. (1999). Forced use of the upper extremity in chronic stroke patients results from a single-blind randomized clinical trial. *Stroke*, 2369-75.
19. Wolf S., Lecraw D., Barton L., Jann B. (1989). Forced use of hemiparetic upper extremities to reverse the effect of learned non-use among chronic stroke and head-injured patient. *Exp Neurol*, 107, 125-32.
20. Dromerick A.W., Edwards D.F., Hahn M. (2000). Does the application of constraint-induced movement therapy during acute rehabilitation reduce arm impairment after ischemic stroke? *Stroke*, 31, 2984-8.
21. Sunderland A., Tinson D.J., Bradley E.L., et al. (1992). Enhanced physical therapy improves recovery of arm function after stroke: A randomized controlled trial. *J Neurol Neurosurg Psychiatry*, 55, 530-5.
22. Dickstein R., Hocherman S., Pillar T., Shaham R. (1988). Stroke rehabilitation. Three exercise therapy approaches. *Phys Ther Rev*, 66,1233-8.

23. Gelber D.A., Josefczyk P.B., Herrman D., Good D.C., Verhulst S.J. (1995). Comparison of two therapy approaches in the rehabilitation of the pure motor hemiparetic stroke patient. *J Neurologic Rehabil*, 9,191-6.
24. Logigian M.K., Samuels M.A., Falconer J., Zagar R. (1983). Clinical exercise trial for stroke patients. *Arch Phys Med Rehabil*, 64, 364-7.
25. Liepert J, Miltner WHR, Bauder H, Sommer M, Dettmers C, Taub E, Weiller C. (1998). Motor cortex plasticity during constraint-induced movement therapy in stroke patients. *Neurosci Lett*, 250, 5-8.
26. Eng J.J., Chu K.S., Kim C.M., Dawson A.S., Carswell A., Mepburn K.E. (2003). Community-based groups exercise program for persons with chronic stroke. *Med Sci Sports Exerc*, 35, 1271-3.
27. Brandsma J.W., Lakerveld-Heyl K., Van Ravensberg C.D., Heerkens Y.F. (1995). Reflection on the definition of impairment and disability as defined by the world health organization, 17(3-4), 119-27.
28. Wolf S.L., Lcatlin P.A., Ellis M., et al. (2001). Assessing Wolf motor function test as outcome measure for research in patients after stroke. *Stroke*, 32, 1635-9.
29. Uswatte G., Taub E. Constraint-induced movement therapy: New approaches to outcome measurement in rehabilitation. In Stuss D.T., Winocur G., Robertson I.H., editors. (1999). *Cognitive neurorehabilitation: A comprehensive approach*. Cambridge: Cambridge University Press.

30. Hatano S. (1976). Experience from a multicentre stroke register: A preliminary report. *Bulletin of the World Health Organization*, 54, 541-53
31. Caplan L.R. (2000). *Caplan's stroke: A clinical approach*, 3rd edition. Butterworth-Heinemann.
32. Gillen G. (2004). *Stroke rehabilitation: A functional-based approach*, 2nd edition. Columbia University. New York, 190-4.
33. Glenn M.B., Whyte J. (1990). *The practical management of spasticity in children and adults*. Philadelphia: Lea & Febiger.
34. Preston L.A., Hecht J.S. (1999). *Spasticity management: Rehabilitation strategies*. Bethesda: American Occupational Therapy Association.
35. Sahrman S.A., Norton B.J. (1977). The relationship of voluntary movement to spasticity in the upper motor neuron syndrome. *Ann Neurol*, 2, 460-5.
36. Fellows S.J., Kaus C., Thilmann A.F. (1994). Voluntary movement at the elbow in spastic hemiparesis. *Ann Neurol*, 36, 397-407.
37. Hummelsheim H., Munch B., Butefisch C, et al. (1994). Influence of sustained stretch on late muscular responses to magnetic brain stimulation in patients with upper motor neuron lesions. *Scand J Rehabil Med*, 26, 3-9.
38. Perry J. Rehabilitation of spasticity. In Feldman RG, Young RR, Koella WP, editors. (1980). *Spasticity: Disordered motor control*, Chicago.
39. Bourbonnais D., Vanden NS (1989). Weakness in patients with hemiparesis. *Am J Occup Ther*, 43, 313-9.

40. Bohannon R.W., Larkin P.A., Smith M.B., et al. (1987). Relationship between static muscle strength deficits and spasticity in stroke patients with hemiparesis. *Phys Ther*, 67, 1068-71.
41. Gowland C., deBruin H., Basmaijian J.V., et al. (1992). Agonist and antagonist activity during voluntary upper-limb movement in patients with stroke. *Phys Ther*, 72, 624-33.
42. Winstein C.J., Rose D.K., Tan S.M., Lewthwaite R., Chui H.C., Azen S.P. (2004). A randomized controlled comparison of upper-extremity rehabilitation strategies in acute stroke: A pilot study of immediate and long-term outcomes. *Arch Phys Med Rehabil*, 85, 620-8.
43. Badics E., Wittmann A., Rupp M., Stabauer B., Zifko U.A. (2002). Systematic muscle building exercises in the rehabilitation of stroke patients. *J Neurologic Rehabil*, 17, 211-14.
44. Kapandji I.A. (1982). *The physiology of the joints, vol 1, The upper limb*. New York: Churchill Livingstone.
45. Botte M.J., Nickel V.L., Akeson W.H. (1988). Spasticity and contracture: Physiologic aspects of formation. *Clin Orthop*, 233, 7-18.
46. Halar E.M., Bell K.R. Contracture and other deleterious effects of immobility. In DeLisa J.B., editor. (1993). *Rehabilitation medicine: Principles and practice*. Philadelphia.

47. Mathiowetz V., Bass-Haugen J. (1994). Motor behavior research: Implications for therapeutic approaches to central nervous system dysfunction. *Am J Occup Ther*, 48, 733-45.
48. Ada L., Canning C.G., Call J.H., et al. (1994). Task-specific training of reaching and manipulation. In Bennett KMB., Castiello U., editors. *Insights into the reach to grasp movements*, Amsterdam.
49. Sabari J.S. (1991). Motor learning concepts applied to activity-based intervention in adults with hemiplegia. *Am J Occup Ther*, 45, 523-30.
50. Jorgensen H.S., Nakayama H., Raaschou H.O., Vive-Larsen J., Stoier M., Olsen T.S. (1995b). Outcome and time course of recovery in stroke. Part I: Outcome. The Copenhagen stroke study. *Arch Phys Med Rehabil*, 76, 399-405.
51. Dromerick A.W., Reding M.J. (1995). Functional outcome for patients with hemiparesis, 8. hemihyesthesia and hemianopsia: does lesion location matter? *Stroke*, 26, 2023.
52. Rogers J.C., Holm M.B., Beach S., Shulz R. (2000). Comparison of ADL/IADL Assessment Methods. Annual Scientific Meeting of the Gerontological Society of America, Washington, DC, November 18.
53. Szekeley G. (1979). Order and plasticity in the nervous system. *Trend in Neuroscience*, 2, 245-49.
54. Goldberger M. (1980). Motor recovery after lesion. *Neuroscience*, 288-91.
55. Rosner B.S. (1970). Brain functions. *Ann.Rev.Psyh*, 21, 555-94.

56. Lashley K.S. (1979). Brain mechanisms and intelligence. Chicago: University of Chicago press.
57. Bliss T.V.P. (1979). Synaptic plasticity in the hippocampus. *Trend in Neuroscience*, 2, 42-5.
58. Laurence S., Stein D.G. (1978). Recovery after brain damage and the concept of localization of function. In *recovery from brain damage* (Finger S., ed). London: Plenum press, 369-407.
59. Luria A.R. (1963). Restoration of function after brain injury. London: Pergamon.
60. Andrews K., Brocklehurst J.C., Richards B., Laycock P.J. (1981). The rate of recovery from stroke-and its measurement. *Int Rehabil Med*, 3, 155-61.
61. Skilbeck C.E., Wade D.T., Hewer R.L. Wood V.A. (1983). Recovery after stroke. *J Neurol Neurosurg Psychiatry*, 46, 5-8.
62. Duncan P.W., Goldstein L.B., Horner R.D., Landsman P.B., Samsa G.P., Matchar D.B. (1994). Similar motor recovery of upper and lower extremities after stroke. *Stroke*, 25, 1181-88.
63. Kwakkel G., Kollen B.J., Wagenaar R.C. (1999). Therapy impact on functional recovery in stroke rehabilitation: a critical review of the literature. *Physiotherapy*, 13, 457-70.
64. Luria A.R. (1963). Restoration of function after brain injury. London: Pergamon.

65. American College of sports medicine. (1990). The recommended quantity and quality of exercise for developing and maintaining cardiorespiratory and muscular fitness in healthy adults. *Med Sci Sports Exerc*;22:265-74
66. Haennel R.G., Quinney H.A., Kappagoda C.T. (1990). Effects of hydraulic circuit training following coronary artery bypass surgery. *Med Sci Sports Exerc*, 23, 158-65
67. Wilmore J.H., Parr R.B., Girandola R.N. (1978). Physiological alternations consequent to circuit weight training. *Med Sci Sports*, 10, 79-84.
68. Bhambhani Y., Rowland G., Farag M. (2005). Effects of circuit training on body composition and peak cardiorespiratory response in patients with moderate to severe traumatic brain injury. *Arch Phys Med Rehabil*, 86, 268-76.
69. Carr J.H., Shepherd R.B. (1990). A motor learning model for rehabilitation of the movement-disabled. In: Ada L., Canning C., editors. *Key issues in neurological physiotherapy*. Oxford: Heinemann Medical, 1-24.
70. Pang M.Y., Harris J.E., Eng J.J. (2006). A community-based upper extremity group exercise program improves motor function and performance of functional activities in chronic stroke: A randomized controlled trial. *Arch Phys Med Rehabil*, 87, 1-9.
71. Dean C.M., Richards C.L., Malouin F. (2000). Task-related circuit training improves performance of locomotor tasks in chronic stroke: A randomized, controlled pilot trial. *Arch Phys Med Rehabil*, 81, 409-17.

72. Anderson B., Burke E.R. (1991). Scientific, medical and practical aspects of stretching. *Clin Sports Med*, 10, 63–86.
73. Bandy W.D., Irion J.M. (1994). The effect of time on static stretch on the flexibility of the hamstring muscles. *Phys Ther*, 74, 845–50.
74. Miltner W.L., Bauder H., Sommer M., Dettmers C., Taub E. (1999). Effects of constraint induced movement therapy on patients with chronic motor deficits after stroke. *Stroke*, 30, 586-92.
75. Taub E., Miller N.E., Novack T.A. (1993). Techniques to improve chronic motor deficit after stroke. *Arch Phys Med Rehabil*, 4, 347-54.
76. Volpe B.T., Kerbs H.I., Hogan N., Edelstein L., Diels C.M., Aisen M.L. (2000). A novel approach to stroke rehabilitation: Robot-aided sensorimotor stimulation. *Neurology*, 54, 1938-44.
77. Volpe B.T., Kerbs H.I., Hogan N. (2001). Is robot-aided sensorimotor training in stroke rehabilitation a realistic option? *Curr Opin Neurol*, 14, 745-52.
78. Lum P.S., Burgar C.G., Shor P.C., Majmundar M., Van der Loos M. (2002). Robot-assisted movement training compared with conventional therapy techniques for the rehabilitation of upper-limb motor function after stroke. *Arch Phys Med Rehabil*, 83, 952-9.
79. Cohen L. (1970). Interaction between limbs during bimanual voluntary activity. *Brain*, 93, 259–72.

80. Whitall J., Waller S., Silver K., Macko R. (2000). Repetitive bilateral arm training with rhythmic auditory cueing improves motor function in chronic hemiparetic stroke. *Stroke*, 31, 2390–5.
81. Poole J.L., Whitney S.L., Hangeland N et al. (1990). The effectiveness of inflatable pressure splints on motor function in stroke patients. *Occup Ther J Res*, 10, 360-6.
82. Langlois S., Pederson L., MacKinnon J.R. (1991). The effects of splinting on the spastic hemiplegic hand: Report of a feasible study. *Can J Occup Ther*, 58, 17-25.
83. Robichaud J.A., Agostinucci J., Vander Linden D.W. (1992). Effect of air-splint application on soleus muscle motoneuron reflex excitability in nondisabled subjects and subjects with cerebrovascular accidents. *Phys Ther*, 72, 176 –85.
84. Sherman S.J., Koshland G.F., Laguna J.F. (2000). Hyper-reflexia without spasticity after unilateral infarct of the medullary pyramid. *J Neurol Sci*, 175, 145-55.
85. McCrea P.H., Eng J.J., Hodgson A.J. (2003). Time and magnitude of torque generation is impaired in both arms following stroke. *Muscle Nerve*, 28, 46-53.
86. Pohl P.S., Duncan P., Perera S. (2002). Rate of isometric knee extension strength development and walking speed after stroke. *J Rehabil Res Dev*, 39, 651-8.
87. Canning C.G., Ada L, O'Dwyer N. (1999). Slowness to develop force contributes to weakness after stroke. *Arch Phys Med Rehabil*, 80, 66-70.

88. Moreland J.D., Goldsmith C.H., Huijbregts M.P. et al. (2003). Progressive resistance strengthening exercises after stroke: a single-blind randomized controlled trial. *Arch Phys Med Rehabil*, 84, 1433-40.
89. Inaba M., Edberg E., Montgomery J. et al. (1973). Effectiveness of functional training, active exercise, and resistive exercise for patients with hemiplegia. *Phys Ther*, 53, 28-35.
90. Patricia K., Sandra A. B. (2005). Exercise-induced changes of the upper extremity in chronic stroke survivors. *Stroke Rehabil*, 12, 58-68.
91. Cannine C.G., Ada L., Adams R., O'Dwyer N.J. (2004). Loss of strength contributes more to physical disability after stroke than loss of dexterity. *Clin Rehabil*, 18, 300-8.
92. Teixeira-Salmela L.F., Nadeau S., McBride I., Olney S.J. (2001). Effects of muscle strengthening and physical conditioning training on temporal, kinematic and kinetic variables during gait in chronic stroke survivors. *J Rehabil Med*, 33, 53-60.
93. Sharp S.A., Brouwer B.J. (1997). Isokinetic strength training of the hemiparetic knee: Effects on function and spasticity. *Arch Phys Med Rehabil*, 78, 1231-6.
94. Engardt M., Knutsson E., Jonsson M., Sternhag M. (1995). Dynamic muscle strength training in stroke patients: Effects on knee extension torque, electromyographic activity and motor function. *Arch Phys Med Rehabil*, 76, 419-25.

95. Fasoli S.E., Kerbs H.I., Stein J., Frontera W.R., Hogan N. (2003). Effects of robotic therapy on motor impairment and recovery in chronic stroke. *Arch Phys Med Rehabil*, 84, 477-82.
96. Patten C, McGill KC, Lateva Z, Rose A. (2000). Common drive among concurrently active motor units in cerebral palsy and post-stroke hemiparesis. 30th Ann Mtg Soc Neurosci.
97. Patten C., Lexell J., Brown H.E. (2004). Weakness and strength training in persons with post-stroke hemiplegia: Rationale, method, and efficacy. *J Rehabil Res Dev*, 41, 293-312.
98. Thilmann A.F., Fellows S.J., Garms E. (1991). The mechanism of spastic muscle hypertonus: Variation in reflex gain over the time course of spasticity. *Brain*, 114, 233-44.
99. Tsai K.H., Yeh C.Y., Chang H.Y., Chen J.J. (2001). Effects of a single session of prolonged muscle stretch on spastic muscle of stroke patients. *Proc Natl SCI Counc Repub China B*, 25, 76-81.
100. Hutchinson K., Winnegge T. (2004). Manual therapy applied to the hemiparetic upper extremity of a chronic poststroke individual: Case report. *J Neurol Phys Ther*, 28, 129-37.
101. Kluding P., Zipp G. (2004). Effect of ankle joint mobilization of ankle mobility and sit-to-stand in subjects with hemiplegia. *J Neurol Phys Ther*, 28, 72-83.

102. Borms J., Van R.P., Santans J.P. (1987). Optimal duration of static stretching exercises for improvement of coxo-femoral flexibility. *J Sports Sci*, 5, 39–47.
103. Roberts J.M., Wilson K. (1999). Effects of stretching duration on active and passive range of motion in the lower extremity. *Br J Sports Med*, 33, 259-63.
104. Ckassen J., Liepert J., Wise S.P., Hallett M., Cohen L.G. (1998). Rapid plasticity of human cortical movement representation induced by practical. *J Neurophysiol*, 79, 1117-23.
105. Taub E., Crago J.E., Uswatte G. (1998). Constraint-induced movement therapy: A new approach to treatment in physical rehabilitation. *Rehabil Psychol*, 43, 152-70.
106. Kwakkel G., Wagenaar R.C., Twisk J., Lankhorst G.J., Koetsier J.C. (1999). Intensity leg and arm training after primary middle-cerebral-artery stroke: A randomized trial. *The Lancet*, 354, 189-94.
107. Pascual-Leone A., Grafman J., Hallett M. (1994). Modulation of cortical motor output maps during development of implicit and explicit knowledge. *Science*, 263, 1287–9.
108. Elbert T., Rockstroh B. (2004). Reorganization of human cerebral cortex: The range of changes following use and injury. *The Neuroscientist*, 10, 129-41
109. Liepert J., Bauder H., Miltner W., Taub E., Weiller C. (2000). Treatment-induced cortical reorganization after stroke in humans. *Stroke*, 31, 1210-6.
110. Jacobs K.M., Donoghue J.P. (1991). Reshaping the cortical motor map by unmasking latent intracortical connections. *Science*, 251, 944 –7.

111. Langhammer B., Stanghelle J.K. (2000). Bobath or motor relearning programme? A comparison of two different approaches of physiotherapy in stroke rehabilitation: A randomized controlled study. *Clin Rehabil*, 14, 361-9.
112. Lennon S. (1996). The Bobath concept: a critical review of the theoretical assumptions that guide physiotherapy practice in stroke rehabilitation. *Phys Ther Rev*, 1, 35-45.
113. Fleiss J.L. (1986) cited in Delitto A., Strube M.J. (1991). Reliability in the clinical setting. *American Physical Therapy Association Research Section Newsletter* 24, 2-8.