

REFERENCES

- Abbas, A.K. and Lichtman, A.H. (2003) Cellular and Molecular Immunology. Saunders, The Curtis Center,Independent Square West, Philadelphia, Pennsylvania.
- Aslund, F. and Beckwith, J. (1999) The thioredoxin superfamily: redundancy, specificity, and gray-area genomics. *J Bacteriol* 181, 1375-9.
- Athappilly, F.K. and Hendrickson, W.A. (1995) Structure of the biotinyl domain of acetyl-coenzyme A carboxylase determined by MAD phasing. *Structure* 3, 1407-19.
- Bach, H., Mazor, Y., Shaky, S., Shoham-Lev, A., Berdichevsky, Y., Gutnick, D.L. and Benhar, I. (2001) Escherichia coli maltose-binding protein as a molecular chaperone for recombinant intracellular cytoplasmic single-chain antibodies. *J Mol Biol* 312, 79-93.
- Baneyx, F. and Mujacic, M. (2004) Recombinant protein folding and misfolding in *Escherichia coli*. *Nat Biotechnol* 22, 1399-408.
- Bardwell, J.C., McGovern, K. and Beckwith, J. (1991) Identification of a protein required for disulfide bond formation in vivo. *Cell* 67, 581-9.
- Beckett, D., Kovaleva, E. and Schatz, P.J. (1999) A minimal peptide substrate in biotin holoenzyme synthetase-catalyzed biotinylation. *Protein Sci* 8, 921-9.

- Bessette, P.H., Aslund, F., Beckwith, J. and Georgiou, G. (1999) Efficient folding of proteins with multiple disulfide bonds in the *Escherichia coli* cytoplasm. Proc Natl Acad Sci U S A 96, 13703-8.
- Biswas, C., Zhang, Y., DeCastro, R., Guo, H., Nakamura, T., Kataoka, H. and Nabeshima, K. (1995) The human tumor cell-derived collagenase stimulatory factor (renamed EMMPRIN) is a member of the immunoglobulin superfamily. Cancer Res 55, 434-9.
- Brodsky, F.M. and Guagliardi, L.E. (1991) The cell biology of antigen processing and presentation. Annu Rev Immunol 9, 707-44.
- Butler, J.E., Ni, L., Brown, W.R., Joshi, K.S., Chang, J., Rosenberg, B. and Voss, E.W., Jr. (1993) The immunochemistry of sandwich ELISAs--VI. Greater than 90% of monoclonal and 75% of polyclonal anti-fluorescyl capture antibodies (CAbs) are denatured by passive adsorption. Mol Immunol 30, 1165-75.
- Butt, T.R., Edavettal, S.C., Hall, J.P. and Mattern, M.R. (2005) SUMO fusion technology for difficult-to-express proteins. Protein Expr Purif 43, 1-9.
- Caudroy, S., Polette, M., Tournier, J.M., Burlet, H., Toole, B., Zucker, S. and Birembaut, P. (1999) Expression of the extracellular matrix metalloproteinase inducer (EMMPRIN) and the matrix metalloproteinase-2 in bronchopulmonary and breast lesions. J Histochem Cytochem 47, 1575-80.
- Chaiet, L. and Wolf, F.J. (1964) The Properties of Streptavidin, a Biotin-Binding Protein Produced by Streptomycetes. Arch Biochem Biophys 106, 1-5.
- Chapman-Smith, A. and Cronan, J.E., Jr. (1999) The enzymatic biotinylation of proteins: a post-translational modification of exceptional specificity. Trends Biochem Sci 24, 359-63.

- Chiampanichayakul, S., Szekeres, A., Khunkaewla, P., Moonsom, S., Leksa, V., Drbal, K., Zlabinger, G.J., Hofer-Warbinek, R., Stockinger, H. and Kasinrerk, W. (2002) Engagement of Na,K-ATPase beta3 subunit by a specific mAb suppresses T and B lymphocyte activation. *Int Immunol* 14, 1407-14.
- Creighton, T.E. (1997) Protein folding coupled to disulphide bond formation. *Biol Chem* 378, 731-44.
- Dalberg, K., Eriksson, E., Enberg, U., Kjellman, M. and Backdahl, M. (2000) Gelatinase A, membrane type 1 matrix metalloproteinase, and extracellular matrix metalloproteinase inducer mRNA expression: correlation with invasive growth of breast cancer. *World J Surg* 24, 334-40.
- De Becker, G., Moulin, V., Pajak, B., Bruck, C., Francotte, M., Thiriart, C., Urbain, J. and Moser, M. (2000) The adjuvant monophosphoryl lipid A increases the function of antigen-presenting cells. *Int Immunol* 12, 807-15.
- de Dios Alche, J. and Dickinson, H. (1998) Affinity chromatographic purification of antibodies to a biotinylated fusion protein expressed in Escherichia coli. *Protein Expr Purif* 12, 138-43.
- De Marco, V., Stier, G., Blandin, S. and de Marco, A. (2004) The solubility and stability of recombinant proteins are increased by their fusion to NusA. *Biochem Biophys Res Commun* 322, 766-71.
- Diamandis, E.P. and Christopoulos, T.K. (1991) The biotin-(strept)avidin system: principles and applications in biotechnology. *Clin Chem* 37, 625-36.
- Fadool, J.M. and Linser, P.J. (1993) 5A11 antigen is a cell recognition molecule which is involved in neuronal-glial interactions in avian neural retina. *Dev Dyn* 196, 252-62.

- Fall, R.R. and Vagelos, P.R. (1972) Acetyl coenzyme A carboxylase. Molecular forms and subunit composition of biotin carboxyl carrier protein. *J Biol Chem* 247, 8005-15.
- Fox, J.D., Kapust, R.B. and Waugh, D.S. (2001) Single amino acid substitutions on the surface of *Escherichia coli* maltose-binding protein can have a profound impact on the solubility of fusion proteins. *Protein Sci* 10, 622-30.
- Frazer, I.H., Quinn, M., Nicklin, J.L., Tan, J., Perrin, L.C., Ng, P., O'Connor, V.M., White, O., Wendt, N., Martin, J., Crowley, J.M., Edwards, S.J., McKenzie, A.W., Mitchell, S.V., Maher, D.W., Pearse, M.J. and Basser, R.L. (2004) Phase 1 study of HPV16-specific immunotherapy with E6E7 fusion protein and ISCOMATRIX adjuvant in women with cervical intraepithelial neoplasia. *Vaccine* 23, 172-81.
- Frenkel, D., Dori, M. and Solomon, B. (2004) Generation of anti-beta-amyloid antibodies via phage display technology. *Vaccine* 22, 2505-8.
- Fuh, G., Mulkerrin, M.G., Bass, S., McFarland, N., Brochier, M., Bourell, J.H., Light, D.R. and Wells, J.A. (1990) The human growth hormone receptor. Secretion from *Escherichia coli* and disulfide bonding pattern of the extracellular binding domain. *J Biol Chem* 265, 3111-5.
- Gerstner, J.A., Bell, J.A. and Cramer, S.M. (1994) Gibbs free energy of adsorption for biomolecules in ion-exchange systems. *Biophys Chem* 52, 97-106.
- Goldstein, M.A. and Doi, R.H. (1995) Prokaryotic promoters in biotechnology. *Biotechnol Annu Rev* 1, 105-28.
- Green, N.M. (1975) Avidin. *Adv Protein Chem* 29, 85-133.

- Guerder, S., Meyerhoff, J. and Flavell, R. (1994) The role of the T cell costimulator B7-1 in autoimmunity and the induction and maintenance of tolerance to peripheral antigen. *Immunity* 1, 155-66.
- Gupta, R.K., Chang, A.C. and Siber, G.R. (1998) Biodegradable polymer microspheres as vaccine adjuvants and delivery systems. *Dev Biol Stand* 92, 63-78.
- Gupta, R.K., Rost, B.E., Relyveld, E. and Siber, G.R. (1995) Adjuvant properties of aluminum and calcium compounds. *Pharm Biotechnol* 6, 229-48.
- Hefti, M.H., Van Vugt-Van der Toorn, C.J., Dixon, R. and Vervoort, J. (2001) A novel purification method for histidine-tagged proteins containing a thrombin cleavage site. *Anal Biochem* 295, 180-5.
- Henkart, P.A. (1994) Lymphocyte-mediated cytotoxicity: two pathways and multiple effector molecules. *Immunity* 1, 343-6.
- Hiniker, A. and Bardwell, J.C. (2003) Disulfide bond isomerization in prokaryotes. *Biochemistry* 42, 1179-85.
- Hochuli, E., Dobeli, H. and Schacher, A. (1987) New metal chelate adsorbent selective for proteins and peptides containing neighbouring histidine residues. *J Chromatogr* 411, 177-84.
- Hockney, R.C. (1994) Recent developments in heterologous protein production in *Escherichia coli*. *Trends Biotechnol* 12, 456-63.
- Hogg, N. and Landis, R.C. (1993) Adhesion molecules in cell interactions. *Curr Opin Immunol* 5, 383-90.
- Holmgren, A. and Aslund, F. (1995) Glutaredoxin. *Methods Enzymol* 252, 283-92.

- Holmgren, A. and Bjornstedt, M. (1995) Thioredoxin and thioredoxin reductase. *Methods Enzymol* 252, 199-208.
- Intasai, N., Arooncharus, P., Kasinrerk, W. and Tayapiwatana, C. (2003) Construction of high-density display of CD147 ectodomain on VCSM13 phage via gpVIII: effects of temperature, IPTG, and helper phage infection-period. *Protein Expr Purif* 32, 323-31.
- Johansen, P., Men, Y., Merkle, H.P. and Gander, B. (2000) Revisiting PLA/PLGA microspheres: an analysis of their potential in parenteral vaccination. *Eur J Pharm Biopharm* 50, 129-46.
- Jurado, P., Ritz, D., Beckwith, J., de Lorenzo, V. and Fernandez, L.A. (2002) Production of functional single-chain Fv antibodies in the cytoplasm of *Escherichia coli*. *J Mol Biol* 320, 1-10.
- Kanekura, T., Miyauchi, T., Tashiro, M. and Muramatsu, T. (1991) Basigin, a new member of the immunoglobulin superfamily: genes in different mammalian species, glycosylation changes in the molecule from adult organs and possible variation in the N-terminal sequences. *Cell Struct Funct* 16, 23-30.
- Kaplan, W., Husler, P., Klump, H., Erhardt, J., Sluis-Cremer, N. and Dirr, H. (1997) Conformational stability of pGEX-expressed *Schistosoma japonicum* glutathione S-transferase: a detoxification enzyme and fusion-protein affinity tag. *Protein Sci* 6, 399-406.
- Kapust, R.B. and Waugh, D.S. (1999) *Escherichia coli* maltose-binding protein is uncommonly effective at promoting the solubility of polypeptides to which it is fused. *Protein Sci* 8, 1668-74.

- Kasinrerk, W., Fiebiger, E., Stefanova, I., Baumruker, T., Knapp, W. and Stockinger, H. (1992) Human leukocyte activation antigen M6, a member of the Ig superfamily, is the species homologue of rat OX-47, mouse basigin, and chicken HT7 molecule. *J Immunol* 149, 847-54.
- Kasinrerk, W., Tokrasinwit, N. and Phunpae, P. (1999) CD147 monoclonal antibodies induce homotypic cell aggregation of monocytic cell line U937 via LFA-1/ICAM-1 pathway. *Immunology* 96, 184-92.
- Kersteen, E.A., Higgin, J.J. and Raines, R.T. (2004) Production of human prolyl 4-hydroxylase in *Escherichia coli*. *Protein Expr Purif* 38, 279-91.
- Khunkeawla, P., Moonsom, S., Staffler, G., Kongtawelert, P. and Kasinrerk, W. (2001) Engagement of CD147 molecule-induced cell aggregation through the activation of protein kinases and reorganization of the cytoskeleton. *Immunobiology* 203, 659-69.
- Kieffhaber, T., Rudolph, R., Kohler, H.H. and Buchner, J. (1991) Protein aggregation in vitro and in vivo: a quantitative model of the kinetic competition between folding and aggregation. *Biotechnology (N Y)* 9, 825-9.
- Koch, C., Staffler, G., Huttinger, R., Hilgert, I., Prager, E., Cerny, J., Steinlein, P., Majdic, O., Horejsi, V. and Stockinger, H. (1999) T cell activation-associated epitopes of CD147 in regulation of the T cell response, and their definition by antibody affinity and antigen density. *Int Immunol* 11, 777-86.
- Li, S.J. and Cronan, J.E., Jr. (1992) The gene encoding the biotin carboxylase subunit of *Escherichia coli* acetyl-CoA carboxylase. *J Biol Chem* 267, 855-63.
- Major, T.C., Liang, L., Lu, X., Rosebury, W. and Bocan, T.M. (2002) Extracellular matrix metalloproteinase inducer (EMMPRIN) is induced upon monocyte

differentiation and is expressed in human atheroma. *Arterioscler Thromb Vasc Biol* 22, 1200-7.

Marmorstein, A.D., Gan, Y.C., Bonilha, V.L., Finnemann, S.C., Csaky, K.G. and Rodriguez-Boulan, E. (1998) Apical polarity of N-CAM and EMMPRIN in retinal pigment epithelium resulting from suppression of basolateral signal recognition. *J Cell Biol* 142, 697-710.

Medzhitov, R. and Janeway, C., Jr. (2000) Innate immunity. *N Engl J Med* 343, 338-44.

Min, D.J., Andrade, J.D. and Stewart, R.J. (1999) Specific immobilization of in vivo biotinylated bacterial luciferase and FMN:NAD(P)H oxidoreductase. *Anal Biochem* 270, 133-9.

Miyauchi, T., Masuzawa, Y. and Muramatsu, T. (1991) The basigin group of the immunoglobulin superfamily: complete conservation of a segment in and around transmembrane domains of human and mouse basigin and chicken HT7 antigen. *J Biochem (Tokyo)* 110, 770-4.

Mogk, A., Mayer, M.P. and Deuerling, E. (2002) Mechanisms of protein folding: molecular chaperones and their application in biotechnology. *Chembiochem* 3, 807-14.

Morein, B. and Bengtsson, K.L. (1999) Immunomodulation by iscoms, immune stimulating complexes. *Methods* 19, 94-102.

Muramatsu, T. and Miyauchi, T. (2003) Basigin (CD147): a multifunctional transmembrane protein involved in reproduction, neural function, inflammation and tumor invasion. *Histol Histopathol* 18, 981-7.

- Nehme, C.L., Fayos, B.E. and Bartles, J.R. (1995) Distribution of the integral plasma membrane glycoprotein CE9 (MRC OX-47) among rat tissues and its induction by diverse stimuli of metabolic activation. *Biochem J* 310 (Pt 2), 693-8.
- Nikaido, H. (1994) Maltose transport system of *Escherichia coli*: an ABC-type transporter. *FEBS Lett* 346, 55-8.
- Peyre, M., Fleck, R., Hockley, D., Gander, B. and Sesardic, D. (2004) In vivo uptake of an experimental microencapsulated diphtheria vaccine following subcutaneous immunisation. *Vaccine* 22, 2430-7.
- Plotkin, S.A. and Orenstein, W.A. (2004) *Vaccines*. Saunders, The Curtis Center,Independent Square West, Philadelphia, Pennsylvania.
- Porath, J., Carlsson, J., Olsson, I. and Belfrage, G. (1975) Metal chelate affinity chromatography, a new approach to protein fractionation. *Nature* 258, 598-9.
- Rao, K.V., He, Y.X. and Kalyanasundaram, R. (2003) Expression of a 28-kilodalton glutathione S-transferase antigen of *Schistosoma mansoni* on the surface of filamentous phages and evaluation of its vaccine potential. *Clin Diagn Lab Immunol* 10, 536-41.
- Reed, K.E. and Cronan, J.E., Jr. (1991) *Escherichia coli* exports previously folded and biotinated protein domains. *J Biol Chem* 266, 11425-8.
- Rietsch, A. and Beckwith, J. (1998) The genetics of disulfide bond metabolism. *Annu Rev Genet* 32, 163-84.
- Rietsch, A., Belin, D., Martin, N. and Beckwith, J. (1996) An in vivo pathway for disulfide bond isomerization in *Escherichia coli*. *Proc Natl Acad Sci U S A* 93, 13048-53.

Robinson, A.S. and King, J. (1997) Disulphide-bonded intermediate on the folding and assembly pathway of a non-disulphide bonded protein. *Nat Struct Biol* 4, 450-5.

Rozkov, A. and Enfors, S.O. (2004) Analysis and control of proteolysis of recombinant proteins in *Escherichia coli*. *Adv Biochem Eng Biotechnol* 89, 163-95.

Salkowski, C.A., Detore, G.R. and Vogel, S.N. (1997) Lipopolysaccharide and monophosphoryl lipid A differentially regulate interleukin-12, gamma interferon, and interleukin-10 mRNA production in murine macrophages. *Infect Immun* 65, 3239-47.

Sambhara, S., Kurichh, A., Miranda, R., Tumpey, T., Rowe, T., Renshaw, M., Arpino, R., Tamane, A., Kandil, A., James, O., Underdown, B., Klein, M., Katz, J. and Burt, D. (2001) Heterosubtypic immunity against human influenza A viruses, including recently emerged avian H5 and H9 viruses, induced by FLU-ISCOM vaccine in mice requires both cytotoxic T-lymphocyte and macrophage function. *Cell Immunol* 211, 143-53.

Sanders, M.T., Brown, L.E., Deliyannis, G. and Pearse, M.J. (2005) ISCOM-based vaccines: the second decade. *Immunol Cell Biol* 83, 119-28.

Santala, V. and Lamminmaki, U. (2004) Production of a biotinylated single-chain antibody fragment in the cytoplasm of *Escherichia coli*. *J Immunol Methods* 284, 165-75.

Sato, K., Kawasaki, H., Nagayama, H., Enomoto, M., Morimoto, C., Tadokoro, K., Juji, T. and Takahashi, T.A. (2001) Signaling events following chemokine

- receptor ligation in human dendritic cells at different developmental stages. *Int Immunol* 13, 167-79.
- Schatz, P.J. (1993) Use of peptide libraries to map the substrate specificity of a peptide-modifying enzyme: a 13 residue consensus peptide specifies biotinylation in *Escherichia coli*. *Biotechnology (N Y)* 11, 1138-43.
- Scholle, M.D., Collart, F.R. and Kay, B.K. (2004) In vivo biotinylated proteins as targets for phage-display selection experiments. *Protein Expr Purif* 37, 243-52.
- Schumann, W. and Ferreira, L.C.S. (2004) Production of recombinant protein in *Escherichia coli*. *Genetics and Molecular Biology* 27, 442-453.
- Seulberger, H., Unger, C.M. and Risau, W. (1992) HT7, Neurothelin, Basigin, gp42 and OX-47--many names for one developmentally regulated immunoglobulin-like surface glycoprotein on blood-brain barrier endothelium, epithelial tissue barriers and neurons. *Neurosci Lett* 140, 93-7.
- Shenoy, B.C. and Wood, H.G. (1988) Purification and properties of the synthetase catalyzing the biotination of the aposubunit of transcarboxylase from *Propionibacterium shermanii*. *Faseb J* 2, 2396-401.
- Sibler, A.P., Kempf, E., Glacet, A., Orfanoudakis, G., Bourel, D. and Weiss, E. (1999) In vivo biotinylated recombinant antibodies: high efficiency of labelling and application to the cloning of active anti-human IgG1 Fab fragments. *J Immunol Methods* 224, 129-40.
- Singh, M. and O'Hagan, D. (1999) Advances in vaccine adjuvants. *Nat Biotechnol* 17, 1075-81.
- Smith, P.A., Tripp, B.C., DiBlasio-Smith, E.A., Lu, Z., LaVallie, E.R. and McCoy, J.M. (1998) A plasmid expression system for quantitative in vivo biotinylation


- of thioredoxin fusion proteins in *Escherichia coli*. Nucleic Acids Res 26, 1414-20.
- Solomon, B. (2005) Generation of anti-beta-amyloid antibodies via phage display technology towards Alzheimer's disease vaccination. Vaccine 23, 2327-30.
- Sorensen, H.P. and Mortensen, K.K. (2005) Advanced genetic strategies for recombinant protein expression in *Escherichia coli*. J Biotechnol 115, 113-28.
- Staffler, G., Szekeres, A., Schutz, G.J., Saemann, M.D., Prager, E., Zeyda, M., Drbal, K., Zlabinger, G.J., Stulnig, T.M. and Stockinger, H. (2003) Selective inhibition of T cell activation via CD147 through novel modulation of lipid rafts. J Immunol 171, 1707-14.
- Stockinger H., Ebel T., Hansmann C., Koch C., Majdic O., Prager E., Patel D. D., Fox D. A., Horejsi V., Sagawa K., and Shen D. C. CD147 (neurothelin/basigin) workshop panel report. In Kishimoto T., Kikutani H., von dem Borne A. E. G.K., Goyert S. M., Mason D. Y., Miyasaka M., Moretta L., Okumura K., Shaw S., Springer T. A., Sugamura K. and Zola, H.,eds, Leukocyte Typing VI. Garland Publishing, New York. 1997: 760.
- Stoltz, J., Ludwig, A. and Sauer, N. (1998) Bacteriophage lambda surface display of a bacterial biotin acceptor domain reveals the minimal peptide size required for biotinylation. FEBS Lett 440, 213-7.
- Tayapiwatana, C., Arooncharus, P. and Kasinrerk, W. (2003) Displaying and epitope mapping of CD147 on VCSM13 phages: influence of *Escherichia coli* strains. J Immunol Methods 281, 177-85.
- Terpe, K. (2003) Overview of tag protein fusions: from molecular and biochemical fundamentals to commercial systems. Appl Microbiol Biotechnol 60, 523-33.

- Thorns, C., Feller, A.C. and Merz, H. (2002) EMMPRIN (CD 174) is expressed in Hodgkin's lymphoma and anaplastic large cell lymphoma. An immunohistochemical study of 60 cases. *Anticancer Res* 22, 1983-6.
- Tsao, K.L., DeBarbieri, B., Michel, H. and Waugh, D.S. (1996) A versatile plasmid expression vector for the production of biotinylated proteins by site-specific, enzymatic modification in *Escherichia coli*. *Gene* 169, 59-64.
- Tucker, J. and Grisshammer, R. (1996) Purification of a rat neurotensin receptor expressed in *Escherichia coli*. *Biochem J* 317 (Pt 3), 891-9.
- Vanhove, M., Guillaume, G., Ledent, P., Richards, J.H., Pain, R.H. and Frere, J.M. (1997) Kinetic and thermodynamic consequences of the removal of the Cys-77-Cys-123 disulphide bond for the folding of TEM-1 beta-lactamase. *Biochem J* 321 (Pt 2), 413-7.
- Vasina, J.A. and Baneyx, F. (1997) Expression of aggregation-prone recombinant proteins at low temperatures: a comparative study of the *Escherichia coli cspA* and *tac* promoter systems. *Protein Expr Purif* 9, 211-8.
- Venturi, M., Seifert, C. and Hunte, C. (2002) High level production of functional antibody Fab fragments in an oxidizing bacterial cytoplasm. *J Mol Biol* 315, 1-8.
- Verma, J.N., Rao, M., Amselem, S., Krzych, U., Alving, C.R., Green, S.J. and Wassef, N.M. (1992) Adjuvant effects of liposomes containing lipid A: enhancement of liposomal antigen presentation and recruitment of macrophages. *Infect Immun* 60, 2438-44.
- Villacres-Eriksson, M., Behboudi, S., Morgan, A.J., Trinchieri, G. and Morein, B. (1997) Immunomodulation by *Quillaja saponaria* adjuvant formulations: in

- vivo stimulation of interleukin 12 and its effects on the antibody response. *Cytokine* 9, 73-82.
- Villacres, M.C., Behboudi, S., Nikkila, T., Lovgren-Bengtsson, K. and Morein, B. (1998) Internalization of iscom-borne antigens and presentation under MHC class I or class II restriction. *Cell Immunol* 185, 30-8.
- Villaverde, A. and Carrio, M.M. (2003) Protein aggregation in recombinant bacteria: biological role of inclusion bodies. *Biotechnol Lett* 25, 1385-95.
- Watson, D.L., Watson, N.A., Fossum, C., Lovgren, K. and Morein, B. (1992) Interactions between immune-stimulating complexes (ISCOMs) and peritoneal mononuclear leucocytes. *Microbiol Immunol* 36, 199-203.
- Watson, W.H., Yang, X., Choi, Y.E., Jones, D.P. and Kehrer, J.P. (2004) Thioredoxin and its role in toxicology. *Toxicol Sci* 78, 3-14.
- Waugh, D.S. (2005) Making the most of affinity tags. *Trends Biotechnol* 23, 316-20.
- Weeratna, R.D., McCluskie, M.J., Xu, Y. and Davis, H.L. (2000) CpG DNA induces stronger immune responses with less toxicity than other adjuvants. *Vaccine* 18, 1755-62.
- Weiss, E., Chatellier, J. and Orfanoudakis, G. (1994) In vivo biotinylated recombinant antibodies: construction, characterization, and application of a bifunctional Fab-BCCP fusion protein produced in *Escherichia coli*. *Protein Expr Purif* 5, 509-17.
- Wilchek, M. and Bayer, E.A. (1990) Introduction to avidin-biotin technology. *Methods Enzymol* 184, 5-13.
- Xiong, S., Wang, Y.F., Ren, X.R., Li, B., Zhang, M.Y., Luo, Y., Zhang, L., Xie, Q.L. and Su, K.Y. (2005) Solubility of disulfide-bonded proteins in the cytoplasm

of *Escherichia coli* and its "oxidizing" mutant. World J Gastroenterol 11, 1077-82.

Yao, X., Wei, D., Soden, C., Jr., Summers, M.F. and Beckett, D. (1997) Structure of the carboxy-terminal fragment of the apo-biotin carboxyl carrier subunit of *Escherichia coli* acetyl-CoA carboxylase. Biochemistry 36, 15089-100.



ลิขสิทธิ์ของมหาวิทยาลัยเชียงใหม่
Copyright © by Chiang Mai University
All rights reserved