

REFERENCES

1. Clark S.L., Hankins G.D. Temporal and demographical trends in cerebral palsy-fact and friction. *Am. J. Obstet. Gynecol* 2003;188:628-633.
2. Yang Y.R., Yen J.G., Wang R.Y., Yen L.L., Lieu F.K. Gait outcome after additional backward walking training in patients with stroke: a randomized controlled trial. *Clin Rehabil* 2005;19(3):264-273.
3. Kuban K.C.K., Leviton A. Cerebral palsy. *N Engl Med* 1994;330:188-189.
4. Morgan D.W., Tseh W, Caputo T.L., Keefer D.J., Craig I.S., Griffith K.B. Longitudinal profile of O₂ uptake during treadmill walking in able-bodied children: the locomotion energy and growth study. *Gait & Posture* 2002;15:230-235.
5. Dodd K.J., Foley S. Partial body-weight-supported treadmill training can improve walking in children with cerebral palsy: a clinical controlled trial. *Dev Med Child Neurol* 2007;47(2):101-105.
6. Schndl M.R., Forstner C., Kern H. Treadmill training with partial body weight support in nonambulatory patients with cerebral palsy. *Arch Phys Med Rehabil* 2000;81:301-306.
7. Macko R.F., Treadmill aerobic exercise training reduces the energy expenditure and cardiovascular demands of hemiparetic gait chronic stroke patients. *Stroke* 1997;28:326-330.
8. In; <http://www.walking.about.com/treadmillworkouts/a/treadmillback.htm>. Walking and Jogging Backwards on the Treadmill; 2008.

9. Masumoto K., Takasugi S., Hotta N., Fujishima K., Iwamoto Y. A comparison of muscle activity and heart rate response during backward and forward walking on an underwater treadmill. *Gait & Posture* 2007;25:222-228.
10. Weng C.S., Wang J., Wang G., Gao L.P. Effectiveness of backward walking treadmill training in lower extremity function after stroke. *Zhonghua Yi Xue Za Zhi* 2006;86(37):2635-2638.
11. Lin J.P. The cerebral palsies: a physiological approach. *J Neurol Neurosurg Psychiatry* 2003;74:23-29.
12. van der Krogt M., M., Doorenbosch C.A.M., Harlaar J. Muscle length and lengthening velocity in voluntary crouch gait. *Gait & Posture* 2006;xx:xxx-xxx.
13. Thomas S.S., Moore C., Kelp-Lenane C., Norris C. Simulated gait pattern: the resulting effect on gait parameter, dynamic electromyography, joint moments, and physiological cost index. *Gait & Posture* 1996;4:100-7.
14. Hoon A.C.K., Johnston M.V. *Diseases of the Nervous System: Clinical Neuroscience and Therapeutic Principles*. 3 ed: Cambridge university; 2002.
15. In; <http://www.eengoeddoel.nl>. Cerebral palsy; 2005.
16. Janet M., Howle W. *Decision Making in Pediatric Neurologic Physical Therapy*. Philadelphia; 1999.
17. Duffy C.M., Hill A.E., Corry L.S., Graham H.K. Energy consumption in children with spina bifida and cerebral palsy: a comparative study. *Dev Med Child Neurol* 1996;38:238-243.
18. Pratanaphon S. *Physical Therapy in children with neurology*. Department of Physical Therapy, Chiang Mai University 1999.


19. Gage J.R. Gait analysis; an essential tool in the treatment of cerebral palsy. Clin Orthop Rel Res 1993;288:126-134.
20. Kramer J.F., Macphail H.E. Relationships among measures of walking efficiency, grossmotor ability and isokinetic strenght in adolescents with cerebral palsy. Pediatr Phys Ther 1994;6:3-8.
21. Mann R. Biomechanics in cerebral palsy. Foot Ankle 1983;4:114-119.
22. Olney S.J., Costigan P.A., Hedden D.M. Mechanical energy patterns in gait of cerebral palsied children with hemiplegia. Phys Ther 1987;67:1348-1354.
23. Waters R.L. Energy expenditure. Gait Analysis: Normal and Pathological Function 1992:443-489.
24. Winter D.A. Knee flexion during stance as a determinant of inefficient walking. Phys Ther 1983;63:331-333.
25. Damiano D.L., Able M.F. Functional outcomes of strength training in spastic cerebral palsy. Arch Phys Med Rehabil 1998;79:119-125.
26. Mossberg K.A., Lintan K.A., Friske K. Ankle-foot orthoses: Effect on energy expenditure of gait in spastic diplegia children. Arch Phys Med rehabill 1990;37:490-4.
27. In; <http://www.cornerstoneceu.com>; 2007
28. Seif-naraghi A.H, Herman R.M. A novel method for locomotion training. J Head Trauma Rehabil 1999;14(2):146-162.
29. Barbeau H. Does neurorehabilitation play a role in the recovery of walking in neurological populations? Annal New York Academic of Sciences 1998;860:377-92.

30. Sidaway B., Anderson J., Danielson G., Martin L., Smith G. Effect of Long-Term Gait Training Using Visual Cues in an Individual with Parkinson disease. *Phys Ther* 2006;86:186-94.
31. Hess S., Bertelt C., Jahnke T.M., Schaffrin A., Baake P., Malezic M. Treadmill Training With Partial Body Weight Support Compared With Physiotherapy in Nonambulatory Hemiparetic Patients *Stroke* 1995;26:976-81.
32. Maltais D., Bar-Or O., Pierrynowski M., Galea V. Repeated treadmill walks affect physiologic responses in children with cerebral palsy *Med Sci Sports Exerc* 2003.
33. Waagfjord J., Levangie K.P., Certo M.C. Effects of treadmill training on gait in a hemiparetic patient. *Phys Ther* 1990;70(9): 558-60.
34. Yoshimoto D., Mohr T., Palmgren A., Frappier J. An EMG study of quiceps and hamstrings activity during forward and backward walking. *Health Affairs & Medical Research* 1995:67-70.
35. Grasso R., Bianchi L., Lacquaniti F. Motor patterns for human gait: backward walking versus forward locomotion. *J. Neurophysiol.* 1998;80:1868-85.
36. Peebles K.C., Woodman A.D., Skinner M.A. The Physiological Cost Index In Elderly Subjects During Treadmill And Floor Walking. *NZ Journal of Physiotherapy* 2003;31:11-6.
37. Russell D.J., Rosenbaum P.L., Avery L.M., Lane M. Gross Motor Function Meseure (GMFM-66 & GMFM-88) User's Manual. Canada: Cambridge univercity press; 2002.

38. Russell D.J., Avery L.M., Rosenbaum P., Palisano R.J. Improve scaling of the Gross Motor Function Measure for children with cerebral palsy: Evidence of reliability and validity. *Physical Therapy* 2000;80:873-885.
39. วุฒิชัย เพิ่มศิริวานิชย์. การออกกำลังกายแบบแอโรบิค; การออกกำลังกายเพื่อสุขภาพและในโรคต่างๆ. กรุงเทพมหานคร: ชานเมืองการพิมพ์; 2004: 81-124.
40. ชูศักดิ์ เวชแพทย์. ผลการฝึกต่อระบบไหลเวียนเลือด; สรีรวิทยาของการออกกำลังกาย. กรุงเทพมหานคร: เทพรัตน์การพิมพ์; 1985:33-35, 137-158.
41. สลิลดา โรจนธรรมณี. สรีรวิทยาการออกกำลังกาย. สมุทรปราการ: ศูนย์เทคโนโลยีการศึกษา มหาวิทยาลัยหัวเฉียวเฉลิมพระเกียรติ; 200:61-72.
42. Raja K., Joseph B., Benjamin S., Minocha V., Rana B. Physiological cost index in cerebral palsy. *J Pediatr Orthop* 2007;27:130-136.
43. MacGregor J. The evaluation of patient performance using long term ambulatory monitoring technique in the domiciliary environment. *Phys Ther* 1981;67(2):30-33.
44. Pohl M., Mehrholz J., Ritschel C., Ruckriem S. Speed-dependent treadmill training in ambulatory hemiparetic stroke patients: a randomized controlled trial. *Stroke* 2002;33:553-558.
45. Flynn T.W., Soutas-Littel R.W. Mechanical power and muscle action during forward and backward running. *J Orthop Sports Phys Ther* 1993;17.

46. Myatt G., Baxter R., Dougherty R., Williams G., Halle J., Stetts D. The cardiopulmonary cost of backward walking at selected speed. *J Orthop Sports Phys Ther* 1995;21:132-138.
47. Vilensky J.A., Gankiewicz E., Gehlsen G. A kinematic comparison of backward and forward walking in human. *J Hum Mov Studies* 1987;13:29-50.
48. Flynn T.W., Connery S.M., Smutok M.A., Zeballos R.J., Weisman I.M. Comparison of cardiopulmonary responses to forward and backward walking and running. *Med Sci Sport Exerc* 1994;26:89-94.
49. Cavanagh P.R., Williams K.R. The effect of stride length variation on oxygen uptake during distance running. *Med Sci Sport Exerc* 1982;14:30-5.
50. Flynn T.W., Soutas-Littel R.W. Patellofemoral joint compressive force in forward and backward running. *J Orthop Sports Phys Ther* 1995;21:277-282.
51. Cherng R.J., Liu C.F., Hong R.B. Effect of treadmill training with body weight support on gait and gross motor function in children with spastic cerebral palsy. *Am J Phys Med Rehabil* 2007;86:548-555.
52. Steinwender G., Saraph V., Zwick E.B., Steinwender W.L. Hip locomotion mechanisms in cerebral palsy crouch gait. *Gait and Posture* 2001;13:78-75.
53. Murray A.W., Robb J.E. The hip in cerebral palsy. *Current Orthopaedics* 2006;20:286-293.
54. Cipriani D.J., Armstrong C.W., Gaul S.G. Backward walking at three levels of treadmill inclination: An electromyographic and kinematic analysis. *JOSPT* 1995;22:95-102.

55. Blackburn M., van Vliet P., Mockett S.P. Reliability of measurements obtained with the Modified Ashworth Scale in the lower extremities of people with stroke. *Phys Ther* 2002;82:25-34.
56. In; <http://www.neurosurgical.ca/ClinicalAssistant/scales/ashworth.html>.
57. McDowell B.C., Kerr C., Parkes J. Validity of a 1 minute walk test for children with Cerebral palsy. *Dev Med Chil Neurol* 2005;47:744-748.



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved