

## เอกสารอ้างอิง

- [1.] Gossuin G, Burtea C, Monseux A, et al. Ferritin-induced relaxation in tissues: an in vitro study. *J Magn Reson Imaging* 2004; 20: 690-696.
- [2.] Brooks RA, Vyzamal J, Goldfarb RB, et al. Relaxometry and magnetometry of ferritin. *Magn Reson Med* 1998; 40: 227-235.
- [3.] Zurlo MG, D.S.P., Borgna-Pignatti C, et al. Survival and causes of death in thalassaemia major. *Lancet*, 1989; 2: 27–30.
- [4.] Olivieri NF, N.D., MacMillan JH, et al. Survival in medically treated patients with homozygous beta-thalassaemia. *N Engl J Med* 1994; 331: 574–578.
- [5.] Anderson LJ, Holden S, Davis B, et al. Cardiovascular T2-star (T2\*) magnetic resonance for the early diagnosis of myocardial iron overload. *Eur Heart J* 2001; 22: 2171-2179.
- [6.] Positano V, Pepe A, Santarelli MF, et al. Multislice multiecho T2\* cardiac magnetic resonance for the detection of heterogeneous myocardial iron distribution in thalassaemia patients. *J Magn Reson Imaging* 2006; 23: 662–668
- [7.] Westwood MA, Wonke B, Maceira AM, et al. Left ventricular diastolic function compared with T2\* cardiovascular magnetic resonance for early detection of myocardial iron overload in thalassaemia major. *J Magn Reson Imaging* 2005; 22: 229-33.
- [8.] ไพรัช สายวิรุณพร, MRI: หลักการและความปลอดภัย พิมพ์ครั้งที่ 1 กรุงเทพฯ : วัฒนาพรินติ้ง; 2551.

- [9.] Harmening DH, et al. Clinical Hematology and Fundamental of Hemostasis, 3rd ed, Philadelphia: F.A. Davis Company, 1997.
- [10.] McKenzie SB, et al., Text Book of Hematology, 2nd ed, Baltimore: Williams & Wilkins, A Waverly Company. 1995.
- [11.] Winichagoon P, Fucharoen S, Wasi P, et al. Severity differences in  $\beta$ -thalassemia/haemoglobin E syndromes: Implication of genetic factors. Br J Haematol 1993; 633-639.
- [12.] Wasi P, Pootrakul P, Fucharoen S, et al. Thalassemia in southeast Asia: determination of different degrees of severity of anemia in thalassemia. Ann N Y Acad Sci 1985; 445: 119-126.
- [13.] Wood JC, Agus A. Onset of cardiac iron loading in pediatric patients with thalassemia major. Haematologica 2008; 93: 917-920.
- [14.] Gabutti V, P.A. Results of long-term iron-chelating therapy. Acta Haematol 1996; 95: 26-36.
- [15.] Conrad ME, Umbreit JN. Pathways of iron absorption. Blood Cells Mol Dis 2002; 29: 336-355.
- [16.] Kumfu S, Chattipakorn S, Fucharoen S, et al. Pathways of iron uptake into cardiomyocytes. Int J Cardiol 2009.
- [17.] Pakbaz Z, Fischer R, Fung E, et al. Serum ferritin underestimates liver iron concentration in transfusion independent thalassemia patients as compared to regularly transfused thalassemia and sickle cell patients. Pediatr Blood Cancer 2007; 49: 329-332.
- [18.] Mavrogeni SI, Markussis V, Kaklamanis L, et al. A comparison of magnetic resonance imaging and cardiac biopsy in the evaluation of heart iron overload

in patients with beta-thalassemia major. *European Journal of Haematology* 2005; 75: 241-247.

- [19.] Ghugre NR, Enriquez CM, Gonzalez I, et al. MRI detects myocardial iron in the human heart. *Magn Reson Med* 2006; 56: 681-686.
- [20.] Liu P, O.N. Iron overload cardiomyopathies: new insights into an old disease. *Cardiovasc Drugs Ther* 1994; 8: 101-110.
- [21.] Olson LJ, E.W., Mccall JT, et al. Cardiac iron deposition in idiopathic hemochromatosis: histologic and analytic assessment of 14 hearts from autopsy. *J Am Coll Cardiol* 1987; 10: 1239-1243.
- [22.] Leonardi B, Margossian R, Colan SD, et al. Relationship of magnetic resonance imaging estimation of myocardial iron to left ventricular systolic and diastolic function in thalassemia. *JACC Cardiovasc Imaging* 2008; 1: 572-578.
- [23.] Vincenzo P, Alessia P, Maria Filomena S, et al. Multislice multiecho T2\* cardiac magnetic resonance for the detection of heterogeneous myocardial iron distribution in thalassaemia patients. *NMR in Biomedicine* 2009. 22: 707-715.
- [24.] Westwood M, Anderson LJ, Firmin DN, et al. A single breath-hold multiecho T2\* cardiovascular magnetic resonance technique for diagnosis of myocardial iron overload. *J Magn Reson Imaging* 2003; 18: 33-39.
- [25.] Bonetti MG, Castriota-Scanderbeg A, Criconia GM, et al. Hepatic iron overload in thalassemic patients: proposal and validation of an MRI method of assessment. *Pediatr Radiol* 1996. 26: 650-656.
- [26.] Ernst O, Sergent G, Bonvarlet P, et al. Hepatic iron overload: diagnosis and quantification with MR imaging. *Am J Roentgenol* 1997; 168: 1205-1208.

- [27.] Argyropoulou M, Kiortsis D, Astrakas L, et al. Liver, bone marrow, pancreas and pituitary gland iron overload in young and adult thalassaemic patients: a T2 relaxometry study. *European Radiology* 2007; 17: 3025-3030.
- [28.] Papanikolaou N, Ghiatas A, Kattamis A, et al. Non-invasive myocardial iron assessment in thalassaemic patients. T2 relaxometry and magnetization transfer ratio measurements. *Acta Radiol* 2000; 41: 348-351.
- [29.] Mavrogeni SI, Gotsis ED, Markussis V, et al, T2 relaxation time study of iron overload in b-thalassemia. *Magma* 1998; 6: 7-12.
- [30.] Pennell D.J. T2\* magnetic resonance and myocardial iron in thalassemia. *Ann N Y Acad Sci* 2005; 1054: 373-378.
- [31.] Positano V, Pepe A, Santarelli MF, et al. Standardized T2\* map of normal human heart in vivo to correct T2\* segmental artefacts. *NMR Biomed* 2007; 20: 578-590.
- [32.] Ramazzotti A, Pepe A, Positano V, et al. Standardized T2\* map of a normal human heart to correct T2\* segmental artefacts; myocardial iron overload and fibrosis in thalassemia intermedia versus thalassemia major patients and electrocardiogram changes in thalassemia major patients. *Hemoglobin* 2008; 32: 97-107.
- [33.] Pepe A, Positano V, Santarelli MF, et al. Multislice multiecho T2\* cardiovascular magnetic resonance for detection of the heterogeneous distribution of myocardial iron overload. *J Magn Reson Imaging* 2006; 23: 662-668.

- [34.] Ramazzotti A, Pepe A, Positano V, et al. Multicenter validation of the magnetic resonance T2\* technique for segmental and global quantification of myocardial iron. *J Magn Reson Imaging* 2009; 30: 62-68.
- [35.] Tanner MA, Galanello R, Dessi C, et al. Myocardial iron loading in patients with thalassemia major on deferoxamine chelation. *J Cardiovasc Magn Reson* 2006; 8: 543-547.
- [36.] He T, Kirk P, Firmin DN, et al. Multi-center transferability of a breath-hold T2 technique for myocardial iron assessment. *J Cardiovasc Magn Reson* 2008; 10: 11.
- [37.] Tanner MA, He T, Westwood MA, et al. Multi-center validation of the transferability of the magnetic resonance T2\* technique for the quantification of tissue iron. *Haematologica* 2006; 91: 1388-1391.
- [38.] He T, Gatehouse PD, Kirk P, et al. Black-blood T2\* technique for myocardial iron measurement in thalassemia. *J Magn Reson Imaging* 2007; 25: 1205-1209.
- [39.] He T, Gatehouse PD, Smith GC, et al. Myocardial T2\* measurements in iron-overloaded thalassemia: An in vivo study to investigate optimal methods of quantification. *Magn Reson Med* 2008; 60: 1082-1089.
- [40.] He T, Smith PD, Mohiaddin GC, et al. Myocardial T2\* measurements in iron-overloaded thalassemia: An ex vivo study to investigate optimal methods of quantification. *Magn Reson Med* 2008; 60: 350-356.
- [41.] Nilesh RG, Cathleen ME, Thomas DC, et al. Improved R2\* measurements in myocardial iron overload. *J Magn Reson Imaging* 2006; 23: 9-16.
- [42.] Firmin D, Keegan J, Navigator echoes in cardiac magnetic resonance. *J Cardiovasc Magn Reson* 2001; 3: 183-193.

- [43.] Jung B, Zaitsev M, Hennig J, et al. Navigator gated high temporal resolution tissue phase mapping of myocardial motion. *Magn Reson Med* 2006; 55: 937-942.
- [44.] Spuentrup E, Bornert P, Botnar RM, et al. Navigator-gated free-breathing three-dimensional balanced fast field echo (TrueFISP) coronary magnetic resonance angiography. *Invest Radiol* 2002; 37: 637-642.
- [45.] Sachs TS, Hu BS, Kohli J, et al. Real-time motion detection in spiral MRI using navigators. *Magn Reson Med* 1994; 32: 639-645.
- [46.] Harrison PM, Arosio P. The ferritins: molecular properties, iron storage function and cellular regulation. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* 1996; 1275: 161-203.
- [47.] Wacker CM, Bock M, Hartlep AW, et al. Changes in myocardial oxygenation and perfusion under pharmacological stress with dipyridamole: assessment using T2\* and T1 measurements. *Magn Reson Med* 1999; 41: 686-695.
- [48.] Reeder SB, Faranesh AZ, Boxerman JL, et al. In vivo measurement of T2\* and field inhomogeneity maps in the human heart at 1.5 T. *Magn Reson Med*, 1998; 39: 988-998.