

REFERENCES

1. Chan KM, Pang WS, Ee CH, Ding YY, Choo P. Epidemiology of falls among the elderly community dwellers in Singapore. *Singapore Med J* 1997; 38: 427-31.
2. Nacreiner NM, Findorff MJ, Wyman JF, McCarthy TC. Circumstances and consequences of falls in community-dwelling older women. *J Womens Health* 2007; 16: 1437-46.
3. Centers for Disease Control and Prevention National Center for Injury Prevention and Control. Falls among older adults: An overview. *Journal* [serial on the Internet]. 2005 Date: Available from: <http://www.cdc.gov/ncipc>
4. Dunne RG, Bergman AB, Rogers LW, Inglin B, Rivara FP. Elderly persons' attitudes towards footwear factor in preventing falls. *Public Health Rep* 1993; 108: 245-8.
5. Fuller GF. Falls in the Elderly. *Am Fam Physician* 2000.
6. Chaiwanichsiri D, Tantisiriwat N, Janchai S. Proper shoe sizes for Thai elderly. *The Foot* 2008; 18: 186-91.
7. Jitapunkul S, Songkhla M, Chayovan N, Chirawatkul A, Choprapawon C, Kachondham Y. Falls and their associated factors: A national survey of the Thai elderly. *J Med Assoc Thai* 1998; 81: 233-42.

8. Resnick B, Junlapeeya P. Falls in a community of older adults: Findings and implications for practice. *Appl Nurs Res* 2004; 17: 81-91.
9. Sherrington C, Benz HB. An evaluation of footwear worn at the time of fall-related hip fracture. *Age Ageing* 2003; 32: 310-4.
10. Fletcher PC, Hirdes JP. Restriction in activity associated with fear of falling among community-based seniors using home care services. *Age Ageing* 2004; 33: 273-9.
11. Zijlstra GAR, Haastregt JCMV, Eijk JTMV, Rossum EV, Stalenhoef PA, Kempen GIJM. Prevalence and correlates of fear of falling, and associated avoidance of activity in the general population of community-living older people. *Age Ageing* 2007; 36: 304-9.
12. Shanthi G, Krishnaswamy B. Risk factors for falls in elderly. *J Indian Acad Geriatrics* 2005; 1: 57-60.
13. Koepsell TD, Wolf ME, Buchner DM, Kukull WA, LaCroix AZ, Tencer AF, et al. Footwear style and risk of falls in older adults. *JAGS* 2004; 52: 1495-501.
14. Menz HB, Lord SR. Footwear and postural stability in older people. *J Am Podiatr Med Assoc* 1999; 89: 346-57.
15. Dyer C, Watkins C, Gould C, Rowe J. Risk-factor assessment for falls: From a written checklist to the penless clinic. *Age Ageing* 1998; 27: 569-72.
16. Barbieri E. Patient falls are not patient accidents. *J Gerontol Nurs* 1983; 9: 165.

17. Gabell A, Simons M, Nayak U. Falls in the elderly: Predisposing causes. *Ergonomics* 1985; 28: 965–75.
18. Menant JC, Perry SD, Steele JR, Menz HB, Munro BJ, Lord SR. Effects of shoe characteristics on dynamic stability when walking on even and uneven surfaces in young and older people. *Arch Phys Med Rehabil* 2008; 89: 1970-6.
19. Menant JC, Steele JR, Menz HB, Munro BJ, Lord SR. Effects of walking surfaces and footwear on temporo-spatial gait parameters in young and older people. *Gait Posture* 2008; 29: 392-7.
20. Hale WA, Delaney MJ, McGaghie WC. Characteristics and predictors of falls in elderly patients. *J Fam Pract* 1992; 34: 577-81.
21. Shumway-Cook A, Woollacott MH. *Motor control: Theory and practical applications*. 2nd ed.: Lippincott Williams&Wilking; 2001.
22. World Health Organization. Definition of an older or elderly person. *Journal [serial on the Internet]*. 2009 Date: Available from: <http://www.who.int/healthinfo/survey/ageingdefnolder/en/>.
23. Orimo H, Ito H, Suzuki T, Araki A, Hosoi T, Sawabe M. Reviewing the definition of “elderly”. *Geriatr Gerontol Int* 2006; 6: 149–58.
24. ราชกิจจานุเบกษา. พระราชบัญญัติผู้สูงอายุ พ.ศ. 2546. In: กระทรวงการพัฒนาศังคมและความมั่นคงของมนุษย์, editor.; 2546.
25. Nevitt MC, Cummings SR, Kidd S, Black D. Risk factors for recurrent nonsyncopal falls. A prospective study. *JAMA* 1989; 261: 2663-8.

26. Zecevic AA, Salmoni AW, Speechley M, Vandervoort AA. Defining a fall and reasons for falling: Comparisons among the views of seniors, health care providers, and the research literature. *Gerontologist* 2006; 46 367–76.
27. Rubenstein L, Josephson K. The epidemiology of falls and syncope. *Clin Geriatr Med* 2002; 18: 141-58.
28. Alexander BH, Rivara FP, Wolf ME. The cost and frequency of hospitalization for fall-related injuries in older adults. *Am J Public Health* 1992; 82: 1020-3.
29. Li W, Keegan THM, Sternfeld B, Sidney S, Quesenberry CP, Kelsey JL. Outdoor falls among middle-aged and older adults: a neglected public health problem. *Am J Public Health* 2006; 96: 1192-200.
30. Assantachai P, Maranetra N. Factors determining hospital admission of Thai elderly by a mailed survey. *J Med Assoc Thai* 2005; 88(): 1051-6.
31. มูลนิธิสถาบันวิจัยและพัฒนาผู้สูงอายุไทย(มส.ผส.). รายงานสถานการณ์ผู้สูงอายุไทย พ.ศ. 2550. กรุงเทพมหานคร: บริษัท ทีคิวพี จำกัด; 2551.
32. Bergland A, Wyller TB. Risk factors for serious fall related injury in elderly women living at home. *Inj Prev* 2004; 10: 308-13.
33. Cummings SR, Nevitt MC, Browner WS, Stone K, Fox KM, Ensrud KE, et al. Risk factors for hip fracture in white women. *N Engl J Med* 1995; 332 767-73.

34. Yardley L, Smith H. A prospective study of the relationship between feared consequences of falling and avoidance of activity in community-living older people. *Gerontologist* 2002; 42: 17-23.
35. Chamonchant D, Gaogasigam C, Boonyong S. A study in fall and balance in Thai elderly population. Bangkok: Chulalongkorn University; 2005.
36. Neyens J, Dijcks B, van Haastregt J, de Witte L, van den Heuvel W, Crebolder H, et al. The development of a multidisciplinary fall risk evaluation tool for demented nursing home patients in the Netherlands. *BMC Public Health* 2006; 6(1): 74.
37. Kenny RA, Rubenstein LZ, Martin FC, Tinetti ME. Guideline for the prevention of falls in older persons. *J Am Geriatr Soc* 2001; 49: 664–72.
38. Menz HB, Morris ME, Lord SR. Footwear characteristics and risk of indoor and outdoor falls in older people. *Gerontology* 2006; 52: 174-80.
39. Hartikainen S, Loönnroos E, Louhivuori K. Medication as a risk factor for falls: Critical systematic review. *J Gerontol* 2007; 62A: 1172–81.
40. Bueno-Cavanillas A, Padilla-Ruiz F, Jiménez-Moleón J, Peinado-Alonso C, Gálvez-Vargas R. Risk factors in falls among the elderly according to extrinsic and intrinsic precipitating causes. *Eur J Epidemiol* 2000; 16: 849-59.
41. Horak F, Macpherson J. Postural orientation and equilibrium. In: Shepard J, Rowell L, editors. *Handbook of physiology, section 12 Exercise: regulation and integration of multiple systems*. New York: Oxford University; 1996. p. 255-92.

42. Nashner L. Evaluation of postural stability, movement, and control. In: Hasson S, editor. *Clinical exercise physiology*. St. Louis: Mosby; 1994. p. 199-234.
43. Akram SB, Frank JS, Patla AE, Allum JHJ. Balance control during continuous rotational perturbations of the support surface. *Gait Posture* 2008; 27: 393–8.
44. Horak FB, Wrisley DM, Frank j. The Balance Evaluation Systems Test (BESTest) to differentiate balance deficits. *Phys Ther* 2009; 89: 484-98.
45. O'Connell M, George K, Stock D. Postural sway and balance testing: A comparison of normal and anterior cruciate ligament deficient knees. *Gait Posture* 1998; 8 136–42.
46. Laessoe U, Voigt M. Anticipatory postural control strategies related to predictive perturbations. *Gait Posture* 2008; 28: 62-8.
47. Menz HB, Sherrington C. The Footwear Assessment Form: A reliable clinical tool to assess footwear characteristics of relevance to postural stability in older adults. *Clin Rehabil* 2000; 14: 657–64.
48. Arnadottir SA, Mercer VS. Effects of footwear on measurements of balance and gait in women between the ages of 65 and 93 years. *Phys Ther* 2000; 80: 17-27.
49. Tencer AF, Koepsell TD, Wolf ME, Frankenfeld CL, Buchner DM, Kukull WA, et al. Biomechanical properties of shoes and risk of falls in older adults. *J Am Geriatr Soc* 2004; 52: 1840-6.
50. Menant JC, Steele JR, Menz HB, Munro BJ, Lord SR. Effects of footwear features on balance and stepping in older people. *Gerontology* 2008; 54: 18–23.

51. Lord S, Bashford G. Shoe characteristics and balance in older women. *J Am Geriatr Soc* 1996; 44: 429–33.
52. Menant JC, Steele JR, Menz HB, Munro BJ, Lord SR. Rapid gait termination: Effects of age, walking surfaces and footwear characteristics. *Gait Posture* 2009; 30: 65–70.
53. Perry SD, Radtke A, Goodwin CR. Influence of footwear midsole material hardness on dynamic balance control during unexpected gait termination. *Gait Posture* 2007; 25: 94–8.
54. Menz H, Lord S, McIntosh A. Slip resistance of casual footwear: Implications for falls in older adults. *Gerontology* 2001; 47: 145–9.
55. Lloyd D, Stevenson M. Measurement of slip resistance of shoes on floor surfaces: Part 2. effect of a beveled heel. *J Occup Health Saf* 1989; 5: 229.
56. Berg K, Wood-Dauphinee S, Williams JI, Gayton D. Measuring balance in the elderly: Preliminary development of an instrument. *Physiother Can* 1989; 41: 304-11.
57. Tinetti M. Performance-oriented assessment of mobility problems in elderly patients. *J Am Geriatr Soc* 1986; 34: 119-26.
58. Bohannon RW, Leary KM. Standing balance and function over the course of acute rehabilitation. *Arch Phys Med Rehabil* 1995; 76: 994–6.
59. Duncan PW, Weiner DK, Chandler J, Studenski S. Functional reach: A new clinical measure of balance. *J Gerontol* 1990; 45: M192-M7.
60. Brauer S, Burns Y, Galley P. Lateral reach: a clinical measure of medio-lateral postural stability. *Physiother Res Int* 1999; 4: 81-8.

61. Isles R, Choy NL, Steer M, Nitz J. Normal values of balance tests in women aged 20–80. *J Am Geriatr Soc* 2004; 52: 1367–72.
62. Shumway-Cook A, Horak FB. Assessing the influence of sensory integration on balance. Suggestions from the field. *Phys Ther* 1986; 66: 1548-9.
63. Whitney SL, Wrisley DM. The influence of footwear on timed balance scores of the modified clinical test of sensory interaction and balance. *Arch Phys Med Rehabil* 2004; 85: 439-43.
64. Wrisley DM, Whitney SL. The effect of foot position on the modified Clinical Test of Sensory Interaction and Balance. *Arch Phys Med Rehabil* 2004; 85: 335-8.
65. Steffen TM, Hacker TA, Mollinger L. Age- and gender-related test performance in community-dwelling elderly people: Six-Minute Walk Test, Berg Balance Scale, Timed Up & Go Test, and Gait Speeds. *Phys Ther* 2002; 82: 128–37.
66. Hedel HJAv, Dietz V, Armin Curt M. Assessment of walking speed and distance in subjects with an incomplete spinal cord injury. *Neurorehabil Neural Repair* 2007; 21: 295 - 301.
67. Oberg T, Karsznia A, Oberg K. Basic gait parameters : Reference data for normal subjects,10-79 years of age. *J Rehabil Res Dev* 1993; 30(2).
68. Podsiadlo D, Richardson S. The timed “Up & Go”: a test of basic functional mobility for frail elderly persons. *J Am Geriatr Soc* 1991; 39: 142–8.
69. Wall JC, Bell C, Campbell S, Davis J. The timed get-up-and-go test revisited : Measurement of the component tasks. *J Rehabil Res Dev* 2000; 37: 109-13.

70. Bohannon RW, Schaubert K. Long-term reliability of the Timed Up-and-Go test among community-dwelling elders. *J Phys Ther Sci* 2005; 17: 93-6.
71. Shumway-Cook A, Brauer S, Woollacott M. Predicting the probability for falls in community-dwelling older adults using the Timed Up & Go test. *Phys Ther* 2000; 80: 896 –903.
72. National Statistical Office of Thailand. Report on the 2007 survey of the older persons in Thailand. 2009 [cited. Available from: www.nso.go.th/].
73. Bohannon RW. Comfortable and maximum walking speed of adults aged 20-79 years: Reference values and determinants. *Age Ageing* 1997; 26: 15-9.
74. Murphy MA, Olson SL, Protas EJ, Overby AR. Screening for falls in community-dwelling elderly. *J Aging Phys Act* 2003; 11: 64-78.