

REFERENCES

1. Faulkner K, Redfern M, Cauley J, Landsittel D, Studenski S, Rosano C, et al. Multitasking: association between poorer performance and a history of recurrent falls. *J Am Geriatr Soc* 2007; 55: 570-6.
2. American Geriatrics Society BGS, and American Academy of Prevention OSPoF. Guideline for the prevention of falls in older Persons. *J Am Geriatr Soc* 2001; 49: 664-72.
3. Soriano TA, DeCherrie LV, Thomas DC. Falls in the community-dwelling older adult: a review for primary-care providers. *Clin Interv Aging* 2007; 4: 545–53.
4. Beauchet O, Dubost V, Allali G, Gonthier R, Hermann F, Kressig R. 'Faster counting while walking'as a predictor of falls in older adults. *Age Ageing* 2007; 36: 418-23.
5. Hollman J, Kovash F, Kubik J, Linbo R. Age-related differences in spatiotemporal markers of gait stability during dual task walking. *Gait Posture* 2006; 26: 113-9.
6. Lajoie Y, Gallagher S. Predicting falls within the elderly community: comparison of postural sway, reaction time, the berg balance scale and the activities-specific comparing fallers and non-fallers. *Arch Gerontol Geriatr* 2004; 38: 11-26.
7. Melzer I, Kurz I, Shahar D, Levi M, Oddsson L. Application of the voluntary step execution test to identify elderly fallers. *Age Ageing* 2007; 36: 532-7.
8. Shumway-Cook A, Baldwin M, Polissar N, Gruber W. Predicting the probability for falls in community-dwelling older adults. *Phys Ther* 1997; 77: 812-8.

9. Lundin-Olsson L, Nyberg L, Gustafson Y. "Stops walking when talking" as a predictor of falls in elderly people. *Lancet* 1997; 349: 617.
10. Shumway-Cook A, Woollacott M. Attentional demands and postural control: the effect of sensory context. *J Gerontol A Biol Sci Med Sci* 2000; 55: M10-6.
11. Shumway-Cook A, Woollacott M, Kerns KA, Baldwin M. The effects of two types of cognitive tasks on postural stability in older adults with and without a history of falls. *J Gerontol Med Sci* 1997; 52A: M232-40.
12. Verghese J, Buschke H, Viola L, Katz M, Hall C, Kuslansky G, et al. Validity of divided attention tasks in predicting falls in older individuals: a preliminary study. *J Am Geriatr Soc* 2002; 50: 1572-6.
13. Beauchet O, Allali G, Annweiler C, Berrut G, Maarouf N, Herrmann FR, et al. Does change in gait while counting backward predict the occurrence of a first fall in older adults? *Gerontology* 2008; 54: 217-23.
14. Beauchet O, Annweiler C, Allali G, Berrut G, Herrmann F, Dubost V. Recurrent falls and dual task-related decrease in walking speed: Is there a relationship? *J Am Geriatr Soc* 2008; 56: 1265-9.
15. Springer S, Giladi N, Peretz C, Yogev G, Simon ES, Hausdorff JM. Dual-tasking effects on gait variability: the role of aging, falls, and executive Function. *Mov Disord* 2006; 21: 950-7.
16. Hausdorff JM, Edelberg HK, Mitchell SL, Goldberger AL, Wei JY. Increased gait unsteadiness in community-dwelling elderly fallers. *Arch Phys Med Rehabil* 1997; 78: 278-83.
17. Beauchet O, Dubost V, Herrmann F, Rabilloud M, Gonthier R, Kressig R. Relationship between dual-task related gait changes and intrinsic risk factors


- for falls among transitional frail older adults. *Aging Clin Exp Res* 2005; 17: 270-5.
18. Hausdorff JM, Schweiger A, Herman T, Yogeve-Seligmann G, Giladi N. Dual-task decrements in gait: contributing factor among healthy older adults. *J Gerontol A Biol Sci Med Sci* 2008; 63A: 1335-43.
 19. Siu K-C, Chou L-S, Mayr U, Donkelaar Pv, Woollacott MH. Attentional mechanisms contributing to balance constraints during gait: The effects of balance impairments. *Brain Res* 2009; 1248: 59-67.
 20. Shumway-Cook A, Woollacott MH. Motor control: translating research into Clinical practice. 3rd ed. Philadelphia: Lippincott Williams&Wilkins; 2007.
 21. Blake AJ, Morgan K, Bendall MJ, Dallosso H, Ebrahim SBJ, Arie THD, et al. Falls by elderly people at home: prevalence and associated factors. *Age Ageing* 1988; 17: 365-72.
 22. Alexander NB. Gait disorders in older adults. *J Am Geriatr Soc* 1996; 49: 434-51.
 23. Priest AW, Salamon KB, Hollman JH. Age-related differences in dual task walking: a cross sectional study. *J Neuroeng Rehabil* 2008; 29: 1-8.
 24. Horak FB. Clinical assessment of balance disorders. *Gait Posture* 1997; 6: 76-84.
 25. Mansfield A, Peters AL, Liu BA, Maki BE. A perturbation-based balance training program for older adults: study protocol for a randomized controlled trial. *BMC Geriatr* 2007; 7: 1-14.
 26. Overstall PW, Exton-Smith AN, Imms FJ, Johnson AL. Falls in the elderly related to postural imbalance. *Br Med J* 1977; 1: 261-4.

27. Tinetti ME, Baker DI, McAvay G, Claus EB, Garrett P, Gottschalk M, et al. A multifactorial intervention to reduce the risk of falling among elderly people living in the community. *N Engl J Med* 1994; 331: 821-7.
28. Trader SE, Newton RA, Cromwell RL. Balance abilities of homebound older adults classified as fallers and nonfallers. *J Geriatr Phys Ther* 2003; 26: 3-8.
29. Brach JS, Studenski S, Perera S, VanSwearingen JM, Newman AB. Stance time and step width variability have unique contributing impairments in older persons. *Gait Posture* 2008; 27: 431-9.
30. Coppin AK, Shumway-Cook A, Saczynski JS, Patel KV, Ble A, Ferrucci L, et al. Association of executive function and performance of dual-task physical tests among older adults: analyses from the InChianti study. *Age Ageing* 2006; 35: 619-24.
31. Bond JM and Morris M. Goal-directed secondary motor tasks: their effects on gait in subjects with Parkinson's disease. *Arch Phys Med Rehabil* 2000; 81: 110-6.
32. Yogev G, Giladi N, Springer S, Simon ES, Hausdorff JM. Dual tasking, gait rhythmicity, and Parkinson's disease: which aspects of gait are attention demanding? *Eur J Neurosci* 2005; 22: 1248-56.
33. Dubost V, Kressig RW, Gonthier R, Herrmann FR, Aminian K, Najafi B and Beauchet O. Relationships between dual-task related changes in stride velocity and stride time variability in healthy older adults. *Hum Mov Sci* 2006; 25: 372-82.
34. Herman T, Giladi N, Gurevich T, Hausdorff JM. Gait instability and fractal dynamics of older adults with a "cautious" gait: why do certain older adults walk fearfully? *Gait Posture* 2005; 21: 178-18.

35. Menz HB, Lord SR, Fitzpatrick RC. A structural equation model relating impaired sensorimotor function, fear of falling and gait patterns in older people. *Gait Posture* 2007; 25: 243–9.
36. Lemke MR, Wendorff T, Mieth B, Buhl K, Linnemann M. Spatiotemporal gait patterns during over ground locomotion in major depression compared with healthy controls. *J Psychiatr Res* 2000; 34: 277-83.
37. Hausdorff JM, Peng C-K, Goldberger AL, Stoll AL. Gait unsteadiness and fall risk in two affective disorders: a preliminary study. *BMC Psychiatry* 2004; 39: 1-7.
38. Fan K, McCandlis BD, Sommer T, Raz A, Posner MI. Testing the efficiency and independence of attentional networks. *J Cog Neurosci* 2002; 14: 340-7.
39. Yogev-Selimann G, Rotem-Galili Y, Mirelman A, Dickstein R, Giladi N, Hausdorff JM. How does explicit prioritization alter walking during dual-task performance? Effects of age and sex on gait speed and variability. *Phys Ther* 2010; 90: 177-86
40. Berg KO, Wood-Dauphine S, Williams JI, Gayton D. Measuring balance in the elderly: preliminary development of an instrument. *Physiother Can* 1989; 41: 304-11.
41. Watson MJ. Refining the Ten-Metre Walking Test for use with neurologically impaired people. *Physiotherapy* 2002; 88: 386-97.
42. Cowley A, Kerr K. A review of clinical balance tools for use with elderly populations. *Crit Rev Phys Rehabil Med* 2003; 15: 167–205.
43. Powell LE, Myers AM. The Activities-Specific Balance Confidence (ABC) Scale. *J Gerontol A Biol Sci Med Sci* 1995; 50: M28-34

44. Yesavage JA, Brink TL, Rose TL, Lum O, Huang V, Adey MB, Leirer VO. Development and validation of a geriatric depression screening scale: A preliminary report. *J Psychiatr Res* 1983; 17: 37-49.
45. Brink TL, Yesavage JA, Lum O, Heersema P, Adey MB, Rose TL. Screening tests for geriatric depression. *Clinical Gerontologist* 1982; 1: 37-44.
46. Train The Brain Forum Committee. Thai Geriatric Depression Scale-TGDS. *Siriraj Hosp Gaz* 1994; 46: 1-9.
47. Faul F, Erdfelder E, Lang AG, Buchner A. G*Power 3: a flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behav Res Methods* 2007; 39: 175-91.
48. Stjernman H, Tysk C, Almer S, Strö m M, Hjortswang H. Factors predicting the outcome of disease activity assessment in crohn's disease. *Inflamm Bowel Dis* 2009; 15: 1859-66.
49. Silsupadol P, Shumway-Cook A, Lugade V, van Donkelaar P, Chou LS, Mayr U, et al. Effects of single-task versus dual-task training on balance performance in older adults: a double-blind, randomized controlled trial. *Arch Phys Med Rehabil* 2009; 90: 381-7.
50. Silsupadol P, Lugade V, Shumway-Cook A, van Donkelaar P, Chou LS, Mayr U, et al. Training-related changes in dual-task walking performance of elderly persons with balance impairment: a double-blind, randomized controlled trial. *Gait Posture* 2009; 29: 634-9.
51. Kramer AF, Larish JF, Strayer DL. Training for attentional control in dual task settings: a comparison of young and old adults. *J Exp Psychol Appl* 1995; 1: 50-76.

52. Huxhold O, Li SC, Schmiedek F, Lindenberger U. Dual tasking postural control: aging and the effects of cognitive demand in conjunction with focus of attention. *Brain Res Bull* 2006; 69: 294–305



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright © by Chiang Mai University
All rights reserved