

REFERENCES

1. Blake AJ, Morgan K, Bendall MJ, Dallosso H, Ebrahim SB, Arie TH, et al. Falls by elderly people at home: prevalence and associated factors. *Age Ageing* 1988; 17: 365-72.
2. Campbell AJ, Reinken J, Allan BC, Martinez GS. Falls in old age: a study of frequency and related clinical factors. *Age Ageing* 1981; 10: 264-70.
3. World Health Organization: Falls prevention in older age; [cited 2010 Sep 12]. Available from:
http://www.who.int/ageing/projects/falls_prevention_older_age/en.
4. American Geriatrics Society, British Geriatrics Society, and American Academy of Orthopaedic Surgeons Panel on Falls Prevention. Guideline for the prevention of falls in older Persons. *J Am Geriatr Soc* 2001; 49: 664-72.
5. Soriano TA, DeCherrie LV, Thomas DC. Falls in the community-dwelling older adult: a review for primary-care providers. *Clin Interv Aging* 2007; 4: 545-53.
6. Beauchet O, Dubost V, Allali G, Gonthier R, Hermann F, Kressig R. 'Faster counting while walking' as a predictor of falls in older adults. *Age Ageing* 2007; 36: 418-23.
7. Hollman J, Kovash F, Kubik J, Linbo R. Age-related differences in spatiotemporal markers of gait stability during dual task walking. *Gait Posture* 2006; 26: 113-9.

8. Lajoie Y, Gallagher S. Predicting falls within the elderly community: comparison of postural sway, reaction time, the berg balance scale and the activities-specific balance confidence comparing fallers and non-fallers. *Arch Gerontol Geriatr* 2004; 38: 11-26.
9. Chang J, Morton S, Rubenstein L, Mojica W, Maglione M, Suttorp M, et al. Interventions for the prevention of falls in older adults: systematic review and meta-analysis of randomized clinical trials. *BMJ* 2004; 328: 680-7.
10. National Center of Injury Prevention and Control Of the Centers for Disease Control and Prevention. Preventing falls: How to develop community-based fall prevention programs for older adults. Atlanta, Georgia. 2008. p. 1-95.
11. Stevens JA, Corso PS, Finkelstein EA, Miller TR. The costs of fatal and non-fatal falls among older adults. *Inj Prev* 2006; 12: 290-5.
12. Rubenstein LZ, Robbins AS, Josephson KR, Schulman BL, Osterweil D. The value of assessing falls in an elderly population. A randomized clinical trial. *Ann Intern Med* 1990; 113: 308-16.
13. Tinetti ME, Baker DI, McAvay G, Claus EB, Garrett P, Gottschalk M, et al. A multifactorial intervention to reduce the risk of falling among elderly people living in the community. *N Engl J Med* 1994; 331: 821-7.
14. Duncan PW, Studenski S, Chandler J, Prescott B. Functional reach: predictive validity in a sample of elderly male veterans. *J Gerontol* 1992; 47: M93-8.
15. Beauchet O, Annweiler C, Allali G, Berrut G, Dubost V. Dual task-related changes in gait performance in older adults: a new way of predicting recurrent falls? *J Am Geriatr Soc* 2008; 56: 181-2.

16. Faulkner K, Redfern M, Cauley J, Landsittel D, Studenski S, Rosano C, et al. Multitasking: association between poorer performance and a history of recurrent falls. *J Am Geriatr Soc* 2007; 55: 570-6.
17. Lundin-Olsson L, Nyberg L, Gustafson Y. Attention, frailty, and falls: the effect of a manual task on basic mobility. *J Am Geriatr Soc* 1998; 46: 758-61.
18. Dubost V, Kressig RW, Gonthier R, Herrmann FR, Aminian K, Najafi B, et al. Relationships between dual-task related changes in stride velocity and stride time variability in healthy older adults. *Hum Mov Sci* 2006; 25: 372-82.
19. Priest AW, Salamon KB, Hollman JH. Age-related differences in dual task walking: a cross sectional study. *J Neuroeng Rehabil* 2008; 29: 1-8.
20. Hausdorff JM, Rios DA, Edelberg HK. Gait variability and fall risk in community-living older adults: a 1-year prospective study. *Arch Phys Med Rehabil* 2001; 82: 1050-6.
21. Springer S, Giladi N, Peretz C, Yogev G, Simon ES, Hausdorff JM. Dual-tasking effects on gait variability: the role of aging, falls, and executive Function. *Mov Disord* 2006; 21: 950-7.
22. Beauchet O, Annweiler C, Allali G, Berrut G, Herrmann F, Dubost V. Recurrent falls and dual task-related decrease in walking speed: Is there a relationship? *J Am Geriatr Soc* 2008; 56: 1265-9.
23. Hausdorff JM. Gait variability: methods, modeling and meaning. *J Neuroeng Rehabil* 2005; 19: 1-9.
24. Siu K-C, Chou L-S, Mayr U, Donkelaar Pv, Woollacott MH. Does inability to allocate attention contribute to balance constraints during gait in older adults? *J Gerontol A Biol Sci Med Sci* 2008; 63: 1364-9.

25. Siu K-C, Chou L-S, Mayr U, Donkelaar Pv, Woollacott MH. Attentional mechanisms contributing to balance constraints during gait: The effects of balance impairments. *Brain Res* 2009; 1248: 59-67.
26. Verghese J, Kuslansky G, Holtzer R, Katz M, Xue X, Buschke H, et al. Walking while talking: effect of task prioritization in the elderly. *Arch Phys Med Rehabil* 2007; 88: 50-3.
27. Yogev-Seligmann G, Rotem-Galili Y, Mirelman A, Dickstein R, Giladi N, Hausdorff JM. How does explicit prioritization alter walking during dual-task performance? Effects of age and sex on gait speed and variability. *Phys Ther* 2010; 90: 177-86.
28. Gill T, Taylor AW, Pengelly A. A population-based survey of factors relating to the prevalence of falls in older people. *Gerontology* 2005; 51: 340-5.
29. Legters K. Fear of falling. *Phys Ther* 2002; 82: 264-72.
30. Tinetti ME, Speechley M. Prevention of falls among the elderly. *N Engl J Med* 1989; 320: 1055-9.
31. Rubenstein LZ. Falls in older people: epidemiology, risk factors and strategies for prevention. *Age Ageing* 2006; 35 Suppl 2: 37-41.
32. Tinetti ME, Mendes de Leon CF, Doucette JT, Baker DI. Fear of falling and fall-related efficacy in relationship to functioning among community-living elders. *J Gerontol* 1994; 49: M140-7.
33. Tinetti ME, Speechley M, Ginter SF. Risk factors for falls among elderly persons living in the community. *N Engl J Med* 1988; 319: 1701-7.

34. Campbell AJ, Borrie MJ, Spears GF, Jackson SL, Brown JS, Fitzgerald JL. Circumstances and consequences of falls experienced by a community population 70 years and over during a prospective study. *Age Ageing* 1990; 19: 136-41
35. Sattin RW. Falls among older persons: a public health perspective. *Annu Rev Public Health* 1992; 13: 489-508.
36. Shumway-Cook A, Woollacott MH, Kerns K. A., Baldwin M. The effects of two types of cognitive tasks on postural stability in older adults with and without a history of falls. *J Gerontol A Biol Sci Med Sci* 1997; 52: M232-M40.
37. Woollacott MH, Shumway-Cook A. Attention and the control of posture and gait: a review of an emerging area of research. *Gait Posture* 2002; 16: 1-14.
38. Yogev-Seligmann G, Hausdorff JM, Giladi N. The role of executive function and attention in gait. *Mov Disord* 2008; 23: 329-42.
39. Beauchet O, Kressig RW, Najafi B, Aminian K, Dubost V, Mourey F. Age-related decline of gait control under a dual-task condition. *J Am Geriatr Soc* 2003; 51: 1187-8.
40. Toulotte C, Thevenon A, Watelain E, Fabre C. Identification of healthy elderly fallers and non-fallers by gait analysis under dual-task conditions. *Clin Rehabil* 2006; 20: 269-76.
41. Ebersbach G, Dimitrijevic MR, Poewe W. Influence of concurrent tasks on gait: a dual-task approach. *Percept Mot Skills* 1995; 81: 107-13.
42. Paul L, Ellis BM, Leese GP, McFadyen AK, McMurray B. The effect of a cognitive or motor task on gait parameters of diabetic patients, with and without neuropathy. *Diabet Med* 2009; 26: 234-9.

43. Armieri A, Holmes JD, Spaulding SJ, Jenkins ME, Johnson AM. Dual task performance in a healthy young adult population: results from a symmetric manipulation of task complexity and articulation. *Gait Posture* 2009; 29: 346-8.
44. Allali G, Kressig RW, Assal F, Herrmann FR, Dubost V, Beauchet O. Changes in gait while backward counting in demented older adults with frontal lobe dysfunction. *Gait Posture* 2007; 26: 572-6.
45. Shumway-Cook A, Baldwin M, Polissar NL, Gruber W. Predicting the probability for falls in community-dwelling older adults. *Phys Ther* 1997; 77: 812-9.
46. Silsupadol P, Shumway-Cook A, Lugade V, van Donkelaar P, Chou LS, Mayr U, et al. Effects of single-task versus dual-task training on balance performance in older adults: a double-blind, randomized controlled trial. *Arch Phys Med Rehabil* 2009; 90: 381-7.
47. Wolf B, Feys H, De W, van der Meer J, Noom M, Aufdemkampe G. Effect of a physical therapeutic intervention for balance problems in the elderly: a single-blind, randomized, controlled multicentre trial. *Clin Rehabil* 2001; 15: 624-36.
48. Siu KC, Lugade V, Chou LS, van Donkelaar P, Woollacott MH. Dual-task interference during obstacle clearance in healthy and balance-impaired older adults. *Aging Clin Exp Res* 2008; 20: 349-54.
49. แบบทดสอบสภาพสมองเบื้องต้น ฉบับภาษาไทย MMSE-Thai 2002. คณะกรรมการจัดทำแบบทดสอบสภาพสมองเบื้องต้น ฉบับภาษาไทย พ.ศ.2542. สถาบันเวชศาสตร์ผู้สูงอายุ กรมการแพทย์ กระทรวงสาธารณสุข.

50. Beck AT, Ward CH, Mendelson M, Mock J, Erbaugh J. An inventory for measuring depression. *Arch Gen Psychiatry* 1961; 4: 561-71.
51. ลัดดา แสนสีหา. ความซึมเศร้า และความคิดอัตโนมัติด้านลบของวัยรุ่นตอนปลาย. วิทยานิพนธ์ปริญญาพยาบาลศาสตรมหาบัณฑิต สาขาวิชาสุขภาพจิตและการพยาบาลจิตเวช บัณฑิตวิทยาลัย มหาวิทยาลัยเชียงใหม่, 2536.
52. Srygley JM, Mirelman A, Herman T, Giladi N, Hausdorff JM. When does walking alter thinking? Age and task associated findings. *Brain Res* 2009; 1253: 92-9.
53. Huxhold O, Li SC, Schmiedek F, Lindenberger U. Dual-tasking postural control: aging and the effects of cognitive demand in conjunction with focus of attention. *Brain Res Bull* 2006; 69: 294-305.
54. Patla AE, Rietdyk S. Visual control of limb trajectory over obstacles during locomotion: effect of obstacle height and width. *Gait Posture* 1993; 1: 45-60.