

TABLE OF CONTENTS

	PAGE
ACKNOWLEDGEMENTS	iii
ABSTRACT (THAI)	iv
ABSTRACT (ENGLISH)	vi
LIST OF TABLES	x
LIST OF FIGURES	xi
CHAPTER I INTRODUCTION	
1.1 Statement and significance of the problem	1
1.2 Anticipated benefits	2
CHAPTER II LITERATURE REVIEW	
2.1 Orthodontic anchorage and its control	3
2.2 Development of temporary anchorage devices (TADs) especially the miniscrew implant as an orthodontic anchorage	4
2.3 Stability assessments of the miniscrew implant	5
2.4 Gingival crevicular fluid (GCF), Peri-implant crevicular fluid (PICF), and Peri-miniscrew implant crevicular fluid (PMICF)	16
2.5 Detection of glycosaminoglycan (GAG) in GCF and PICF by Enzyme-linked immunosorbent assay (ELISA) with monoclonal antibody (mAb)	17
CHAPTER III MATERIALS AND METHODS	
3.1 Materials	20
3.2 Methods	22

	PAGE
CHAPTER IV RESULTS	
4.1 Part I : Five-week period	29
4.2 Part II: Ten-week period	31
CHAPTER V DISCUSSION	35
CHAPTER VI CONCLUSION	39
BIBLIOGRAPHY	40
APPENDIX	50
CIRRICULUM VITAE	54

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
 Copyright© by Chiang Mai University
 All rights reserved

LIST OF TABLES

TABLE	PAGE
A.1 CS epitope (WF6 epitope) levels (ng/ml) and the median CS epitope (WF6 epitope) levels (ng/ml) in PMICF samples during the unloaded and the loaded period from ten miniscrew implants in Part I	50
A.2 Distribution of CS epitope (WF6 epitope) levels in PMICF samples collected in Part I	50
A.3 Comparison of CS epitope (WF6 epitope) levels (ng/ml) between the unloaded and the loaded group in Part I	51
A.4 CS epitope (WF6 epitope) levels (ng/ml) and the median CS epitope (WF6 epitope) levels (ng/ml) in PMICF samples during the unloaded and the loaded period from ten miniscrew implants in Part II	52
A.5 Distribution of CS epitope (WF6 epitope) levels in PMICF samples collected in Part II	52
A.6 Comparison of CS epitope (WF6 epitope) levels (ng/ml) between the unloaded and the loaded group in Part II	53

LIST OF FIGURES

FIGURE	PAGE
3.1 Diagram of the experimental design	23
3.2 Miniscrew implant placement procedure	24
3.3 Sentalloy closed coil spring (50 g) was used to connect the miniscrew implant head and the canine bracket	25
3.4 Peri-miniscrew implant crevicular fluid (PMICF) collection.	26
3.5 Clinical mobility assessment of the miniscrew implant	26
4.1 The median CS epitope (WF6 epitope) levels (ng/ml) in PMICF samples during the unloaded (n = 40) and the loaded (n = 38) periods	30
4.2 Distribution of CS epitope (WF6 epitope) levels in Part I of the experiment (five-week period) during the unloaded (n = 40) and the loaded (n = 38) periods	31
4.3 The median CS epitope (WF6 epitope) levels (ng/ml) in PMICF samples during the unloaded (n = 40) and the loaded (n = 82) periods	33
4.4 Distribution of CS epitope (WF6 epitope) levels in Part II of the experiment (ten-week period) during the unloaded (n = 40) and the loaded (n = 82) periods	34