

BIBLIOGRAPHY

- Ahmad NN, Ala-Kokko L, Knowlton RG, Weaver EJ, Maguire JI, Tasman W, Prockop DJ. A stop codon in the gene for type II procollagen (*COL2A1*) causes one variant of arthro-ophthalmopathy (the Stickler syndrome). *Am J Hum Genet* 1990; 47 (suppl.): A206 only.
- Ahmad NN, Ala-Kokko L, Knowlton RG, Jimenez SA, Weaver EJ, Maguire JI, Tasman W, Prockop DJ. Stop codon in the procollagen II gene (*COL2A1*) in a family with the Stickler syndrome (arthro-ophthalmopathy). *Proc Nat Acad Sci* 1991; 88: 6624-6627.
- Ahmad W, Brancolini V, ul Haque FM, Lam H, ul Hague HS, Haider M, Maimon A, Aita VM, Owen J, Brown D, Zegarelli DJ, Ahmad M, Ott J, and Christiano AM. A locus for autosomal recessive hypodontia with associated dental anomalies maps to chromosome 16q12.1. *Am J Hum Genet* 1998; 62: 987-991.
- Alkuraya FS, Saadi I, Lund JJ, Turbe-Doan A, Morton CC, Mass RL. *SUMO1* haploinsufficiency leads to cleft lip palate. *Science* 2006; 313: 1751.
- Andreou AM, Pauws E, Jones MC, Singh MK, Bussen M, Douney K, Moore GE, Kispert A, Brosens JJ, Stanier P. *TBX22* missense mutations found in patients with X-linked cleft palate affect DNA binding, sumoylation, and transcriptional repression. *Am J Hum Genet* 2007; 81: 700-712.
- Ardinger HH, Beutow KH, Bell GI. Association of genetic variation of the transforming growth factor alpha gene with cleft lip and palate. *Am J Hum Genet* 1989; 45: 34-53.

Baccetti T. A controlled study of associated dental anomalies. *Angle Orthod* 1998; 68: 267-274.

Ballard JL, Auer CE, Khoury JC. Ankyloglossia: Assessment, incidence, and effect of frenuloplasty on the breast-feeding dyad. *Pediatrics* 2002; 110: e63.

Bayes M, Hartung AJ, Ezer S, Pispa J, Thesleff I, Srivastava AK. The anhidrotic ectodermal dysplasia gene (EDA) undergoes alternative splicing and encodes ectodysplasin-A with deletion mutations in collagenous repeats. *Hum Mol Genet* 1998; 7: 1661-1669.

Bender PL. Genetics of cleft lip and palate. *J Pediat Nurs* 2000; 15: 242-249.

Bitgood MJ, McMahon AP. Hedgehog and *Bmp* genes are coexpressed at many diverse sites of cell-cell interaction in the mouse embryo. *Dev Biol* 1995; 172: 126-138.

Bittencourt MA, Bolognese AM. Epithelial alterations of secondary palate formation. *Braz Dent J* 2000; 11: 117-126.

Bougeard G, Hadj-Rabia S, Faivre L, Sarafan-Vasseur N, Frebourg T. The Rapp-Hodgkin syndrome results from mutations of the *TP63* gene. *Eur J Hum Genet* 2003; 11: 1700-1704.

Braybrook C, Doudney K, Marciano ACB, Arnason A, Bjornsson A, Patton MA, Goodfellow PJ, Moore GE, Stanier P. The T-box transcription factor gene *TBX22* is mutated in X-linked cleft palate and ankyloglossia. *Nat Genet* 2001; 29: 179-183.

Braybrook C, Lisgo S, Doudney K, Henderson D, Marciano ACB, Strachan T, Patton MA, Villard L, Moore GE, Stanier P, Lindsay S. Craniofacial expression of

human and murine *TBX22* correlates with the cleft palate and ankyloglossia phenotype observed in CPX patients. *Hum Mol Genet* 2002; 11: 2793-2804.

Brunet CL, Sharpe PM, Ferguson MW. Inhibition of *TGF-b3* (but not *TGF-b1* and *TGF-b2*) activity prevents normal mouse embryonic palate fusion. *Int J Dev Biol* 1995; 39: 345–355.

Buehler BA, Rao V, Finnell RH. Biochemical and molecular teratology of fetal hydantoin syndrome. *Neurol Clin* 1994; 12: 741-748.

Burdick AB, Lian M, Zhuohua D, Ning G. Van der Woude syndrome in two families in Chaina. *J Craniofac Genet Dev Biol* 1987; 7: 413-418.

Carette MJ, Ferguson MW. The fate of medial edge epithelial cells during palatal fusion in vitro: an analysis by Dil labelling and confocal microscopy. *Development* 1992; 114: 379-388.

Carinci F, Scapoli L, Palmieri A, Zollino I, Pezzetti F. Human genetic factors in non-syndromic cleft lip and palate: An update. *Int J Pediatr Otorhinolaryngol* 2007; 71: 1509-1519.

Carlson BM. Human embryology and developmental biology, 2004.

Carreira S, Dexter TJ, Yavuzer U, Easty DJ, Goding CR. Brachyury-related transcription factor *Tbx2* and repression of the melanocyte-specific TRP-1 promoter. *Mol Cell Biol* 1998; 18: 5099-5108.

Carroll DK, Carroll JS, Leong CO, Cheng F, Brown M, Mills AA, Brugge JS, Ellisen LW. *p63* regulates an adhesion programme and cell survival in epithelial cells. *Nat Cell Biol* 2006; 8: 551-561.

Casey ES, O'Reilly MA, Conlon RL, Smith JC. The T-box transcription factor Brachyury regulates expression of *eFGF* through binding to a non-palindromic response element. *Development* 1998;125: 3887-3894.

Celli J, Dujif P, Hamel BC, Bamshad M, Kramer B, Smits AP, Newbury- Ecob R, Hennekam RC, van Buggenhout G, van Haeringen A, Qoods CG, van Essen AJ, de Waal R, Vriend G, Haber DA, Yang A, McKeon F, Brunner HG, van Bokhoven H. Heterozygous germline mutations in the p53 homolog p63 are the cause of EEC syndrome. *Cell* 1999; 99: 143-153.

Chaabouni M, Smaoui N, Benneji N, M'rad R, Jemaa LB, Hachicha S, Chaabouni H. Mutation analysis of *TBX22* reveals new mutation in Tunisian CPX family. *Clin Dysmorphol* 2004; 41: 68-74.

Chi S, Godfrey K. Cleft lip and cleft palate in New South Wales. *The Medical J Australia* 1970; 1172.

Chuangsuwanich A, Aojanepong C, Muangsombut S, Tongpiew P. Epidemiology of cleft lip and palate in Thailand. *Ann Plast Surg* 1998; 41: 7-10.

Cobourne MT, Hardcastle Z, Sharpe PT. Sonic hedgehog regulates epithelial proliferation and cell survival in the developing tooth germ. *J Dent Res* 2001; 80: 1974-1979.

Cobourne MT. The complex genetics of cleft lip and palate. *Eur J Orthod* 2004; 26: 7-16.

Cuervo R, Valencia C, Chandraratna RA, Covarrubias L. Programmed cell death is required for palate shelf fusion and is regulated by retinoic acid. *Dev Biol* 2002; 245: 145-156.

Cui XM, Warburton D, Zhao J, Crowe DL, Shuler CF. Immunohistochemical localization of TGFB type II receptor and *TGFB3* during palatogenesis *in vivo* and *in vitro*. *Int J Dev Biol* 1998; 42: 817-820.

Culiat CT, Stubbs L, Nicholls RD, Montgomery CS, Russell LB, Johnson DK, Rinchik EM. Concordance between isolated cleft palate in mice and alterations within a region including the gene encoding the $\beta 3$ subunit of the type A γ -aminobutyric acid receptor. *Proc Natl Acad Sci U S A* 1993; 90: 5105–5109.

Culiat CT, Stubbs LJ, Woychik RP, Russell LB, Johnson DK, Rinchik EM. Deficiency of the $\beta 3$ subunit of the type A γ -aminobutyric acid receptor causes cleft palate in mice. *Nat Genet* 1995; 11: 344–346.

Das P, Stockton DW, Bauer C, Shaffer LG, D'Souza RN, Wright T. Haploinsufficiency of *PAX9* is associated with autosomal dominant hypodontia. *Hum Genet* 2002; 110: 371-376.

Daugaad-Jensen J, Nodal M, Skovgaard LT, Kjaer I. Comparison of the pattern of agenesis in the primary and permanent dentitions in a population characterized by agenesis in the primary dentition. *Int J Pediatr Dent* 1997; 7: 143-148.

Davis PJ. Hypodontia and hyperdontia of permanent teeth in Hong Kong schoolchildren. *Community Dent Oral Epidemiol* 1987; 15: 218-220.

Dechkunakorn S, Chaiwat J, Sawaengkit P. Congenital absence and loss of teeth in an orthodontic patient group. *J Dent Assoc Thai* 1990; 40: 165-176.

Derijcke A, Eerens A, Carels C. The incidence of orofacial clefts: a review. *British J Oral and Maxillofac Surg* 1996; 34: 488-494.

Ding H, Wu X, Boström H, Kim I, Wong N, Tsoi B, O'Rourke M, Koh GH, Soriano P, Betsholtz C, Hart TC, Marazita ML, Field LL, Tam PPL, Nagy A. A specific

- requirement for *PDGF-C* in palate formation and *PDGFR- α* signaling. *Nat Genet* 2004; 36: 1111–1116.
- Dobrovolskaia-Zavadksis N. Sur la mortification spontanee de la queue chez la souris nouveau-née et sur l'existence d'un caractere heriditaire non-viable. *C R Soc Biol* 1927; 97: 114-116.
- Dudas M, Li WY, Kim J, Yang A, Kaartinen V. Palatal fusion- Where do the midline cells go? A review on cleft palate, a major human birth defect. *Acta histochemica* 2007; 109: 1-14.
- Endo T, Ozoe R, Yoshino S, Shimooka S. Hypodontia patterns and variations in craniofacial morphology in Japanese orthodontic patients. *Angle Orthod* 2006; 76: 996-1003.
- Erlich HA, Arnheim N. Genetic analysis using the polymerase chain reaction. *Ann Rev Genet* 1992; 26: 479-506.
- Ferguson MWJ, Honig LS. Epithelial-mesenchymal interaction during vertebrate palatogenesis. *Curr Top Dev Biol* 1984; 19: 137-164.
- Ferguson MWJ. Palate development. *Development* 1988; 103: 41–61.
- Ferrante MI, Giorgio G, Feather SA, Bulfone A, Wright V, Ghiani M, Selicorni A, Gammaro L, Scolari F, Woolf AS, Sylvie O, Le Marec B, Malcolm S, Winter R, Ballabio A, Franco B. Identification of the gene for oral-facial-digital type I syndrome. *Am J Hum Genet* 2001; 68: 569-576.
- Fiechett JE, Hay ED. Medial edge epithelium transforms to mesenchyme after embryonic palatal shelves fuse. *Dev Biol* 1989; 131: 455-474.
- Fitzpatrick DR, Dnhez F, Kondaiah P, Akhurst RJ. Differential expression of *TGF β* isoforms in murine palatogenesis. *Development* 1990; 109: 585-595.

Fogh-Andersen P. Incidence of cleft lip and palate: constant or increasing? *Acta Chir Scand* 1961; 122: 106–111.

Freidman HI, Sayetta RB, Coston GN, Hussey JR. Symbolic representation of cleft lip and palate. *Cleft Palate Craniofac J* 1991; 28: 252-260.

Godfrey K, Chowchuen B. Investigating the concurrences of oral clefts and other related birth defects. *Srinagarind Med J* 2001; 16: 27-36.

Gorlin RJ, Cohen NM, Hannekam R. syndromes of the had and neck. Oxford University Press, New York 2001.

Gorski SM, Adams KJ, Birch PH, Friedman JM, Goodfellow PJ. The gene responsible for X-linked cleft palate (CPX) in a British Columbian native kindred is localized between PGK1 and DXYS1. *Am J Hum Genet* 1992; 50: 1129-1136.

Gritli-Linde A. Molecular control of secondary palate development. *Developmental Biology* 2007; 301: 309–326.

Hall HE, Chan ER, Collins A, Judis L, Shirley S, Surti U, Hoffner L, Cockwell AE, Jacobs PA, Hassold TJ. The origin of trisomy 13. *Am J Med Genet* 2007; 143: 2242-2248.

Hecht JT, Wang Y, Blanton SH, Daiger SP. Van der Woude syndrome and non-syndromic cleft lip and palate. *Am J Hum Genet* 1992; 51: 442-444.

Hillard SA, Yu L, Gu S, Zhang Z, Chen YP. Regional regulation of palatal growth and patterning along the anterior-posterior axis in mice. *J Anat* 2005; 207: 655-667.

Hocevar-Boltezar I. Ear, nose and voice problems in children with orofacial clefts. *J Laryngol Otol* 2006; 120: 276-281.

Holtgrave EA, Stoltenburg-Didinger G. Apoptotic epithelial cell death: a prerequisite for palatal fusion. An in vivo study in rabbits. *J Craniomaxillofac Surg* 2002; 30: 329-336.

Homanics GE, DeLorey TM, Firestone LL, Quinlan JJ, Handforth A, Harrison NL, Krasowski MD, Rick CE, Korpi ER, Mäkelä R, Brilliant MH, Hagiwara N, Ferguson C, Snyder K, Olsen RW. Mice devoid of γ -aminobutyrate type A receptor $\beta 3$ subunit have epilepsy, cleft palate, and hypersensitive behavior. *Proc Natl Acad Sci U S A* 1997; 94: 4143-4148.

Hsueh YP, Want TF, Yang FC, Sheng M. Nuclear translocation and transcription regulation by the membrane-associated guanylate kinase CASK/LIN-2. *Nature* 2000; 404: 298-302.

Javinen S, Lehtinen L. Supernumerary and congenitally missing primary teeth in Finish children. An epidemiologic study. *Acta Odontol Scand* 1981; 39: 83-86.

Jumlongras D, Bei M, Stimson JM, Wang WF, DePalma SR, Seidman CE. A nonsense mutation in *MSX1* causes Witkop syndrome. *Am J Hum Genet* 2001; 69: 67-74.

Jumlongras D, Lin JY, Chapra A, Seidman CE, Seidman JG, Maas RL. A novel missense mutation in the paired domain of *PAX9* causes non-syndromic oligodontia. *Hum Genet* 2004; 114: 242-249.

Kaartinen V, Voncken JW, Shule C, Warburton DBD, Heisterkamp N, Groffen J. Abnormal lung development and cleft palate in mice lacking *TGFB3* indicates defects of epithelial-mesenchymal interaction. *Nat Genet* 1995; 11: 415-421.

Kaartinen V, Cui X, Heisterkamp N, Groffen J, Shuler CF. Transforming growth factor-B3 regulates transdifferentiation of medial edge epithelium during palatal

- fusion and associated degradation of the basement membrane. *Dev Dyn* 1997; 209: 255-260.
- Kanno K, Suzuki Y, Yamada A, Aoki Y, Kure S. Association between non-syndromic cleft lip with or without cleft palate and the glutamic acid decarboxylase 67 gene in the Japanese population. *Am J Med Genet* 2004; 127: 11-16.
- Kantaputra PN, Sumitsawan Y, Scutte BC, Tochareontanaphol C. Van der Woude Syndrome with sensorineural hearing loss, large craniofacial sinuses, dental pulp stones, and minor limb anomalies. *Am J Med Genet* 2002; 108: 275-280.
- Kapadia H, Frazier-Bowers S, Ogawa T, D'Souza RN. Molecular characterization of a novel *PAX9* missense mutation causing posterior tooth agenesis. *Eur J Hum Genet* 2006; 14: 403-409.
- Kayano S, Kure S, Suzuki Y, Kanno K, Aoki Y, Kondo S, Schutte BC, Murray JC, Yamada A, Matsubara Y. Novel *IRF6* mutations in Japanese patients with van der Woude syndrome: two missense mutations (R45Q and P396S) and a 17-kb deletion. *J Hum Genet* 2003; 48: 622-628.
- Kernahan DA, Stark RB. A new classification for cleft lip and palate. *Plast Reconstr Surg* 1958; 22: 435.
- Kim NY, Baek SH. Cleft sidedness and congenitally missing or malformed permanent maxillary lateral incisors in Korean patients with unilateral cleft lip and alveolus or unilateral cleft lip and palate. *Am J Orthod Dentofac Orthop* 2006; 130: 752-758.
- Kispert A, Herrmann BG. The Brachyury gene encodes a novel DNA binding protein. *EMBO J* 1993; 12: 3211-3220.

- Klein ML, Nieminen P, Lammi L, Niebuhr E, Kreiborg S. Novel mutation of the initiation codon of *PAX9* causes oligodontia. *J Dent Res* 2005; 84: 43-47.
- Kondo S, Schtte BC, Richardson RJ, Bjork BC, Knight AS, Watanabe Y, Howard E, Ferreira de Lima RLL, Daack-Hirsch S, Sander A, McDonald-McGinn DM, Zackai EH, and 14 others. Mutation in *IRF6* cause Van der Woude and Popliteal pterygium syndromes. (Letter) *Nature Genet* 2002; 32: 285-289.
- Koster MI, Kim S, Mills AA, DeMayo FG, Roop DR. *p63* is the molecular switch for initiation of an epithelial stratification program. *Genes Dev Growth Differ* 2004; 18: 126–131.
- Kotlow LA. Ankyloglossia (tongue-tie): A diagnostic and treatment quandary. *Quintessence Int* 1999; 30: 259-262.
- Kulkarni AB, Huh C, Becker D, Geiser A, Lyght M, Flanders KC, Roberts AB, Sporn MB, Ward JM, Karlsson S. Transforming growth factor- β 1 null mutation in mice causes excessive inflammatory response and early death. *Proc Natl Acad Sci USA* 1993; 90: 770–774.
- Kupietzky A, Botzer E. Ankyloglossia in the infant and young child: clinical suggestions for diagnosis and management. *Pediatr Dent* 2005; 27: 40-46.
- Lalakia ML, Messner AH. Ankyloglossia: does it matter? *Pediatr Clin N Am* 2003; 50: 381-397.
- Lammi L, Halonen K, Pirinen S, Thesleff I, Arte S, Nieminen P. A missense mutation in *PAX9* in a family with distinct phenotype of oligodontia. *Eur J Hum Genet* 2003; 11: 866-871.

- Lan Y, Ovitt CE, Cho ES, Maltby KM, Wang Q, Jiang R. Odd-skipped related (*Osr2*) encodes a key intrinsic regulator of secondary palate growth and morphogenesis. *Development* 2004; 131: 3207-3216.
- Laugier-Anfossi F, Villard L. Molecular characterisation of a new human T-box gene (*TBX22*) located in Xq21.1 encoding a protein containing a truncated T-domain. *Gene* 2000; 255: 289-296.
- Leoyklang P, Siriwan P, Shotelersuk V. A mutation of the *p63* gene in non-syndromic cleft lip. *J Med Genet* 2006; 43: e28.
- Lewis R. Human genetics concepts and applications. 2007: 377-383.
- Li QY, Newbury-Ecob RA, Terrett JA, Wilson DI, Curtis ARJ, Yi CH, Gebuhr T, Bullen PJ, Robson SC, Strachan T, Bonnet D, Lyonnet S, Young ID, Raeburn JA, Buckler AJ, Law DJ, Brook JD. Holt-Oram syndrome is caused by mutations in TBX5, a member of the Brachyury (T) gene family. *Nat Genet*. 1997;15:21-9.
- Lidral AC, Murray JC, Beutow KH, Basart AM, Schearer H, Shiang R, Naval A, Layda E, Magee K, Magee W. Studies of the candidate genes *TGFB2*, *MSX1*, *TGFA*, and *TGFB3* in the etiology of cleft lip and palate in the Philippines. *Cleft Palate Craniofac J* 1997; 34: 1-6.
- Lidral AC, Romitti PA, Basart AM, Doetschman T, Leysens NJ, Daack-Hirsch S, Semina EV, Johnson LR, Machida J, Burds A, Parnell TJ, Rubenstein JL, Murray JC. Association of *MSX1* and *TGFB3* with non-syndromic clefting in humans. *Am J Hum Genet* 1998; 63: 557-568.
- Lidral AC, Reising BC. The role of *MSX1* in human tooth agenesis. *J Dent Res* 2002; 81: 274-278.

Lowry RB. Sex linked cleft palate in a British Columbia Indian family. *Pediatrics* 1970; 46: 123-128.

Lu DW, Shi B, Wang HJ, Zheng Q. The comparative study of craniofacial structural characteristic of individuals with different types of cleft palate. *Ann Plast Surg* 2007; 59: 382-387.

Marcano ACB, Doudney K, Braybrook C, Squires R, Patton MA, Lees M, Richieri-Costa A, Lideral AC, Murray JC, Moore GE, Stanier P. *TBX22* mutations are a frequent cause of cleft palate. *J Med Genet* 2004; 41: 68-74.

Martinez-Alvarez C, Tudela C, Perez MJ, O'Kane S, Puerta J, Ferguson MW. Medial edge epithelial cell fate during palatal fusion. *Dev Biol* 2000; 220: 343-357.

Martinez-Alvarez C, Blanco MJ, Perez R, Rabadan MA, Aparicio M, Resel E, Martinez T, Nieto MA. Snail family members and cell survival in physiological and pathological cleft palates. *Dev Biol* 2004; 265: 207-218.

Mattheeuws N, Dermaut L, and Martens G. Has hypodontia increased in Caucasians during the 20th century? A meta-analysis. *Eur J Orthod* 2004; 26: 99-103.

McGrath JA, Dujif PH, Doetsch V, Irvine AD, de Waal R, Vanmolkot KR, Wessagowit V, Kelly A, Atherton DJ, Griffiths WA, Orlows SJ, van Haeringen A, Ausems MG, Yang A, McKeon F, Bamshad MA, Brunner HG, Hamel BC, van Bokhoven H. Hay-Wells syndrome is caused by heterozygous missense mutations in the SAM domain of *p63*. *Hum Mol Genet* 2001; 10: 221–229.

Mcleod NHM, Arana-Urioste ML, Saeed NR. Birth prevalene of cleft lip and palate in sucre, Bolivia. *Cleft Palate Craniofac J* 2004; 41: 195-198.

Messner AH, Lalakea ML. Ankyloglossia: incidence and associated feeding difficulties. *Arch Otolaryngol Head Neck Surg* 2000; 126: 36-39.

- Mills A, Zheng B, Wang XJ, Vogel H, Roop DR, Bradley A. *p63* is a *p53* homologue required for limb and epidermal morphogenesis. *Nature* 1999; 398: 708–713.
- Moore GE, Ivens A, Chambers J, Bjornsson A, Arnason A, Jensson O, Williamson R. The application of molecular genetics to detection of craniofacial abnormality. *Development* 1988; 103: 233-239.
- Mossey P. Epidemiology underpinning research in the aetiology of orofacial clefts. *Orthod Craniofac Res* 2007; 10: 114-120.
- Mostowska A, Biedziak B, Trzeciak WH. A novel c.581C>T transition localized in a highly conserved homeobox sequence of *MSX1*: is it responsible for oligodontia? *J Appl Genet* 2006; 47: 159-164.
- Munger RG, Romitti PA, Daack-Hirsch S, Burns TL, Murray JC, Hanson J. Maternal alcohol use and risk of orofacial cleft birth defects. *Teratology* 1996; 54: 27-33.
- Murray JC D, aack-Hirsch S, Beutow KH. Clinical and epidemiologic studies of cleft lip and palate in the Phillipines. *Cleft Palate Craniofac J* 1997; 34: 7-10.
- Natsume N, Kawaia T, Kohamab G, Teshimac T, Kochic S, Ohashid Y, Enomotoe S, Ishiiie M, Shigematsu T (deceased), Nakano Y, Matsuyag T, Kogog M, Yoshimurah Y, Ohishii M, Nakamura N, Katsukij T, Gotoj M, Shimizuk M, Yanagisawak S, Mimural T, Sunakawam H. Incidence of cleft lip or palate in 303738 Japanese babies born between 1994 and 1995. *British J Oral and Maxillofac Surg* 2000; 38: 605-607.
- Nawshad A, Lagamba D, Hay ED. Transforming growth factor beta (TGFbeta) signalling in palatal growth, apoptosis and epithelial mesenchymal transformation (EMT). *Arch Oral Biol* 2004; 49: 675-589.

- Nieminen P, Pekkanen M, Åberg T, and Thesleff I. A graphical www-database on gene expression in tooth. *European Journal of Oral Sciences* 1998; 106: 7-11.
- Nieminen P, Arte S, Tanner D, Paulin L, Alaluusua S, Thesleff I. Identification of a nonsense mutation in the *PAX9* gene in molar oligodontia. *Eur J Hum Genet* 2001; 9: 743-746.
- Nik-Hussein NN. Hypodontia in the permanent dentition: a study of its prevalence in Malaysian children. *Aust Orthod J* 1989; 11: 93-95.
- Nodal M, Kjaer I, and Solow B. Craniofacial morphology in patients with multiple congenitally missing permanent teeth. *Eur J Orthod* 1994; 16: 104-109.
- Packham EA, Brook JD. T-box genes in human disorders. *Hum Mol Genet* 2003; 12: 37-44.
- Pelton RW, Hogan BL, Miller DA, Moses HL. Differential expression of genes encoding TGF- β 1, β 2 and β 3 during murine palate formation. *Dev Biol* 1990; 141: 456-460.
- Persson C, Lohmander A, Elander A. Speech in children with an isolated cleft palate: a longitudinal perspective. *Cleft Palate Craniofac J* 2006; 43: 295-309.
- Polder BJ, Hof MAV, Van der Linder FPGM, Kuijpers-Jagtman AM. A meta-analysis of the prevalence of dental agenesis of permanent teeth. *Community Dent Oral Epidemiol* 2004; 32: 217-226.
- Proetzel G, Pawlowski SA, Wiles MV, Yin M, Boivin GP, Howles PN, Ding J, Ferguson MWJ, Doetschman T. Transforming growth factor- β 3 is required for secondary palate fusion. *Nat Genet* 1995; 11: 409-414.
- Rahoma AH. Complete cleft palate: a new scheme of treatment. *The Int J Plastic Surg* 2007 ;4.

- Rattanasiri T, Junthathamrongwat N, Apiwan thanakul S, Wongkam C, Chowchuen B. The incidence of cleft lip and palate in Srinakarind hospital, 1990-1999. *Srinakarind Med J* 2001; 16: 3-7.
- Reardon W, Winter RM, Rutland P, Pulleyn LJ, Jones BM, Malcolm S. Mutations in the fibroblast growth factor receptor 2 gene cause Crouzon syndrome. *Nature Genet* 1994; 8: 98-103.
- Rice R, Spencer-Dene B, Connor EC, Gritli-Linde A, McMahon AP, Dickson C, Thesleff I, Rice DP. Disruption of *Fgf10/Fgfr2b*-coordinated epithelial-mesenchymal interactions causes cleft palate. *J Clin Invest* 2004; 113: 1692-1700.
- Rice R, Connor E, Rice DPC. Expression patterns of hedgehog signaling pathway members during palate development. *Gene Expression Patterns* 2006; 6: 206-221.
- Richard ME. Feeding the newborn with cleft lip and/or palate: the enlargement, stimulate, swallow, rest (ESSR) method. *J Pediatr Nurs* 1991; 6: 317-321.
- Ritthakol W. The incidence of cleft lip and palate in Songklanakarin hospital. *J Dent Assoc Thai* 2001; 51: 29-37.
- Rizos M, Spyopoulos MN. Van der Woude syndrome: a review. Cardinal signs, epidemiology, associated features, differential diagnosis, expressivity, genetic counselling and treatment. *Eur J Orthod* 2004; 26: 17-24.
- Ruangritt C, Phrasertthang P, Banpho Y, Lamduan W, Giathamnuay S, Nuwantha A. Incidence cleft lip and cleft palate in 3 hospitals in Khon Kaen. Khon kaen : Department of orthodontics, Faculty of Dentistry, Khon Kaen University. 1993.
- Sanford LP, Ormsby I, Groot ACG, Sariola H, Friedman R, Boivin GP, Cardell EL, Doetschman T. *TGF-β2* knockout mice have multiple developmental defects that

are nonoverlapping with other *TGF-β* knockout phenotype. *Development* 1997; 124: 2659–2670.

Scapoli L, Martinelli M, Pezzetti F, Carinci F, Bodo M, Tognon M, Carinci P. Linkage disequilibrium between *GABRB3* gene and non-syndromic familial cleft lip with or without cleft palate. *Hum Genet* 2002; 110: 15–20.

Schupbach PM, Schroeder HE. Cell release from the palatal shelves and the fusion line. *J Biol Buccale* 1983; 11: 227-241.

Schutte BC, Murray JC. The many faces and factors of orofacial clefts. *Hum Mol Genet* 1999; 8: 1853-1859.

Seielstad M, Bekele E, Ibrahim M, Toure A, Traore M. Aview of modern human origin from Y chromosome microsatellite variation. *Genome Research* 1999; 9: 558-567.

Semina EV, Reiter R, Leysens NJ, Alward WL, Small KW, Datson NA. Cloning and characterization of a novel bicoid-related homeobox transcription factor gene, RIEG, involved in Rieger syndrome. *Nat Genet* 1996; 14: 392-399.

Shaw GM, Wasserman CR, Lammer EJ, O'Malley CD, Murray JC, Basart AM,Tolarova MM. Orofacial clefts, parental cigarette smoking, and transforming growth factor-alpha gene variants. *Am J Hum Genet* 1996; 58: 551–561.

Shotelersuk V, Janklat S, Siriwan P, Tongkobpatch S. De novo missense mutation, S541Y, in the *p63* gene underlying Rapp-Hodgkin ectodermal dysplasia syndrome. *Clin Exp Dermatol* 2005; 30: 282–285.

- Shuler CF, Guo Y, Majumder A, Luo RY. Molecular and morphologic changes during the epithelial-mesenchymal transformation of palatal shelf medial edge epithelium in vitro. *Int J Dev Biol* 1991; 35: 463-472.
- Shuler CF, Halpern DE, Guo Y, Sank AC. Medial edgeepithelium fate traced by cell lineage annalysis during epithelial-mesenchymeal transformation in vivo. *Dev Biol* 1992; 154: 318-330.
- Shull MM, Ormsby I, Kier AB, Pawlowski S, Diebold RJ, Yin M, Allen R, Sidman C, Proetzel G, Calvin D, Annunziata N, Doetschman T. Targeted disruption of the mouse transforming growth factor- β 1 gene results in multifocal inflammatory disease. *Nature* 1992; 359: 693–699.
- Slavotinek A, Poyser L, Wallace A, Martin F, Gaunt L, Kingston H. Two unique patients with trisomy 18 mosaicism and molecular marker studies. *Am J Med Genet* 2003; 117: 282-288.
- Slayton RL, Williums L, Murray JC, Wheeler JJ, Lidral AC, Nishimura CJ. Genetic association studies of cleft lip and/or palate with hypodontia outside the cleft region. *Cleft Pal craniofac J* 2003; 40: 274-279.
- Slootweg PJ. Dental pathology, 2007:1-4.
- Sozen MA, Suzuki K , Tolarova MM, Bustos T, Iglesias JEF, Spritz RA. Mutation of *PVRL1* is associated with sporadic, non-syndromic cleft lip/palate in northern Venezuela. *Nat Genet* 2001; 29: 141-142.
- Sperber GH. Craniofacial development, 2001.
- Stanier P, Moore GE. Genetics of cleft lip and palate: syndromic genes contribute to the incidence of non-syndromic clefts. *Hum Mol Genet* 2004; 13: R73-R81.

Stevenson RE, Hall JG. Human malformations and related anomalies. 2nd ed Oxford University Press 2006; 405-406.

Stockton DW, Das P, Goldenberg M, D'Souza RN, Patel PI. Mutation of *PAX9* is associated with oligodontia. *Nat Genet* 2000; 24: 18-19.

Suphapeetiporn, K Tongkobpatch S, Siriwan P, Shotelersuk V. *TBX22* mutations are a frequent cause of non-syndromic cleft palate in the Thai population. *Clin Genet* 2007; 72: 478-483.

Suzuki K, Hu D, Bustos T, Zlotogora J, Richieri-Costa A, Helms JA. Mutations of *PVRL1*, encoding a cell-cell adhesion molecule/herpesvirus receptor, in cleft lip/palate-ectodermal dysplasia. *Nat Genet* 2000; 25: 427-430.

Tada M, Casey ES, Fairclough L, Smith JC. *Bix1*, a direct target of xenopus T-box genes, causes formation of ventral mesoderm and endoderm. *Development* 1998; 125: 3997-4006.

Tada M, Smith JC. T-targets: clues to understanding the functions of T-box proteins. *Dev Growth Differ* 2001; 43: 1-11.

Takigawa T, Shiota K. Terminal differentiation of palatal medial edge epithelial cells in vitro is not necessarily dependent on palatal shelf contact and midline epithelial seam formation. *Int J Dev Biol* 2004; 48: 307-317.

Tan KI. Incidence and epidemiology of cleft lip and palate in Singapore. *Ann Acad Med Singapore* 1988; 17: 311-314.

Taya Y, O'Kane S, Ferguson MWJ. Pathogenesis of cleft palate in *TGF-b3* knockout mice. *Development* 1999; 126: 3869-3879.

Thesleff I. Genetic basis of tooth development and dental defects. *Acta Odontol Scand* 2000; 58: 191-194.

- Tucker A, Sharpe PT. The cutting-edge of mammalian development; how the embryo makes teeth. *Nat Rev Genet* 2004; 5: 499-508.
- Turner SR, Rumsey N, Sandy JR. Psychological aspects of cleft lip and palate. *Eur J Orthod* 1998; 20: 407-415.
- van Bokhoven H, Hamel BC, Bamshad M, Sangiorgi E, Gurrieri F, Duijf PH. p63 Gene mutations in eec syndrome, limb-mammary syndrome, and isolated split hand-split foot malformation suggest a genotype-phenotype correlation. *Am J Hum Genet* 2001; 69: 481-492.
- van den Boogaard MJ, Dorland M, Beemer FA, van Amstel HK. *MSX1* mutation is associated with orofacial clefting and tooth agenesis in humans. *Nat Genet* 2000; 24: 342-343.
- Varju P, Katarova Z, Madarasz E, Szabo G. *GABA* signaling during development: new data and old questions. *Cell Tissue Res* 2001; 305: 239–246.
- Wang X, Liu J, Zhang H, Xiao M, Li J, Yang C, Lin X, Wu Z, Hu L, Kong X. Novel mutations in the *IRF6* gene for Van der Woude syndrome. *Hum Genet* 2003; 113: 382-386.
- Wilkie AOM SS, Oldridge M, Poole MD, Ashworth GJ, Hockley AD, Hayward RD, David DJ, Pulley LJ, Rutland P, Malcolm S, Winter RM, Reardon W. Apert syndrome results from localized mutations of *FGFR2* and is allelic with Crouzon syndrome. *Nature Genet* 1995; 9: 165-172.
- Williams WN, Waldron CM. Assessment of lingual function when ankyloglossia (tongue-tie) is suspected. *J Am Dent Assoc* 1985; 110: 353-326.
- Wilson V, Colon F. The T-box family. *Genome Biology* 2002; 3: 3008.1-3008.7.
- Winter PC, Hickey GI, Fletcher HL. Genetics, 2002: 255-263.

- Wise CA, Chiang LC, Paznekas WA, Sharma M, Musy MM, Ashley JA, Lovett M, Jabs EW. *TCOF1* gene encodes a putative nucleolar phosphoprotein that exhibits mutations in Treacher Collins syndrome throughout its coding region. *Proc Nat Acad Sci* 1997; 94: 3110-3115.
- Wong FWL, King NM. A review of the rate of occurrence of cleft lip and palate in Chinese people. *HKMJ* 1997; 3: 96-100.
- Yang A SR, Sun D, Kaghad M, Walker N, Bronson RT, Tabin C, Sharpe A, Caput D, Crum C, McKeon F. *p63* is essential for regenerative proliferation in limb, craniofacial and epithelial development. *Nature* 1999; 398: 714–718.
- Yu L, Gu S, Alappat S, Song Y, Yan M, Zhang X, Zhang G, Jiang Y, Zhang Z, Zhang Y, Chen Y. *Shox2*-deficient mice exhibit a rare type of incomplete clefting of the secondary palate. *Development* 2005; 132: 4397–4406.
- Zhang Z, Song Y, Zhao X, Zhang X, Fermin C, Chen YP. Rescue of cleft palate in *Msx1*-deficient mice by transgenic *Bmp4* reveals a network of *BMP* and *Shh* signaling in the regulation of mammalian palatogenesis. *Development* 2002; 129: 4135–4146.