

REFERENCES

- [1] Mikesell P, Nusstein J, Reader A, Beck M, Weaver J. A comparison of articaine and lidocaine for inferior alveolar nerve blocks. *J Endod* 2005;31:265-70.
- [2] Lai TN, Lin CP, Kok SH, Yang PJ, Kuo YS, Lan WH, et al. Evaluation of mandibular block using a standardized method. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2006;102:462-8.
- [3] Dunbar D, Reader A, Nist R, Beck M, Meyers WJ. Anesthetic efficacy of the intraosseous injection after an inferior alveolar nerve block. *J Endod* 1996;22:481-6.
- [4] Simon F, Reader A, Drum M, Nusstein J, Beck M. A prospective, randomized single-blind study of the anesthetic efficacy of the inferior alveolar nerve block administered with a peripheral nerve stimulator. *J Endod* 2010;36:429-33.
- [5] Claffey E, Reader A, Nusstein J, Beck M, Weaver J. Anesthetic efficacy of articaine for inferior alveolar nerve blocks in patients with irreversible pulpitis. *J Endod* 2004;30:568-71.
- [6] Tortamano IP, Siviero M, Costa CG, Buscariolo IA, Armonia PL. A comparison of the anesthetic efficacy of articaine and lidocaine in patients with irreversible pulpitis. *J Endod* 2009;35:165-8.
- [7] Matthews R, Drum M, Reader A, Nusstein J, Beck M. Articaine for supplemental buccal mandibular infiltration anesthesia in patients with irreversible pulpitis when the inferior alveolar nerve block fails. *J Endod* 2009;35:343-6.

- [8] Poorni S, Veniashok B, Senthilkumar AD, Indira R, Ramachandran S. Anesthetic efficacy of four percent articaine for pulpal anesthesia by using inferior alveolar nerve block and buccal infiltration techniques in patients with irreversible pulpitis: a prospective randomized double-blind clinical trial. *J Endod* 2011;37:1603-7.
- [9] Parirokh M, Satvati SA, Sharifi R, Rekabi AR, Gorjestani H, Nakhaee N, et al. Efficacy of combining a buccal infiltration with an inferior alveolar nerve block for mandibular molars with irreversible pulpitis. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2010;109:468-73.
- [10] Kanaa MD, Whitworth JM, Meechan JG. A prospective randomized trial of different supplementary local anesthetic techniques after failure of inferior alveolar nerve block in patients with irreversible pulpitis in mandibular teeth. *J Endod* 2012;38:421-5.
- [11] Khademi AA, Saatchi M, Minaiyan M, Rostamizadeh N, Sharafi F. Effect of preoperative alprazolam on the success of inferior alveolar nerve block for teeth with irreversible pulpitis. *J Endod* 2012;38:1337-9.
- [12] Monteiro MR, Groppo FC, Haiter-Neto F, Volpato MC, Almeida JF. 4% articaine buccal infiltration versus 2% lidocaine inferior alveolar nerve block for emergency root canal treatment in mandibular molars with irreversible pulpits: a randomized clinical study. *Int Endod J* 2014.
- [13] Hargreaves KM, Keiser K. Local anesthetic failure in endodontics. *Endod Topics* 2002;1:26-39.
- [14] Srinivasan N, Kavitha M, Loganathan CS, Padmini G. Comparison of anesthetic efficacy of 4% articaine and 2% lidocaine for maxillary buccal infiltration in patients with irreversible pulpitis. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2009;107:133-6.

- [15] Aggarwal V, Singla M, Miglani S, Kohli S, Singh S. Comparative evaluation of 1.8 mL and 3.6 mL of 2% lidocaine with 1:200,000 epinephrine for inferior alveolar nerve block in patients with irreversible pulpitis: a prospective, randomized single-blind study. *J Endod* 2012;38:753-6.
- [16] Fowler S, Reader A. Is a Volume of 3.6 mL Better than 1.8 mL for Inferior Alveolar Nerve Blocks in Patients with Symptomatic Irreversible Pulpitis? *J Endod* 2013;39:970-2.
- [17] Fan S, Chen WL, Pan CB, Huang ZQ, Xian MQ, Yang ZH, et al. Anesthetic efficacy of inferior alveolar nerve block plus buccal infiltration or periodontal ligament injections with articaine in patients with irreversible pulpitis in the mandibular first molar. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2009;108:e89-93.
- [18] Kanaa MD, Whitworth JM, Corbett IP, Meechan JG. Articaine buccal infiltration enhances the effectiveness of lidocaine inferior alveolar nerve block. *Int Endod J* 2009;42:238-46.
- [19] Meechan JG. Supplementary routes to local anaesthesia. *Int Endod J* 2002;35:885-96.
- [20] Nusstein J, Kennedy S, Reader A, Beck M, Weaver J. Anesthetic efficacy of the supplemental X-tip intraosseous injection in patients with irreversible pulpitis. *J Endod* 2003;29:724-8.
- [21] Nusstein J, Claffey E, Reader A, Beck M, Weaver J. Anesthetic effectiveness of the supplemental intraligamentary injection, administered with a computer-controlled local anesthetic delivery system, in patients with irreversible pulpitis. *J Endod* 2005;31:354-8.

- [22] Feck AS, Goodchild JH. The use of anxiolytic medications to supplement local anesthesia in the anxious patient. *Compend Contin Educ Dent* 2005;26:183-6, 8, 90; quiz 91, 209.
- [23] Ianiro SR, Jeanssone BG, McNeal SF, Eleazer PD. The effect of preoperative acetaminophen or a combination of acetaminophen and Ibuprofen on the success of inferior alveolar nerve block for teeth with irreversible pulpitis. *J Endod* 2007;33:11-4.
- [24] Parirokh M, Ashouri R, Rekabi AR, Nakhaee N, Pardakhti A, Askarifard S, et al. The effect of premedication with ibuprofen and indomethacin on the success of inferior alveolar nerve block for teeth with irreversible pulpitis. *J Endod* 2010;36:1450-4.
- [25] Simpson M, Drum M, Nusstein J, Reader A, Beck M. Effect of combination of preoperative ibuprofen/acetaminophen on the success of the inferior alveolar nerve block in patients with symptomatic irreversible pulpitis. *J Endod* 2011;37:593-7.
- [26] Stanley W, Drum M, Nusstein J, Reader A, Beck M. Effect of nitrous oxide on the efficacy of the inferior alveolar nerve block in patients with symptomatic irreversible pulpitis. *J Endod* 2012;38:565-9.
- [27] Khabbaz MG, Anastasiadis PL, Sykaras SN. Determination of endotoxins in caries: association with pulpal pain. *Int Endod J* 2000;33:132-7.
- [28] Bernick S. Effect of aging on the nerve supply to human teeth. *J Dent Res* 1967;46:694-9.
- [29] Bernick S, Nedelman C. Effect of aging on the human pulp. *J Endod* 1975;1:88-94.

- [30] Ikawa M, Komatsu H, Ikawa K, Mayanagi H, Shimauchi H. Age-related changes in the human pulpal blood flow measured by laser Doppler flowmetry. Dent Traumatol 2003;19:36-40.
- [31] Morse DR, Esposito JV, Schoor RS. A radiographic study of aging changes of the dental pulp and dentin in normal teeth. Quintessence Int 1993;24:329-33.
- [32] Johnsen DC, Harshbarger J, Rymer HD. Quantitative assessment of neural development in human premolars. Anat Rec 1983;205:421-9.
- [33] Michaelson PL, Holland GR. Is pulpitis painful? Int Endod J 2002;35:829-32.
- [34] Meechan JG. Practical Dental Local Anesthesia. 2nd ed. Wilson NHF, editor. London: Quintessence Publishing Co.Ltd.; 2010.
- [35] Reader ANJ, Drum M. Successful local anesthesia for restorative dentistry and endodontics. 1st ed. USA: Quinessence Publishing Co. Inc.; 2011.
- [36] Meechan JG. How to overcome failed local anaesthesia. Br Dent J 1999;186:15-20.
- [37] Madan GA, Madan SG, Madan AD. Failure of inferior alveolar nerve block: exploring the alternatives. Journal of the American Dental Association (1939) 2002;133:843-6.
- [38] Ashkenazi M, Taubman L, Gavish A. Age-associated changes of the mandibular foramen position in anteroposterior dimension and of the mandibular angle in dry human mandibles. Anat Rec (Hoboken) 2011;294:1319-25.
- [39] Kanno CM, de Oliveira JA, Cannon M, Carvalho AA. The mandibular lingula's position in children as a reference to inferior alveolar nerve block. J Dent Child (Chic) 2005;72:56-60.

- [40] Aps JK. Number of accessory or nutrient canals in the human mandible. *Clin Oral Investig* 2014;18:671-6.
- [41] Blanton PL, Jeske AH. The key to profound local anesthesia: neuroanatomy. *J Am Dent Assoc* 2003;134:753-60.
- [42] Wilson S, Johns P, Fuller PM. The inferior alveolar and mylohyoid nerves: an anatomic study and relationship to local anesthesia of the anterior mandibular teeth. *J Am Dent Assoc* 1984;108:350-2.
- [43] Bennett S, Townsend G. Distribution of the mylohyoid nerve: anatomical variability and clinical implications. *Aust Endod J* 2001;27:109-11.
- [44] Vreeland DL, Reader A, Beck M, Meyers W, Weaver J. An evaluation of volumes and concentrations of lidocaine in human inferior alveolar nerve block. *J Endod* 1989;15:6-12.
- [45] Klingberg G, Raadal M, Arnrup K. Dental fear and behavior management problems. In: Koch G, Poulsen S, editors. *Pediatric Dentistry A Clinical Approach*. 2nd ed: Blackwell publishing Ltd.; 2009. p. 32-43.
- [46] van Wijk AJ, Makkes PC. Highly anxious dental patients report more pain during dental injections. *Br Dent J* 2008;205:E7; discussion 142-3.
- [47] Kuscu OO, Akyuz S. Is it the injection device or the anxiety experienced that causes pain during dental local anaesthesia? *Int J Paediatr Dent* 2008;18:139-45.
- [48] Nakai Y, Milgrom P, Mancl L, Coldwell SE, Domoto PK, Ramsay DS. Effectiveness of local anesthesia in pediatric dental practice. *J Am Dent Assoc* 2000;131:1699-705.
- [49] Okawa K, Ichinohe T, Kaneko Y. Anxiety may enhance pain during dental treatment. *Bull Tokyo Dent Coll* 2005;46:51-8.

- [50] Eli I, Svensson P. The multidimensional nature of pain. In: Bergenholz G, Horsted-Bindslev P, Reit C, editors. Textbook of Endodontontology. 2nd ed: Blackwell Publishing Ltd.; 2010. p. 277-89.
- [51] Bruno KF, Silva JA, Silva TA, Batista AC, Alencar AH, Estrela C. Characterization of inflammatory cell infiltrate in human dental pulpitis. Int Endod J 2010;43:1013-21.
- [52] Gibbs JL, Hargreaves KM. Mechanisms of odontogenic and non-odontogenic pain. In: Ingle JI, Bakland LK, Baumgartner JC, editors. Ingle's Endodontics. 6th ed: People's Medical Publishing House; 2008. p. 376-91.
- [53] Nusstein JM, Reader A, Drum M. Local anesthesia strategies for the patient with a "hot" tooth. Dent Clin North Am 2010;54:237-47.
- [54] Modaresi J, Mozayeni MA, Dianat O. Comparing the quality of anaesthesia in normal and inflamed teeth by pulp testing. Aust Endod J 2005;31:120-2.
- [55] Dreven LJ, Reader A, Beck M, Meyers WJ, Weaver J. An evaluation of an electric pulp tester as a measure of analgesia in human vital teeth. J Endod 1987;13:233-8.
- [56] Byers MR, Taylor PE, Khayat BG, Kimberly CL. Effects of injury and inflammation on pulpal and periapical nerves. J Endod 1990;16:78-84.
- [57] Woolf CJ. Central sensitization: implications for the diagnosis and treatment of pain. Pain 2011;152:S2-15.
- [58] Hahn CL, Liewehr FR. Relationships between caries bacteria, host responses, and clinical signs and symptoms of pulpitis. J Endod 2007;33:213-9.
- [59] Haase A, Reader A, Nusstein J, Beck M, Drum M. Comparing anesthetic efficacy of articaine versus lidocaine as a supplemental buccal infiltration of

the mandibular first molar after an inferior alveolar nerve block. J Am Dent Assoc 2008;139:1228-35.

- [60] Kanaa MD, Whitworth JM, Corbett IP, Meechan JG. Articaine and lidocaine mandibular buccal infiltration anesthesia: a prospective randomized double-blind cross-over study. J Endod 2006;32:296-8.
- [61] Robertson D, Nusstein J, Reader A, Beck M, McCartney M. The anesthetic efficacy of articaine in buccal infiltration of mandibular posterior teeth. J Am Dent Assoc 2007;138:1104-12.
- [62] Brandt RG, Anderson PF, McDonald NJ, Sohn W, Peters MC. The pulpal anesthetic efficacy of articaine versus lidocaine in dentistry: a meta-analysis. J Am Dent Assoc 2011;142:493-504.
- [63] Yapp KE, Hopcraft MS, Parashos P. Articaine: a review of the literature. Br Dent J 2011;210:323-9.
- [64] Wright GZ, Weinberger SJ, Friedman CS, Plotzke OB. The use of articaine local anesthesia in children under 4 years of age—a retrospective report. Anesth Prog 1989;36:268.
- [65] Leith R, Lynch K, O'Connell AC. Articaine use in children: a review. Eur Arch Paediatr Dent 2012;13:293-6.
- [66] Replogle K, Reader A, Nist R, Beck M, Weaver J, Meyers WJ. Anesthetic efficacy of the intraosseous injection of 2% lidocaine (1:100,000 epinephrine) and 3% mepivacaine in mandibular first molars. Oral Surg Oral Med Oral Pathol Oral Radiol Endod 1997;83:30-7.
- [67] Zarei M, Ghoddusi J, Sharifi E, Forghani M, Afkhami F, Marouzi P. Comparison of the anaesthetic efficacy of and heart rate changes after periodontal ligament or intraosseous X-Tip injection in mandibular molars: a randomized controlled clinical trial. Int Endod J 2012;45:921-6.

- [68] Aps JK. Intraosseous Local Anesthesia in Dentistry Makes Sense. *Int J Clin Anesthesiol* 2013;1.
- [69] Reisman D, Reader A, Nist R, Beck M, Weaver J. Anesthetic efficacy of the supplemental intraosseous injection of 3% mepivacaine in irreversible pulpitis. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 1997;84:676-82.
- [70] AAPD. Guideline on use of local anesthesia for pediatric dental patients. Reference manual 2009;34:183-9.
- [71] Endo T, Gabka J, Taubenheim L. Intraligamentary anesthesia: benefits and limitations. *Quintessence Int* 2008;39:e15-25.
- [72] Walton RE, Abbott BJ. Periodontal ligament injection: a clinical evaluation. *J Am Dent Assoc* 1981;103:571-5.
- [73] Smith GN, Walton RE. Periodontal ligament injection: distribution of injected solutions. *Oral Surg Oral Med Oral Pathol* 1983;55:232-8.
- [74] Boopathi T, Sebeena M, Sivakumar K, Harikaran J, Karthick K, Raj A. Supplemental pulpal anesthesia for mandibular teeth. *J Pharm Bioallied Sci* 2013;5:S103-8.
- [75] Malamed SF. The periodontal ligament (PDL) injection: an alternative to inferior alveolar nerve block. *Oral Surg Oral Med Oral Pathol* 1982;53:117-21.
- [76] Smith GN, Walton RE, Abbott BJ. Clinical evaluation of periodontal ligament anesthesia using a pressure syringe. *J Am Dent Assoc* 1983;107:953-6.
- [77] Malamed SF. Handbook of local anesthesia. 3rd ed. USA: Mosby-Year Book, Inc.; 1990.

- [78] Nusstein J, Burns Y, Reader A, Beck M, Weaver J. Injection pain and postinjection pain of the palatal-anterior superior alveolar injection, administered with the Wand Plus system, comparing 2% lidocaine with 1:100,000 epinephrine to 3% mepivacaine. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2004;97:164-72.
- [79] Brannstrom M, Lindskog S, Nordenvall KJ. Enamel hypoplasia in permanent teeth induced by periodontal ligament anesthesia of primary teeth. *J Am Dent Assoc* 1984;109:735-6.
- [80] Maki K, Miller A, Okano T, Shibasaki Y. Changes in Cortical Bone Mineralization in the Developing Mandible: A Three-Dimensional Quantitative Computed Tomography Study. *J Bone Miner Res* 2000;15:700-9.
- [81] Tranasi M, Sberna MT, Zizzari V, D'Apolito G, Mastrangelo F, Salini L, et al. Microarray evaluation of age-related changes in human dental pulp. *J Endod* 2009;35:1211-7.
- [82] Sixou JL, Barbosa-Rogier ME. Efficacy of intraosseous injections of anesthetic in children and adolescents. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2008;106:173-8.
- [83] Levin LG. Pulp and periradicular testing. *J Endod* 2013;39:S13-9.
- [84] Jafarzadeh H, Abbott PV. Review of pulp sensibility tests. Part I: general information and thermal tests. *Int Endod J* 2010;43:738-62.
- [85] Olgart L. Excitation of intradental sensory units by pharmacological agents. *Acta Physiol Scand* 1974;92:48-55.
- [86] Gopikrishna V, Pradeep G, Venkateshbabu N. Assessment of pulp vitality: a review. *Int J Paediatr Dent* 2009;19:3-15.

- [87] Trowbridge HO, Franks M, Korostoff E, Emling R. Sensory response to thermal stimulation in human teeth. *J Endod* 1980;6:405-12.
- [88] Jones DM. Effect of the type carrier used on the results of dichlorodifluoromethane application to teeth. *J Endod* 1999;25:692-4.
- [89] Jones VR, Rivera EM, Walton RE. Comparison of carbon dioxide versus refrigerant spray to determine pulpal responsiveness. *J Endod* 2002;28:531-3.
- [90] Fuss Z, Trowbridge H, Bender IB, Rickoff B, Sorin S. Assessment of reliability of electrical and thermal pulp testing agents. *J Endod* 1986;12:301-5.
- [91] Augsburger RA, Peters DD. In vitro effects of ice, skin refrigerant, and CO₂ snow on intrapulpal temperature. *J Endod* 1981;7:110-6.
- [92] Chen E, Abbott PV. Evaluation of accuracy, reliability, and repeatability of five dental pulp tests. *J Endod* 2011;37:1619-23.
- [93] Weisleder R, Yamauchi S, Caplan DJ, Trope M, Teixeira FB. The validity of pulp testing: a clinical study. *J Am Dent Assoc* 2009;140:1013-7.
- [94] Grossman LI. Endodontic practice. 10th ed: Philadelphia: Lea&Febiger; 1981.
- [95] Kulild JC. Diagnostic testing. In: Ingle JI, Bakland LK, Baumgartner JC, editors. *Ingle's Endodontics*. USA: People's Medical Publishing House; 2008. p. 532-53.
- [96] Mumford J. Evaluation of gutta-percha and ethyl chloride in pulp testing. *Br Dent J* 1964;116:338-42.

- [97] Pantera EA, Jr., Anderson RW, Pantera CT. Reliability of electric pulp testing after pulpal testing with dichlorodifluoromethane. *J Endod* 1993;19:312-4.
- [98] Narhi M, Virtanen A, Kuhta J, Huopaniemi T. Electrical stimulation of teeth with a pulp tester in the cat. *Scand J Dent Res* 1979;87:32-8.
- [99] Klein H. Pulp responses to an electric pulp stimulator in the developing permanent anterior dentition. *ASDC J Dent Child* 1978;45:199-202.
- [100] Bernick S. Differences in Nerve Distribution between Erupted and Non-Erupted Human Teeth. *J Dent Res* 1964;43:406-11.
- [101] Ehrmann EH. Pulp testers and pulp testing with particular reference to the use of dry ice. *Aust Dent J* 1977;22:272-9.
- [102] Fulling HJ, Andreasen JO. Influence of maturation status and tooth type of permanent teeth upon electrometric and thermal pulp testing. *Scand J Dent Res* 1976;84:286-90.
- [103] Bender IB, Landau MA, Fonsecca S, Trowbridge HO. The optimum placement-site of the electrode in electric pulp testing of the 12 anterior teeth. *J Am Dent Assoc* 1989;118:305-10.
- [104] Cooley RL, Robison SF. Variables associated with electric pulp testing. *Oral Surg Oral Med Oral Pathol* 1980;50:66-73.
- [105] Harris SC. Problems of experimental algesimetry and a comment on the tooth pulp method. *J Chronic Dis* 1956;4:52-8.
- [106] Certosimo A, Archer R. A clinical evaluation of the electric pulp tester as an indicator of local anesthesia. *Oper Dent* 1995;21:25-30.

- [107] Petersson K, Soderstrom C, Kiani-Anaraki M, Levy G. Evaluation of the ability of thermal and electrical tests to register pulp vitality. *Endod Dent Traumatol* 1999;15:127-31.
- [108] Al-Namankany A, De Souza M, Ashley P. Evidence-based dentistry: analysis of dental anxiety scales for children. *Br Dent J* 2012;212:219-22.
- [109] Buchanan H, Niven N. Validation of a Facial Image Scale to assess child dental anxiety. *Int J Paediatr Dent* 2002;12:47-52.
- [110] Tomlinson D, von Baeyer CL, Stinson JN, Sung L. A systematic review of faces scales for the self-report of pain intensity in children. *Pediatrics* 2010;126:e1168-98.
- [111] Chambers CT, Hardial J, Craig KD, Montgomery C. Faces scales for the measurement of postoperative pain intensity in children following minor surgery. *Clin J Pain* 2005;21:277-85.
- [112] Chambers CT, Giesbrecht K, Craig KD, Bennett SM, Huntsman E. A comparison of faces scales for the measurement of pediatric pain: children's and parents' ratings. *Pain* 1999;83:25-35.
- [113] Luffy R, Grove SK. Examining the validity, reliability, and preference of three pediatric pain measurement tools in African-American children. *Pediatr Nurs* 2003;29:54-9.
- [114] Chambers CT, Craig KD. An intrusive impact of anchors in children's faces pain scales. *Pain* 1998;78:27-37.
- [115] Champion GD. Comment on Stanford EA et al.: The role of developmental factors in predicting young children's use of a self-report scale for pain. *Pain* 2006;120:16-23. *Pain* 2006;124:360-1; author reply 1-2.
- [116] von Baeyer CL. Children's self-reports of pain intensity: scale selection, limitations and interpretation. *Pain Res Manag* 2006;11:157-62.

- [117] Demirjian A, Goldstein H, Tanner JM. A new system of dental age assessment. *Hum Biol* 1973;45:211-27.
- [118] Lin J, Chandler NP. Electric pulp testing: a review. *Int Endod J* 2008;41:365-74.
- [119] AAE Consensus Conference Recommended Diagnostic Terminology. *J Endod* 2009;35:1634.
- [120] Rodd HD, Boissonade FM. Innervation of human tooth pulp in relation to caries and dentition type. *J Dent Res* 2001;80:389-93.
- [121] Rodd HD, Boissonade FM. Immunocytochemical investigation of immune cells within human primary and permanent tooth pulp. *Int J Paediatr Dent* 2006;16:2-9.
- [122] Azize PM, Humphreys A, Cattani A. The impact of language on the expression and assessment of pain in children. *Intensive Crit Care Nurs* 2011;27:235-43.
- [123] Zero DT, Zandona AF, Vail MM, Spolnik KJ. Dental caries and pulpal disease. *Dent Clin North Am* 2011;55:29-46.
- [124] Singh SK, Kanaparthi A, Kanaparthi R, Pillai A, Sandhu G. Geriatric Endodontic. *Journal of Orofacial Research* 2013;3:191-6.
- [125] Jespersen JJ, Hellstein J, Williamson A, Johnson WT, Qian F. Evaluation of dental pulp sensibility tests in a clinical setting. *J Endod* 2014;40:351-4.
- [126] Tortamano IP, Siviero M, Lee S, Sampaio RM, Simone JL, Rocha RG. Onset and duration period of pulpal anesthesia of articaine and lidocaine in inferior alveolar nerve block. *Braz Dent J* 2013;24:371-4.
- [127] Asfour MA, Millar BJ, Smith PB. An assessment of the reliability of pulp testing deciduous teeth. *Int J Paediatr Dent* 1996;6:163-6.

[128] Pitt Ford TR, Patel S. Technical equipment for assessment of dental pulp status. Endod Topics 2004;7:2-13.



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved