

## REFERENCES

1. Hoy DG, Protani M, De R, Buchbinder R. The epidemiology of neck pain. Best Prac Res Clin Rheumotol 2010;24:783-92.
2. Chiu TT, Ku WY, Lee MH, Sum WK, Wan MP, Wong CY, et al. A study on the prevalence of and risk factors for neck pain among university academic staff in Hong Kong. J Occup Rehabil. 2002;12:77-91.
3. Aoyagia K, Rossb PD, Huangb C, Wasnichb RD, Hayashic T, Ti T. Prevalence of joint pain is higher among women in rural Japan than urban Japanese-American women in Hawaii. Ann Rheum Dis. 1999;58:315-9.
4. Leboeuf-Yde C, Nielsen J, Kyvik KO, Fejer R, Hartvigsen J. Pain in the lumbar, thoracic or cervical regions: do age and gender matter? a population-based study of 34,902 Danish twins 20-71 years of age. BMC Musculoskelet Disord. 2009;10:39. doi:10.1186/1471-2474-10-39.
5. Skillgate E, Magnusson C, Lundberg M, Hallqvist J. The age- and sex-specific occurrence of bothersome neck pain in the general population - results from the Stockholm public health cohort. BMC Musculoskeletal Disord. 2012;13:185. doi:10.1186/1471-2474-13-185.
6. Kanchanomai S, Janwantanakul P, Pensri P, Jiamjarasrangsi W. Risk factors for the onset and persistence of neck pain in undergraduate students: 1-year prospective cohort study. BMC public health. 2011;11:566. doi:10.1186/1471-2458-11-566.
7. Tuzun EH. Quality of life in chronic musculoskeletal pain. Best Prac Clin Rheumotol. 2007;21:567-79.
8. Daffner SD, Hilibrand AS, Hanscom BS . Impact of neck and arm pain on overall health status. Spine. 2003;28:2030-5.

9. Cagnie B, Danneels L, Van Tiggelen D, De Loose V, Cambier D. Individual and work related risk factors for neck pain among office workers: a cross sectional study. *Eur Spine J.* 2007;16:679-86.
10. Zakharova-Luneva E, Jull G, Johnston V, O'Leary S. Altered trapezius muscle behavior in individuals with neck pain and clinical signs of scapular dysfunction. *J Manipulative Physiol Ther.* 2012;35:346-53.
11. Falla D, Jull G, Hodges P. Patients with neck pain demonstrate reduced electromyographic activity of the deep cervical flexor muscles during performance of the craniocervical flexion test. *Spine.* 2004;29:2108-14.
12. Farina D, Arendt-Nielsen L, Merletti R, Graven-Nielsen T. Effect of experimental muscle pain on motor unit firing rate and conduction velocity. *J Neurophysiol.* 2004;91:1250-9.
13. Falla D, Farina D. Muscle fiber conduction velocity of the upper trapezius muscle during dynamic contraction of the upper limb in patients with chronic neck pain. *Pain.* 2005;116:138-45.
14. Falla D, Arendt-Nielsen L, Farina D. The pain-induced change in relative activation of upper trapezius muscle regions is independent of the site of noxious stimulation. *Clin Neurophysiol.* 2009;120:150-7.
15. Wegner S, Jull G, O'Leary S, Johnston V. The effect of a scapular postural correction strategy on trapezius activity in patients with neck pain. *Man Ther.* 2010;15:562-6.
16. Petersen SM, Wyatt SN. Lower trapezius muscle strength in individuals with unilateral neck pain. *J Orthop Sports Phys Ther.* 2011;41:260-5.
17. Johnson G, Bogduk N, Noweitzke A, House D. Anatomy and actions of the trapezius muscle. *Clin Biomech.* 1994;9:44-50.

18. Kibler WB. The role of the scapula in athletic shoulder function. Am J Sports Med. 1998;26:325-37.
19. Paine RM, Voight M. The role of the scapula. J Orthop Sports Phys Ther. 1993;18:386-91.
20. Phadke V, Camargo P, Ludewig P. Scapular and rotator cuff muscle activity during arm elevation: a review of normal function and alterations with shoulder impingement. Res Bras Fisioter. 2009;13:1-9.
21. Shi J, Zheng YP, Chen X, Huang QH. Assessment of muscle fatigue using sonomyography: muscle thickness change detected from ultrasound images. Med Eng Phys. 2007;29:472-9.
22. Rezasoltani A, Ahmadipoor A, Khademi-Kalantari K, Javanshir K. The sign of unilateral neck semispinalis capitis muscle atrophy in patients with chronic non-specific neck pain. J Back Musculoskelet Rehabil. 2012;25: 67-72.
23. Kristjansson E. Reliability of ultrasonography for the cervical multifidus muscle in asymptomatic and symptomatic subjects. Man Ther. 2004;9:83-8.
24. Rezasoltani A, Ali-Reza A, Khosro K-K, Abbass R. Preliminary study of neck muscle size and strength measurements in females with chronic non-specific neck pain and healthy control subjects. Man Ther. 2010;15:400-3.
25. Muñoz-Muñoz S, Muñoz-García MT, Alburquerque-Sendín F, Arroyo-Morales M, Fernández-de-las-Peñas C. Myofascial trigger points, pain, disability, and sleep quality in individuals with mechanical neck pain. J Manipulative Physiol Ther. 2012;35:608-13.
26. Ståhl M, Mikkelsson M, Kautiainen H, Häkkinen A, Ylinen J, Salminen JJ. Neck pain in adolescence. a 4-year follow-up of pain-free preadolescents. Pain. 2004;110:427-31.

27. Sjors A, Larsson B, Persson AL, Gerdle B. An increased response to experimental muscle pain is related to psychological status in women with chronic non-traumatic neck-shoulder pain. *BMC Musculoskelet Disord.* 2011;12:230. doi:10.1186/1471-2474-12-230.
28. Delcker A, Walker F, Caress J, Hunt C, Tegeler C. In vitro measurement of muscle volume with 3-dimensional ultrasound. *Eur J Ultrasound.* 1999;9: 185-90.
29. Javanshir K, Mohseni-Bandpei MA, Rezasoltani A, Amiri M, Rahgozar M. Ultrasonography of longus colli muscle: a reliability study on healthy subjects and patients with chronic neck pain. *J Bodyw Mov Ther.* 2011;15:50-6.
30. Talbott NR, Witt DW. Ultrasound imaging of the serratus anterior muscle at rest and during contraction. *Clin Physiol Funct Imaging.* 2012;1-9.
31. O'Sullivan C, Bentman S, Bennett K, Stokes M. Rehabilitative ultrasound imaging of the lower trapezius muscle: technical description and reliability. *J Orthop Sports Phys Ther.* 2007;37:620-6.
32. O'Sullivan C, Meaney J, Boyle G, Gormley J, Stokes M. The validity of rehabilitative ultrasound imaging for measurement of trapezius muscle thickness. *Man Ther.* 2009;14:572-8.
33. Cagnie B, Derese E, Vandamme L, Verstraete K, Cambier D, Danneels L. Validity and reliability of ultrasonography for the longus colli in asymptomatic subjects. *Man Ther.* 2009;14:421-6.
34. Fernandez-de-Las-Penas C, Hernandez-Barrera V, Alonso-Blanco C, Palacios-Cena D, Carrasco-Garrido P, Jimenez-Sanchez S, et al. Prevalence of neck and low back pain in community-dwelling adults in spain: a population-based national study. *Spine.* 2011;36:213-9.

35. Fejer R, Kyvik KO, Hartvigsen J. The prevalence of neck pain in the world population: a systematic critical review of the literature. *Eur Spine J.* 2006;15:834-43.
36. Côté P, Cassidy JD, Carroll L. The saskatchewan health and back pain survey: the prevalence of neck pain and related disability in saskatchewan adults. *Spine.* 1998;23:1689-98.
37. Young A, Stokes M, Crowe M. Size and strength of the quadriceps muscles of old and young women. *Eur J Clin Invest.* 1984;14:282-7.
38. Guzman J, Haldeman S, Carroll LJ, Carragee EJ, Hurwitz EL, Peloso P, et al. Clinical practice implications of the bone and joint decade 2000–2010 task force on neck pain and its associated disorders: from concepts and findings to recommendations. *J Manipulative Physiol Ther.* 2009;32:227-43.
39. Ariens GA, van Mechelen W, Bongers PM, Bouter LM, van der Wal G. Physical risk factors for neck pain. *Scand J Work Environ Health.* 2000;26:7-19.
40. Côté P, Velde G, David Cassidy J, Carroll L, Hogg-Johnson S, Holm L, et al. The burden and determinants of neck pain in workers. *Eur Spine J.* 2008;17:60-74.
41. Hogg-Johnson S, van der Velde G, Carroll LJ, Holm LW, Cassidy JD, Guzman J, et al. The burden and determinants of neck pain in the general population: results of the bone and joint decade 2000–2010 task force on neck pain and its associated disorders. *J Manipulative Physiol Ther.* 2009;32:46-60.
42. Cagnie B, Dirks R, Schouten M, Parlevliet T, Cambier D, Danneels L. Functional reorganization of cervical flexor activity because of induced muscle pain evaluated by muscle functional magnetic resonance imaging. *Man Ther.* 2011;16:470-5.

43. Jesus-Moraleida FR, Ferreira PH, Pereira LSM, Vasconcelos CM, Ferreira ML. Ultrasonographic analysis of the neck flexor muscles in patients with chronic neck pain and changes after cervical spine mobilization. *J Manipulative Physiol Ther.* 2011;34:514-24.
44. Saavedra-Hernandez M, Castro-Sanchez AM, Cuesta-Vargas AI, Cleland JA, Fernandez-de-las-Penas C, Arroyo-Morales M. The contribution of previous episodes of pain, pain intensity, physical impairment, and pain-related fear to disability in patients with chronic mechanical neck pain. *Am J Phys Med Rehabil.* 2012;91:1070-6.
45. Park KN, Kwon OY, Ha SM, Kim SJ, Choi HJ, Weon JH. Comparison of electromyographic activity and range of neck motion in violin students with and without neck pain during playing. *Med Probl Perform Art.* 2012;27:188-92.
46. Guo LY, Lee SY, Lin CF, Yang CH, Hou YY, Wu WL, et al. Three-dimensional characteristics of neck movements in subjects with mechanical neck disorder. *J Back Musculoskelet.* 2012;25:47-53.
47. Jull G, Bogduk N, Marsland A. The accuracy of manual diagnosis for cervical zygapophysial joint pain syndromes. *Med J Aust.* 1988;148:233-6.
48. Weber Hellstenius SA. Recurrent neck pain and headaches in preadolescents associated with mechanical dysfunction of the cervical spine: a cross-sectional observational study with 131 students. *J Manipulative Physiol Ther.* 2009;32:625-34.
49. Lord SM, Barnsley L, Wallis BJ, Bogduk N. Chronic cervical zygapophysial joint pain after whiplash: a placebo-controlled prevalence study. *Spine.* 1996;21:1737-44.
50. O'Leary S, Vicenzino B, Jull G. A new method of isometric dynamometry for the craniocervical flexor muscles. *Phys Ther.* 2005;85:556-64.

51. O'Leary S, Jull G, Kim M, Vicenzino B. Cranio-cervical flexor muscle impairment at maximal, moderate, and low loads is a feature of neck pain. *Man Ther.* 2007;12:34-9.
52. Falla D, Jull G, Rainoldi A, Merletti R. Neck flexor muscle fatigue is side specific in patients with unilateral neck pain. *Eur J Pain.* 2004;8:71-7.
53. Falla D, Rainoldi A, Jull G, Stavrou G, Tsao H. Lack of correlation between sternocleidomastoid and scalene muscle fatigability and duration of symptoms in chronic neck pain patients. *Neurophysiol Clin.* 2004;34: 159-65.
54. Jull G, O'Leary S, Falla D. Clinical assessment of the deep cervical flexor muscles: the craniocervical flexion test. *J Manipulative Physiol Ther.* 2008;31:525-33.
55. Field S, Treleaven J, Jull G. Standing balance: a comparison between idiopathic and whiplash-induced neck pain. *Man Ther.* 2008;13:183-91.
56. Hurwitz EL, Carragee EJ, van der Velde G, Carroll LJ, Nordin M, Guzman J, et al. Treatment of neck pain: noninvasive interventions: results of the bone and joint decade 2000–2010 task force on neck pain and its associated disorders. *J Manipulative Physiol Ther.* 2009;32:141-75.
57. Carroll LJ, Cassidy JD, Peloso PM, Giles-Smith L, Cheng CS, Greenhalgh SW, et al. Methods for the best evidence synthesis on neck pain and its associated disorders: the bone and joint decade 2000-2010 task force on neck pain and its associated disorders. *J Manipulative Physiol Ther.* 2009;32:39-45.
58. Merskey H, Bogduk N, editors. Classification of chronic pain. Descriptions of chronic pain syndromes and definitions of pain terms. 2<sup>nd</sup> ed. Seattle: IASP Press; 1994:11-6.

59. Guzman J, Hurwitz EL, Carroll LJ, Haldeman S, Côté P, Carragee EJ, et al. A new conceptual model of neck pain: linking onset, course, and care: the bone and joint decade 2000–2010 task force on neck pain and its associated disorders. *J Manipulative Physiol Ther.* 2009;32:17-28.
60. Gross A, Miller J, D'Sylva J, et al. Manipulation or mobilisation for neck pain: a cochrane review. *Man Ther.* 2010;15:315-33.
61. Spitzer WO, Skovron ML, Salmi LR, Cassidy JD, Duranceau J, Suissa S, et al. Scientific monograph of the quebec task force on whiplash-associated disorders: redefining "whiplash" and its management. *Spine.* 1995;20: 1-73.
62. Dorshimer GW, Kelly M. Cervical pain in the athlete: common conditions and treatment. *Pri care.* 2005;32:231-43.
63. Bongers PM, Ijmker S, van den Heuvel S, Blatter BM. Epidemiology of work related neck and upper limb problems: psychosocial and personal risk factors (part I) and effective interventions from a bio behavioural perspective (part II). *J Occup Rehabil.* 2006;16:279-302.
64. Carroll LJ, Hogg-Johnson S, van der Velde G, Haldeman S, Holm LW, Carragee EJ, et al. Course and prognostic factors for neck pain in the general population: results of the bone and joint decade 2000-2010 task force on neck pain and its associated disorders. *J Manipulative Physiol Ther.* 2009;32:87-96.
65. Guzman J, Haldeman S, Carroll LJ, Carragee EJ, Hurwitz EL, Peloso P, et al. Clinical practice implications of the bone and joint decade 2000-2010 task force on neck pain and its associated disorders: from concepts and findings to recommendations. *Spine.* 2008;33:199-213.
66. Schünke M, Schulte E, Ross LM, Schumacher U, Lamperti ED. Thieme atlas of anatomy: general anatomy and musculoskeletal system: New York: Thieme Medical Publishers; 2006.

67. Kendall FP, Kendall-McCreary E, Provance PG, McIntrye-Rodgers M, Romani WA. Muscles: testing and function with posture and pain. 5<sup>th</sup> ed. Baltimore: Lippincott Williams & Wilkins; 2005.
68. Mottram SL. Dynamic stability of the scapula. *Man Ther*. 1997;2:123-31.
69. Ludewig PM, Cook TM. Alterations in shoulder kinematics and associated muscle activity in people with symptoms of shoulder impingement. *Phys Ther*. 2000;80:276-91.
70. Ekstrom RA, Donatelli RA, Soderberg GL. Surface electromyographic analysis of exercise for the trapezius and serratus anterior muscles. *J Orthop Sports Phys Ther*. 2003;33:247-58.
71. McCabe RA, Orishimo KF, McHugh MP, Nicholas SJ. Surface electromyographic analysis of the lower trapezius muscle during exercises performed below ninety degrees of shoulder elevation in healthy subjects. *N Am J Sports Phys Ther*. 2007;2:34-43.
72. Kinney E, Wusthoff J, Zyck A, Hatzel B, Vaughn D, Strickler T. Activation of the trapezius muscle during varied forms of kendall exercises. *Phys Ther Sport*. 2008;9:3-8.
73. Yoshizaki K, Hamada J, Tamai K, Sahara R, Fujiwara T, Fujimoto T. Analysis of the scapulohumeral rhythm and electromyography of the shoulder muscles during elevation and lowering: comparison of dominant and nondominant shoulders. *J Shoulder Elbow Surg*. 2009;18:756-63.
74. Lukasiewicz AC, McClure P, Michener L, Pratt N, Sennett B. Comparison of 3-dimensional scapular position and orientation between subjects with and without shoulder impingement. *J Orthop Sport Phys Ther*. 1999;29: 574-86.

75. Pontillo M, Orishimo KF, Kremenic IJ, McHugh MP, Mullaney MJ, Tyler TF. Shoulder musculature activity and stabilization during upper extremity weight-bearing activities. *N Am J Sports Phys Ther.* 2007;2:90-6.
76. Westgaard RH, Vasseljen O, Holte KA. Trapezius muscle activity as a risk indicator for shoulder and neck pain in female service workers with low biomechanical exposure. *Ergonomics.* 2001;44:339-53.
77. Helgadottir H, Kristjansson E, Einarsson E, Karduna A, Jonsson H. Altered activity of the serratus anterior during unilateral arm elevation in patients with cervical disorders. *J Electromyogr Kinesiol.* 2011;21:947-53.
78. Weon JH, Oh JS, Cynn HS, Kim YW, Kwon OY, Yi CH. Influence of forward head posture on scapular upward rotators during isometric shoulder flexion. *J Bodyw Mov Ther.* 2010;14:367-74.
79. Gibbs V, Cole D, Sassano A. Ultrasound physics and technology: how, why and when. China: Churchill Livingstone; 2009.
80. Hides JA, Stokes MJ, Saide M, Jull GA, Cooper DH. Evidence of lumbar multifidus muscle wasting ipsilateral to symptoms in patients with acute subacute low back pain. *Spine* 1994;19:165-72.
81. Hides JA, Richardson C, Jull GA, Davies S. Ultrasound imaging in rehabilitation. *Aust J Physiother.* 1995;41:187-93.
82. Whittaker JL, Stokes M. Ultrasound imaging and muscle function. *J Orthop Sports Phys Ther.* 2011;41:572-80.
83. Kremkau FW. Diagnostic ultrasound: principles and instruments. Winston-Salem: W.B. Saunders Company; 2006.
84. O'Neill J. Musculoskeletal ultrasound: anatomy and technique. New York: Springer; 2008.

85. Thoirs K, English C. Ultrasound measures of muscle thickness: intra-examiner reliability and influence of body position. *Clin Physiol Funct Imaging*. 2009;29:440-6.
86. Bentman S, O'Sullivan C, Stokes M. Thickness of the middle trapezius muscle measured by rehabilitative ultrasound imaging: description of the technique and reliability study. *Clin Physiol Funct Imaging*. 2010;30: 426-31.
87. Koppenhaver SL, Hebert JJ, Fritz JM, Parent EC, Teyhen DS, Magel JS. Reliability of rehabilitative ultrasound imaging of the transversus abdominis and lumbar multifidus muscles. *Arch Phys Med Rehabil*. 2009;90:87-94.
88. Dupont AC, Sauerbrei EE, Fenton PV, Shragge PC, Loeb GE, Richmond FJ. Real time sonography to estimate muscle thickness: comparison with MRI and CT. *J Clin Ultrasound*. 2001;29:230-6.
89. Teyhen DS, Rieger JL, Westrick BR, Miller AC, Molly JM, Childs JD. Changes in deep abdominal muscle thickness during common trunk-strengthening exercises using ultrasound imaging. *J Orthop Sports Phys Ther*. 2008;38:596-605.
90. Whittaker JL, Warner MB, Stokes MJ. Induced transducer orientation during ultrasound imaging: effects on abdominal muscle thickness and bladder position. *Ultrasound Med Biol*. 2009;35:1803-11.
91. Uppot RN, Sahani DV, Hahn PF, Gervais D, Mueller PR. Impact of obesity on medical imaging and image-guided intervention. *Am J Roentgenol*. 2007;188:433-40.
92. Koo TK, Wong C, Zheng Y. Reliability of sonomyography for pectoralis major thickness measurement. *J Manipulative Physiol Ther*. 2010;33: 386-94.

93. Karen M, Sinead E. Ultrasound measurement of the size of the anterior tibial muscle group: the effect of exercise and leg dominance. *Sports Med Arthrosc Rehabil Ther Technol.* 2011;3:18. doi:10.1186/1758-2555-3-18.
94. Barber L, Barrett R, Lichtwark G. Validity and reliability of a simple ultrasound approach to measure medial gastrocnemius muscle length. *J Anat.* 2011;218:637-42.
95. Teyhen DS, George SZ, Dugan JL, Williamson J, Neilson BD, Childs JD. Inter-rater reliability of ultrasound imaging of the trunk musculature among novice raters. *J Ultrasound Med.* 2011;30:347-56.
96. Roscoe JT. Fundamental research statistics for the behavioural sciences. 2<sup>nd</sup> ed. New York: Holt Rinehart & Winston; 1975:211-4.
97. Boonstra AM, Schiphorst Preuper HR, Reneman MF, Posthumus JB, Stewart RE. Reliability and validity of the visual analogue scale for disability in patients with chronic musculoskeletal pain. *Int J Rehabil Res.* 2008;31:165-9.
98. Vernon H, Mior S. The Neck Disability Index: a study of reliability and validity. *J Manipulative Physiol Ther.* 1991;14:409-15.
99. Uthaikhup S, Paungmali A, Pirunsan U. Validation of Thai versions of the neck disability index and neck pain and disability scale in patients with neck pain. *Spine.* 2011;36:1415-21.
100. O'Sullivan C, McCarthy Persson U, Blake C, Stokes M. Rehabilitative ultrasound measurement of trapezius muscle contractile states in people with mild shoulder pain. *Man Ther.* 2012;17:139-44.
101. Portney LG, Watkins MP. Foundations of clinical research applications to practice. New Jersey: Prentice Hall Inc; 2000:560-7.

102. Danneels LA, Vanderstraeten GG, Cambier DC, Witvrouw EE, De Cuyper HJ. CT imaging of trunk muscles in chronic low back pain patients and healthy control subjects. *Eur Spine J.* 2000;9:266-72.
103. Hides J, Gilmore C, Stanton W, Bohlscheid E. Multifidus size and symmetry among chronic LBP and healthy asymptomatic subjects. *Man Ther.* 2008;13: 43-9.
104. Barker KL, Shamley DR, Jackson D. Changes in the cross-sectional area of multifidus and psoas in patients with unilateral back pain: the relationship to pain and disability. *Spine.* 2004;29:515-9.
105. McCreesh K, Egan S. Ultrasound measurement of the size of the anterior tibial muscle group: the effect of exercise and leg dominance. *Sports Med Arthrosc Rehabil Ther Technol.* 2011;3:18-23.
106. Malanga GA, Chu SK, Ramirez Del Toro J, Karnaugh RD, Dentico R, Komaroff E. Sonographic evaluation of supraspinatus cross-sectional area in collegiate baseball players. *PM R.* 2012;4:488-92.
107. Pang BS, Ying M. Sonographic measurement of achilles tendons in asymptomatic subjects: variation with age, body height, and dominance of ankle. *J Ultrasound Med.* 2006;25:1291-6.
108. Springer BA, Mielcarek BJ, Nesfield TK, Teyhen DS. Relationships among lateral abdominal muscles, gender, body mass index, and hand dominance. *J Orthop Sports Phys Ther.* 2006;36:289-97.
109. Lund JP, Donga R, Widmer CG, Stohler CS. The pain-adaptation model: a discussion of the relationship between chronic musculoskeletal pain and motor activity. *Can J Physiol Pharmacol.* 1991;69:683-94.
110. Hodges PW, Tucker K. Moving differently in pain: a new theory to explain the adaptation to pain. *Pain.* 2011;152:S90-8.

111. Janda V. Muscles and motor control in cervicogenic disorder: assessment and management. In: Grant R, editor. Physical therapy of the cervical and thoracic spine. 2<sup>nd</sup> ed. New York: Churchill Livingstone; 1994. p. 195-215.
112. Cools AM, Declercq GA, Cambier DC, Mahieu NN, Witvrouw EE. Trapezius activity and intramuscular balance during isokinetic exercise in overhead athletes with impingement symptoms. *Scand J Med Sci Sports*. 2007;17:25-33.
113. Constand M, MacDermid J. Effects of neck pain on reaching overhead and reading: a case-control study of long and short neck flexion. *BMC Sports Sci Med Rehabil*. 2013;5:1-6.
114. Lindstroem R, Graven-Nielsen T, Falla D. Current pain and fear of pain contribute to reduced maximum voluntary contraction of neck muscles in patients with chronic neck pain. *Arch Phys Med Rehabil*. 2012;93:2042-8.
115. Falla D, Bilenkij G, Jull G. Patients with chronic neck pain demonstrate altered patterns of muscle activation during performance of a functional upper limb task. *Spine*. 2004;29:1436-40.
116. Sterling M, Jull G, Wright A. The effect of musculoskeletal pain on motor activity and control. *J Pain*. 2001;2:135-45.
117. Javanshir K, Rezasoltani A, Mohseni-Bandpei MA, Amiri M, Ortega-Santiago R, Fernandez-de-Las-Penas C. Ultrasound assessment of bilateral longus colli muscles in subjects with chronic bilateral neck pain. *Am J Phys Med Rehabil*. 2011;90:293-301.
118. Oatis CA. Kinesiology: the mechanics and pathomechanics of human movement. Philadelphia: Lippincott Williams and Wilkins; 2004.
119. Faulkner JA, Larkin LM, Claflin DR, Brooks SV. Age-related changes in the structure and function of skeletal muscles. *Clin Exp Pharmacol Physiol*. 2007;34:1091-6.